

Association of Posterior reversible encephalopathy syndrome with Renal failure.

Abstract

Background:

Posterior reversible encephalopathy syndrome is a clinicoradiological entity that is characterized by variable associations of seizure activity, consciousness impairment, headaches, visual abnormalities, nausea, vomiting and focal neurological signs. No large data exists on the association of Posterior reversible encephalopathy syndrome with renal failure.

Material and Methods:

This case series of five patients was collected to know the association of two conditions and evaluate outcome. All these patients were enrolled in a tertiary care hospital over a period of two years. . No informed consent was sought because the study was merely observational and did not demand deviations from standard.

Results:

Four patients presented with acute renal failure and one patient had established End stage renal disease. All the Patients had Hypertention at presentation besides other risk factors. All the patients had Magnetic Resonance imaging documented posterior reversible encephalopathy syndrome. Despite intensive management, two patients did not show any compromise in the control of blood pressure and died. This observed showed that association of renal failure increases mortality in patients with Posterior reversible encephalopathy syndrome. However, our series is limited by small number of patients and hence no definite conclusions can be drawn from this observation.

Conclusion:

Posterior reversible encephalopathy syndrome should be considered in any patient of Renal failure with neurological manifestations and aggressive management is warranted.

22

23 **Key-words:** Posterior reversible encephalopathy syndrome; Magnetic Resonance Imaging; Renal failure.

24 **Introduction**

25 Posterior Reversible Encephalopathy Syndrome (PRES), characterized by transient neurological symptoms, including headache, altered mental
26 status, seizures and visual impairment, is a well-known condition occurring in some patients hospitalised for acute illness.(1) Neuroradiological
27 study usually shows oedema involving the cerebral posterior regions (1–4). The most common causes of PRES are hypertensive encephalopathy,
28 preeclampsia, eclampsia, cyclosporine A (CSA) neurotoxicity (5,6 4–6). The distinctiveness of PRES is its reversibility both in terms of the clinical
29 and radiological abnormalities after institution of appropriate treatment and removal of the precipitating factors. Although PRES is reversible once
30 treatment is instituted, delayed diagnosis and therapy can result in permanent brain damage and neurological sequelae, such as chronic epilepsy
31 (1, 2). Most of the literature available on PRES is from case reports and case series. PRES in patients with renal failure is not commonly described
32 in literature. We could not find any large series describing the condition in patients with renal failure. This case series of 5 patients was collected to
33 evaluate the course of disease and evaluate outcome in patients with renal failure.

34

35 **Material and Methods:**

36 All the patients were enrolled in a tertiary care hospital over a period of two years. Patients were taken from emergency department, intensive
37 care unit (ICU) and specialty wards. Patient data was collected at the time of admission and during hospital stay. The data collected include
38 patient demographics, clinical characteristics, laboratory parameters which include total leukocyte count, platelet count, kidney function tests,

39 urine examination and cultures. The clinical presentation and laboratory evaluation of the patients is shown in Table 1. Posterior Reversible
40 Encephalopathy Syndrome (PRES) was diagnosed by Magnetic Resonance Imaging (MRI) of Brain.

41 Diagnostic workup and therapy was at the discretion of attending doctor. No informed consent was sought because the study was merely
42 observational and did not demand deviations from standard clinical care. The outcome was defined by clinical and radiological improvement or
43 death.

44 **Clinical Presentation:**

45 All the patients presented with headache, altered sensorium and hypertention. The hypertention in four patients was first time detected and
46 one patient with known ESRD was on suboptimal doses of antihypertensives with poor compliance Case no 1 was a 34 year old female having
47 ESRD and was on dialysis protocol. She was on suboptimal doses of antihypertensives with uncontrolled blood pressure record. She presented
48 with accelerated hypertention with clinical presentation shown in the table 1. MRI of the brain showed hyperintense lesions in parieto-occipital
49 regions and was suggestive of PRES (Figure 1).

50 Case no. 2 was a 14 year old boy who presented with GTCS and unconsciousness. Patient had high blood pressure, mild renal failure, and active
51 urinary sediment. Patient gave history of upper respiratory tract infections two weeks before. Imaging studies showed hyperintense lesions in
52 Parieto-occipital and cerebellar regions (Figure 2). The clinical presentation, biochemical profile and outcome of other patients is shown in table
53 1.

54 **Laboratory evaluation:**

55 All the patients had renal failure. Two patients had Total leukocyte count in the upper range of normal and two patients had leukocytosis. All
56 the the patients had MRI documented PRES. Follow up MRI in three patients was normal.

57 **Management and outcome:**

58 All the patients were managed in ICU settings. Intensive management was pursued for control of hypertention which included Nitroglycerine
59 infusion, Intravenous (I.V) Labetolol (SOS) and oral antihypertensive agents. Four patients (case no 1, 3, 4 and 5) were empirically started on
60 antibiotics as septic screen revealed sterile cultures. Two patients (case no 2 and 5) received I.V diuretics in addition. Three patients (1, 2 and 5)
61 improved fully clinically and radiologically. Two patients required ventilatory support and died even after intensive management.

62 **Discussion**

63 PRES has been frequently associated with hypertensive emergencies, pre-eclampsia and treatment with immunosuppressive drugs. Less
64 common causes include uremia, digitoxin toxicity and dialysis disequilibrium (7, 8). The association of PRES and renal failure has been reported
65 in few case reports only (9, 10). No large data exists on this association. We report this first case series of PRES associated with renal failure. The
66 presentation of PRES, its association with renal failure, course and outcome was studied. The reversibility of the syndrome was confirmed both
67 clinically as well as by neuroimaging studies in three cases. In the rest of two cases, course of illness was complicated by severe sepsis and septic
68 shock. Both patients died within 48 hrs of admission. The likely cause of PRES in our patients was accelerated hypertention and uremia. The
69 cause of renal failure in three patients was thought due to hypertention and septicemia. In one patient (case no 2) however, renal failure was
70 likely due to acute glomerulonephritis (postinfectious) and hypertention.

71 All the patients had Hypertention at presentation. Four patients presented with acute renal failure and one patient had established ESRD. Two
72 patients were managed in ICU and developed resistant hypertention with severe sepsis syndrome. Hypertention and sepsis was well controlled
73 in other three patients and all three patients recovered fully. It was therefore observed that control of hypertention, aggressive treatment of
74 infection and supportive care contributes towards favorable outcome.

75 The favored pathogenic theory for PRES suggests autoregulatory disturbance with hyperperfusion, resulting in the shifting of fluid from the
76 intravascular compartment and consequent vasogenic edema (2). It is unknown why the posterior circulation is preferentially affected. A
77 possible explanation is the lower sympathetic innervation of posterior cerebral arterial circulation, with a consequent reduced autoregulation of
78 already impaired cerebral areas (11). Acute renal failure otherwise carries high mortality and morbidity (12, 13). Further studies are needed to
79 know whether increased mortality is due to complications by renal failure and uremia or combinations of two conditions per se.

80

81 The PRES presenting in renal failure with varied presentation and in different age groups suggests that the syndrome should be kept in the
82 differential diagnosis of patients presenting with altered sensorium, seizures and hypertensive crisis irrespective of age. It will greatly help in
83 patient management and better outcome. This study signifies that although PRES is a reversible entity but when complicated by conditions like
84 renal failure, resistant hypertension or sepsis, condition may prove fatal.

85

86 **References**

87 1. Hinchey J, Chaves C, Appignani B, Breen J, Pao L, Wang A et al. A reversible posteriorencephalopathy syndrome. *N Engl J Med* 1996; 334: 494–
88 500

89

90 2. Kwon S, Jahoon K, Sangkwon L. Clinical spectrum of Reversible Posterior Leukoencephalopathy Syndrome. *Pediatr Neurol* 2001; 24:
91 361–64

92

93 3. Covarrubias DJ, Luetmer PH, Campeau NG. Posterior Reversible Encephalopathy Syndrome: Prognostic Utility of Quantitative Diffusion-
94 Weighted MR Images. *Am J Neuroradiol* 2002; 23: 1038–48

95

96 4. Casey SO, Sampaio RC, Michel E, Truwit CL. Posterior reversible encephalopathy syndrome: utility of fluid-attenuated inversion recovery
97 MR imaging in the detection of cortical and subcortical lesions. *Am J Neuroradiol* 2000; 21: 1199–206

98

99 5. Servillo G, Striano P, Striano S et al: Posterior Reversible Encephalopathy Syndrome (PRES) in obstetric critically ill patients.

100 *Intensive Care Med* 2003; 29: 2323–26

101

102 6. Sheth KN, Wu GF, Messe SR et al. Dialysis disequilibrium: another reversible posterior leukoencephalopathy syndrome? Clin Neurol
103 Neurosurg 2003; 105: 249–52

104 7. Kwon S, Jahoon K, Sangkwon L. Clinical spectrum of Reversible Posterior Leukoencephalopathy Syndrome. Pediatr Neurol 2001; 24:361–64
105

106 8. Covarrubias DJ, Luetmer PH, Campeau NG. Posterior Reversible Encephalopathy Syndrome: Prognostic Utility of Quantitative Diffusion-
107 Weighted MR Images. Am J Neuroradiol 2002; 23: 1038–48

108
109 9. Jorge S, Lopes JA, De Almeida E, Martins Prata M. Posterior reversible encephalopathy syndrome (PRES) and chronic kidney disease.
110 Nefrologia 2007; 27: 650

111
112 10. Burrus TM, Mandrekar J, Wijidicks EF, Rabinstein AA. Renal failure and posterior reversible encephalopathy syndrome in patients with
113 thrombotic thrombocytopenic purpura. Arch Neurol 2010; 67: 831-4
114

115 11. Schwartz RB, Feske SK, Polak JF et al. Preeclampsia-eclampsia: clinical and neuroradiographic correlates and insight into the pathogenesis of
116 hypertensive encephalopathy. Radiology 2000; 217: 371–76

117 12. Xue JL, Daniels F, Star RA, Kimmel PL, Eggers PW, Molitoris BA, et al. Incidence and mortality of acute renal failure in Medicare beneficiaries,
118 1992 to 2001. J Am Soc Nephrol 2006; 17: 1135–42

119 13. Levy EM, Viscoli CM, Horwitz RI. The effect of acute renal failure on mortality: a cohort analysis. JAMA.1996; 275: 1489-1494

120

121

122

123 **Table 1: Clinical and Biochemical characteristics of studied population**

Patient data	1	2	3	4	5
SEX/AGE (years)	Female/34	Male/ 11	Female/58	Female/ 22	Male /50
PRESENTATION	Headache, Vomiting, Seizures, altered sensorium (7 days) / Hypertention for 8 years, End stage renal disease on dialysis for 4 years.	Altered sensorium, Generalized swelling of the body, decreased urine output and Headache (2 weeks).	Altered sensorium, Headache and Vomiting (10 days) / received chemotherapy 4 years back for breast cancer, in remission.	Fever, Headache, altered sensorium and seizures (5 days).	Headache , altered sensorium, decreased urine output and abdominal pain (5 days) / Hypertention for 6 years
B.P (mmHg)	230/120	170/110	220/110	220/120	180/100
CREATININE (mg/dl)	7.8	2.5	3.5	6.4	3.5
UREA (mg/dl)	225	55	95	176	147
HEMOGLOBIN (g/dl)	9.4	11.6	11.3	9.2	10.2
WHITE BLOOD CELLS ($\times 10^9/L$)	10.31	22.8	9.8	14.6	10.2
PLATELET COUNT/ mm^3	153,000	318,000	185,000	121,000	123,000
PYURIA	Absent	Positive	Positive	Positive	Absent
PROTEINURIA	Negative	Negative	Negative	Negative	Negative
OUTCOME	Improved	Improved	Died	Died	Improved
FOLLOW UP MRI	Normal	Normal	Not applicable	Not applicable	Normal

124 B.P – Blood Pressure; MRI – Magnetic Resonance Imaging


125

Figure 1

A1

MRI Brain at presentation

T2W Images




126

127 MRI,n

128

Figure 2

MRI Head



ct pressn