

**Original Research Article****Antimicrobial and anti-inflammatory effects of some marine red algae isolated from Quseir, Red Sea, Egypt.****ABSTRACT**

**Aims:** This study aimed to analyze the bioactivities of two different species of marine red algae crude extracts; *Acanthophoraspicifera* (M.Vahl) Borgese and *Digenea simplex* (Wulfen) C.Agardh.

**Place & duration of study:** Sample: Quseir city, Red Sea, Egypt at (April, 2015), extraction and biological study; pharmacognosy department, summer 2015, Kasr El-Eini Teaching Hospitals, Cairo University, Egypt.

**Methodology:** The crude extract screened against some human pathogenic bacteria including ten Gram positive, ten Gram negative strains and unicellular and filamentous fungi. The antimicrobial activity was done through the agar well diffusion method. The anti-inflammatory effect of the crude extract on inflamed liver cells was evaluated by measuring SOD, MDA, catalase, GSH, IL-6 and TNF-alpha in the serum of treated Wistar albino rats compared to positive and negative control groups.

**Results:** The antimicrobial study showed that both *Acanthophoraspicifera* and *Digenea simplex* extracts showed the highest zone of inhibition against *Streptococcus agalactiae* with clear zones ( $22.5 \pm 0.58$  mm and  $23.1 \pm 0.58$  mm) as Gram positive bacteria compared to Ampicillin as positive control. But with Gram negative bacteria *Acanthophoraspicifera* showed highest zone of inhibition against *Pseudomonas aeruginosa* ( $22.3 \pm 2.1$  mm) and *Digenea simplex* showed highest zone of inhibition against *Serratiamarcescens* ( $24.1 \pm 0.58$  mm) compared to Gentamycin as positive control. Finally, the effect of both *Acanthophoraspicifera* and *Digenea simplex* against fungi showed highest zone of inhibition ( $23.8 \pm 0.63$  mm and  $22.4 \pm 2.1$  mm) against *Geotricum candidum*. On the other hand results revealed that all methanol extracts had equally potent anti-inflammatory effects on inflamed liver rats.

**Conclusion:** *Acanthophoraspicifera* (M.Vahl) Borgese and *Digenea simplex* (Wulfen) C.Agardh might be a good source of anti-inflammatory and antibacterial activity.

**Keywords:** Marine red macro-algae, Antimicrobial, Anti-inflammation.

**1. INTRODUCTION**

Marine macro algae are one of nature's most biologically active resources, as they hold a wealth of bioactive compounds. Many compounds isolated from marine macroalgae have demonstrated various biological activities, such as antibacterial activity, antioxidant potential, anti-inflammatory properties, anticoagulant activities, antiviral activities, apoptotic activities, and prebiotic activity [1-6]. Marine macro algae are commonly classified into three main groups based on their pigmentation; Phaeophyta (brown algae), Chlorophyta (green algae), and Rhodophyta (red algae). Macroalgae was considered as a rich source of dietary fiber, minerals, lipids, proteins, omega-3 fatty acids, essential amino acids, polysaccharides, and vitamins A, B, C, and E. Various bioactive compounds from marine organisms have been experimentally tested to comprehensively study the biological effects of recently developed drugs [7]. Studies on the bioactivities of marine algae have revealed numerous health-promoting effects, including anti-oxidative, anti-inflammatory, antimicrobial, and anti-cancer effects. Seaweeds are an excellent source of components such as polysaccharides, tannins, flavonoids, phenolic acids, bromophenols, and carotenoids have exhibits different biological activities, relating on their solubility and polarity, different solvents shows different antimicrobial activity. Chemical compounds should be extracted from different seaweeds in order to optimize their antibacterial and antifungal activity by selecting the best solvent system [8]. Oxidative stress plays important role in endothelial dysfunction [9], lung disease [10], gastrointestinal dysfunction [11], and atherosclerosis [12], all of which involve inflammatory reactions. Many marine natural products that contain antioxidants were known to have anti-inflammatory effects [13-15]. This study was carried out to figure the antimicrobial and anti-inflammatory effects of 2 different algal species.

## 2. MATERIAL AND METHODS

### 2.1. Collection and Identification of marine macroalgae

The fresh algal species were collected from the inter-tidal region of Quseir city (Figure 1) which located on the west coastal area of Red Sea shore between longitude  $34^{\circ} 17' E$  and latitude  $26^{\circ} 06' N$  during the spring year 2015. Collected sample was immediately brought to the laboratory in new plastic bags containing pond water to prevent evaporation. Algal material was washed thoroughly with tap water and distilled water to remove extraneous materials and shade-dried for 5 days and oven dried at  $60^{\circ}C$  until constant weight was obtained, then was grind into fine powder using electric mixer and stored at  $4^{\circ}C$  for future use. Algal species were identified according to [16-19].



Fig.1 Map show the studied area [Quseir city, Red Sea, Egypt]

### 2.2. Extraction of selected algal species:

Powdered marine algae (500 g, each) were extracted with 70% methanol by percolation till exhaustion. The alcoholic extract in each case was evaporated under reduced pressure to obtain a semisolid residue.

### 2.3. Antimicrobial activity

The hydroalcoholic extracts of the selected species were investigated for their antimicrobial activities using Agar well diffusion and Meller-Henton against gram positive and gram negative bacteria and used Sab. dextrose agar against some fungi. All tested microorganisms were kindly supplied from Biotechnology Research Center, Al-Azhar University (for boys), Cairo, Egypt.

#### 2.3.1. Gram positive bacteria:

*Staphylococcus aureus* (RCMB 010027), *Staphylococcus epidermidis* (RCMB 010024), *Streptococcus sanguis* (RCMB 01001 71-3), *Streptococcus pyogenes* (RCMB 01001 74-2),

*Streptococcus agalactiae*(RCMB 01001 73-2), *Bacillus subtilis* (RCMB 01001 69-3), *Enterococcus faecalis*(RCMB 01001 54-2),*Corynebacteriumdiphtheriae*(RCMB 01001 26-7), *Micrococcus luteus*(RCMB 01001 76-9), *Methicillin-resistant Staphylococcus aureus* MRSA (RCMB 01001 94-5).

### 2.3.2. **Gram negative bacteria:**

*Escherichia coli* (RCMB 01002 52-6), *Proteus mirabilis* (RCMB 01002 54-2), *Acinetobacterbaumannii* (RCMB 01002 82-9),*Klebsiella pneumonia*(RCMB 01002 23-5), *Pseudomonas aeruginosa* (RCMB 01002 43-5), *Serratiaplymuthica*(RCMB 01002 75-3), *Serratiamarcescens*(RCMB 01002 75b-8), *Salmonella typhi*(RCMB 01002 15-4), *Enterobacter cloacae* (RCMB 01002 64-5), *Shigelladysenteriae*(RCMB 01002 41-8).

### 2.3.3. **Unicellular fungi & Filamentous fungi:**

*Aspergillus fumigatus* (RCMB 02568), *Syncephalastrum racemosum* (RCMB 05922), *Geotricum candidum* (RCMB 05097), *Candida albicans* (RCMB 05036), *Aspergillus niger* (RCMB 02724), *Cryptococcus neoformans* (RCMB 05642), *Candida trobicalis* (RCMB05239), *Penicillium expansum* (RCMB 01924), *Microsporium canis* (RCMB 0834),*Trichophyton mentagrophytes* (RCMB 0925).

### 2.3.4. **Methods of measuring antimicrobial activity:**

#### 2.3.4.1. **Well-diffusion method for anti-bacterial activity: (Mueller Henton)**

The solution of 50 mg/ml of each sample in DMSO (Dimethyl sulphoxide) was prepared for testing against bacteria. Centrifuged pellets of bacteria from 24h old culture containing approximately 104-106 CFU/ml (Colony forming unit per ml) were spread on the surface of Nutrient agar (type tone 1%, Yeast extract 0.5%, agar 1%, 100 ml of distilled water, PH 7.0) which autoclaved under 121°C for at least 20 min. Wells were created in medium with the help of a sterile metallic bores and then cooled down to 45°C. The activity was determined by measuring the diameter of the inhibition zone (in mm). 100µl of the tested samples (100 mg/ml) were loaded into the wells of the plates. All samples were prepared in DMSO. DMSO was loaded as control. The plates were kept for incubation at 37°C for 24h and then the plates were examined for the formation of zone of inhibition. Each inhibition zone was measured three times by caliper to get an average value. The test was performed three times for each bacterium culture. Ampicillin and Gentamycin were used as antibacterial standard drugs[20].

#### 2.3.4.2. **Well-diffusion method for anti-fungal activity: (Sabourad dextrose)**

The antifungal activity was investigated by agar well diffusion method by the following procedure: Sabourad dextrose agar plates: A homogenous mixture of glucose-peptone-agar (40: 10: 15) was sterilized by autoclaving at 121°C for 20 min. The sterilized solution (25 ml) was poured in each sterilizedpetridish in laminar flow and left for 20 min to form the solidified sabourad dextrose agar plate. These plates were inverted and kept at 30 °C in incubator to remove the moisture and check for any contamination. Antifungal assay: Fungalstrain was grown in 5ml Sabourad dextrose broth (glucose: peptone; 40: 10) for 3-4 days to achieve 105 CFU/ml cells. The fungal culture (0.1 ml) was spread out uniformly on the Sabourad dextrose agar plates. Now small wells of size (4mm×20mm) were cut into the plates with the help of well cutter and bottom of the wells were sealed with 0.8% soft agar to prevent the flow of test sample at the bottom of the well. 100µl of the tested samples (10mg/ml) were loaded into wells of the plates. All samples was prepared in dimethyl sulfoxide (DMSO), DMSO was loaded as control. The plates were kept for incubation at 30 °C for 3-4 days and then the plates were examined for the formation of zone of inhibition. Each inhibition zone was measured three times by caliper to get an average value. The test was performed three times for each fungus. Amphotericin B was used as antifungal standard drug [20].

## 2.4. **Anti-inflammatory activity**

Animal: Wister albino rats weighing between (140-200 g) were obtained from animal house of National Research Center, Dokki – Giza, Egypt. They were housed in well protected polypropylene cages and maintained in standard laboratory conditions in an air conditioned area at 25±2°C in 10:14 hr light dark cycle and allowed to take standard laboratory feed and tap water [21]. The study was approved by **Institutional Animal Ethics Committee of National Research Center** that the animal doesn't suffering at any stage of experiment and maintained in accordance with the Guide for the Care and Use of Laboratory Animals. The inflamed liver was induced in rats according to the described method[22].The animals were divided into three groups consisted of six animals; Normal control (A), Diseased or inflamed Liver cancer (positive control receiving saline) (B), Inflamed liver rats were treated by injection with methanol extract of the first extract (*Acanthoporaspecifica*)(C) and the second extract (*Digenea simplex*)(D).The methanol extract (0.1 g of each sample, separately) mixed or dissolved with 1ml of phosphate buffer and every day injected 100 µm to each rat for 15 days. After 15 days, rats were sacrificed and blood samples were collected

by puncture the sublingual vein in clean and dry test tube. Allow clotting for 10 minutes before centrifuging at 3000 rpm for serum separation. The separated serum was stored at  $-80^{\circ}\text{C}$  for further determinations of the following tests: SOD, MDA, catalase, GSH, IL-6 and TNF-alpha. The anti-inflammatory activity was observed on Wistar albino rats after injection by the extracts of the marine algal species (Table 4) by measuring the parameters in the normal and diseased states.

### 3. RESULTS AND DISCUSSION

#### 3.1. Collection and Identification

Isolated algal species: Two algal samples of red algae *Acanthophoraspicifera* (M. Vahl) Borgese (Fig. 2) and *Digenea simplex* (Wulfen) C. Agardh (Fig. 3) were collected from Red Sea, Egypt (Quseir city).



Figure 2. *Acanthophoraspicifera* Figure 3. *Digenea simplex*

#### 3.2. Antimicrobial activity

Results summarised in the tables (1-3) showed the antimicrobial screening of the crude extracts of two macro algae against some human pathogenic bacteria as well as some unicellular and filamentous fungi. Both *A.spicifera* and *D.simplex* extracts showed similar potent inhibitory growth activities against three Gram +ve bacteria [*Streptococcus agalactiae*, *Streptococcus pyogenes* and *Streptococcus sanguis*] with zone of inhibition ranging from  $[23.1 \pm 0.58$  to  $20.6 \pm 0.63$  mm] and showed moderate activities with [*Corynebacterium diphtheriae*, *Bacillus subtilis* and *Staphylococcus aureus*] with inhibition zones ranging from  $[20.1 \pm 1.5$  to  $16.3 \pm 2.1$  mm]. Also the crude extracts were found to be more active than the positive control Ampicillin, ( $22.3 \pm 1.5$  mm), against *Streptococcus agalactiae* which showing inhibition zone  $[22.5 \pm 0.58$  mm with *A.spicifera* and  $23.1 \pm 0.58$  mm with *D.simplex*]. *Staphylococcus epidermidis*, *Enterococcus faecalis*, *Micrococcus luteus* and *Methicillin-resistant Staphylococcus aureus* are resistance to the activity of the algae compared to the positive control Ampicillin and Vancomycin, this was on the contrary of previously published activity of *A.spicifera* against MRSA [23]. On the other hand the two algal species showing activities against seven Gram -ve bacteria [*Escherichia coli*, *Proteus mirabilis*, *Klebsiella pneumonia*, *Pseudomonas aeruginosa*, *Serratia plymuthica*, *Serratia marcescens* and *Enterobacter cloacae*] with zones of inhibition ranging from  $[17.2 \pm 1.5$  to  $24.1 \pm 0.58$  mm] similar to previously reported [24]. Both algal extracts showed higher selective activity against *Pseudomonas aeruginosa* with zone of inhibition  $[22.3 \pm 2.1$  mm with *A.spicifera* and  $22.9 \pm 0.63$  mm with *D.simplex*] compared to the positive control Gentamycin ( $20.6 \pm 1.5$  mm) and against *Serratia marcescens* with zone of inhibition  $[22.3 \pm 1.5$  mm with *A.spicifera* and  $24.1 \pm 0.58$  mm with *D.simplex*] compared to the same positive control with ( $20.4 \pm 0.58$  mm of inhibition zone). *Acinetobacter baumannii*, *Salmonella typhi* and *Shigella dysenteriae* are resistance to the effect of the selected algae compared to the positive control Gentamycin. Previous results were different from the reported one that showed the crude extract of *Digenea simplex* as a weak growth inhibition zones ranging from 6 to 8 mm against *Staphylococcus aureus*, *Enterococcus faecalis* and *Pseudomonas aeruginosa* [25]. Discussing the antifungal activity, the algal extracts were found to be active on the selected species of filamentous fungi against six

fungi [*Aspergillus fumigatus*, *Syncephalastrum racemosum*, *Geotricum candidum*, *Aspergillus niger*, *Candida tropicalis* and *Microsporum canis*] with zones of inhibition ranging from [18.6±0.63 to 23.8±0.63 mm]. *Geotricum candidum* and *Aspergillus niger* are the most sensitive fungal species against both *A. spicifera* with clear zones [23.8±0.63 and 20.6±0.58 mm] and *D. simplex* with clear zones [22.4±2.1 and 21.2±0.63 mm] compared to the positive control Amphotericin B with zone of inhibition [20.3±1.5 and 20.3±0.58 mm]. *Candida albicans*, *Cryptococcus neoformans*, *Penicillium expansum* and *Trichophyton mentagrophytes* were resistance to the selected algae compared to the positive control Amphotericin B, on the other hand this result was differ from previous reported where the inhibition zone (6 mm) was recorded in methanol extract against *C. albicans* [26].

**Table 1. Antimicrobial activity of the studied algal species against Gram +ve bacteria.**

Sample	<i>Acanthophora spicifera</i>	<i>Digenea simplex</i>	St.
Gram positive bacteria			Ampicillin
<i>Staphylococcus aureus</i> (RCMB 010027)	16.3±2.1	17.1±2.1	22±1.0
<i>Staphylococcus epidermidis</i> (RCMB 010024)	NA	NA	23±1.0
<i>Streptococcus sanguis</i> (RCMB 01001 71-3)	20.6±0.63	21.2±0.63	21.7±1.5
<i>Streptococcus pyogenes</i> (RCMB 01001 74-2)	21.4±1.2	21.6±1.2	22.7±1.5
<i>Streptococcus agalactiae</i> (RCMB 01001 73-2)	22.5±0.58	23.1±0.58	22.3±1.5
<i>Bacillus subtilis</i> (RCMB 01001 69-3)	18.3±0.63	19.3±1.2	25.3±1.5
<i>Enterococcus faecalis</i> (RCMB 01001 54-2)	NA	NA	19.3±0.58
<i>Corynebacterium diphtheria</i> (RCMB 01001 26-7)	19.3±1.2	20.1±1.5	20±1.0
<i>Micrococcus luteus</i> (RCMB 01001 76-9)	NA	NA	19.6±1.5
Methicillin-resistant microorganisms			vancomycine
Methicillin-resistant <i>Staphylococcus aureus</i> MRSA (RCMB 01001 94-5)	NA	NA	21.6±2.1

**Table 2. Antimicrobial activity of the studied algal species against Gram -ve bacteria.**

Sample	<i>Acanthophora spicifera</i>	<i>Digenea simplex</i>	St.
Gram negative bacteria			Gentamycine
<i>Escherichia coli</i> (RCMB 01002 52-6)	18.9 ± 1.2	19.1± 1.2	20.3± 0.85
<i>Proteus mirabilis</i> (RCMB 01002 54-2)	17.2± 1.5	17.8± 1.5	21.2± 1.2
<i>Acinetobacter baumannii</i> (RCMB 01002 82-9)	NA	NA	23.4± 1.2

<i>Klebsiella pneumonia</i> (RCMB 01002 23-5)	21.4 ± 1.2	21.6 ± 0.72	27.2± 2.1
<i>Pseudomonas aeruginosa</i> (RCMB 01002 43-5)	22.3 ± 2.1	22.9± 0.63	20.6± 1.5
<i>Serratia plymuthica</i> (RCMB 01002 75-3)	20.2 ± 0.72	20.8 ± 2.1	22.3± 0.58
<i>Serratiamarcescens</i> (RCMB 01002 75b-8)	22.3 ± 1.5	24.1 ± 0.58	20.4± 0.58
<i>Salmonella typhi</i> (RCMB 01002 15-4)	NA	NA	21.1± 0.72
<i>Enterobacter cloacae</i> (RCMB 01002 64-5)	21.3 ± 0.58	21.8± 1.5	22.4± 2.1
<i>Shigelladysenteriae</i> (RCMB 01002 41-8)	NA	NA	21.3± 1.5

**Table 3. Antimicrobial activity of the studied algal species against the tested fungi**

Sample	<i>Acanthophora spicifera</i>	<i>Digenea simplex</i>	St.
Fungi			Amphotericin B
<i>Aspergillus fumigatus</i> (RCMB 02568)	21.3±1.2	22.4±0.58	25.7±1.5
<i>Syncephalastrum racemosum</i> (RCMB 05922)	19.3±1.2	20.3±1.2	24.3±1.2
<i>Geotricum candidum</i> (RCMB 05097)	23.8±0.63	22.4±2.1	20.3±1.5
<i>Candida albicans</i> (RCMB 05036)	NA	NA	21.3±1.5
<i>Aspergillus niger</i> (RCMB 02724)	20.6±0.58	21.2±0.63	20.3±0.58
<i>Cryptococcus neoformans</i> (RCMB 05642)	NA	NA	21±1.0
<i>Candida tropicalis</i> (RCMB 05239 )	18.6±0.63	20.3±1.2	23.7±2.0
<i>Penicillium expansum</i> (RCMB 01924)	NA	NA	21.7±2.0
<i>Microsporium canis</i> (RCMB 0834)	20.3±1.2	21.9±0.58	23.3±1.5
<i>Trichophyton mentagrophytes</i> (RCMB 0925)	NA	NA	21.3±1.5

Data in table 1,2,3 are represented as mean zone of inhibition ± standard deviation.

RCMB: Regional Center of Mycology and Biotechnology Antimicrobial unit test organism.

NA: No activity.

### 3.3. Anti-inflammatory activity

In this study table and figure no. 4 represent the levels of SOD, MDA, Catalase, GSH, IL-6 and TNF-Alpha in the liver tissue homogenates of the normal and treated groups. Each parameter showed significant different degree in the disease state compared to the normal 'control' state. As it observed significant decreased in the levels of SOD from (72.03±6.49 to 5.93±0.31 U/ml), Catalase from (75.68±2.6 to 6.96±0.31 U/ml) and GSH from (58.85±3.6 to 5.03±0.29µmol/ml). Also observed that significant increased in the inflamed liver state of MDA from (20.31±0.90 to 133.56±5.10nmol/ml), IL-6 from (15.51±1.04 to 144.1±5.3Pg/ml) and TNF-Alpha from (23.89±1.07 to 257.56±11.85Pg/ml). After treatment of inflamed liver rats with crude extracts using a dose of 500mg/kg of each drug, observed that the difference in the parameters levels relatively equal to the normal control results.

**Table 4. Anti-inflammatory activity of the studied algal species.**

parameter						
Group	SOD (U/ml)	MAD (nmol/ml)	Catalase (U/ml)	GSH (μmol/ml)	IL-6 (Pg/ml)	TNF-Alpha (Pg/ml)
Group A	72.03±6.49	20.31±0.90	75.68±2.6	58.85±3.6	15.51±1.04	23.89±1.07
Group B	5.93±0.31	133.56±5.10	6.96±0.31	5.03±0.29	144.1±5.3	257.56±11.85
Group C ( <i>Acanthoporaspi</i> <i>cifera</i> )	69.18±0.51	16.61±0.30	70.46±0.40	56.01±0.58	18.35±1.03	29.08±1.04
Group D ( <i>Digenea simplex</i> )	70.18±0.73	18.25±0.29	72.3±0.36	56.63±0.33	19.63±1.02	26.58±1.95
F-Prob	P<0.0001	P<0.0001	P <0.0001	P<0.0001	P<0.0001	P<0.0001
LSD at5%	9.68	7.67	3.98	5.43	8.26	15.16
LSD at1%	13.20	10.46	5.43	7.41	11.27	22.32


Data in table 4 expressed as Mean ± S.E.M = Mean values ± Standard error of means of six experiments. Number of animals in each group is six.

Statistical evaluation of results was performed using the one-way ANOVA and least significant difference was determined at the level of LSD at 5% and LSD at 1%.

Test drugs: significant from normal control, P < 0.0001.

SOD: Superoxide Dismutase, MAD: Malonaldehyde, GSH: Glutathione, IL-6: Interleukin-6, TNF-α: Tumor necrosis factor.

Group A: Normal control, Group B: Diseased or inflamed Liver cancer (positive control receiving saline), Group C and D: Inflamed liver rats were treated by injection with methanol extract of the first drug (*Acanthoporaspicifera*) and the second drug (*Digenea simplex*).



**Fig. 4: Anti-inflammatory activity of *Acanthophora spicifera* and *Digenea simplex* on inflamed liver Wister Albino rats.**

#### 4. CONCLUSION

The bioactive components from seaweeds crude extract have antimicrobial activity on some human pathogenic bacterial species and unicellular and filamentous fungi and anti-inflammatory effect which might protect the human health against some oxidative stress which attack DNA, proteins and membrane lipids and induced cellular damage.

#### Reference

1. Haroun, B. M., Sharaf, A.M. and Ibraheem B. Evaluation of natural productions in some common Egyptian marine algae. *J. Union. Arab Biol., B, Botany*, 1995, 2, 137-153.
2. O' Sullivan, L., B. Murphy, P. McLoughlin, P. Duggan, P.G. Lawlor, H. Hughes, and G.E. Gardiner. 2010. Prebiotics from marine macroalgae for human and animal health applications. *Marine Drugs* 8(7): 2038–2064.
3. Ibraheem, B.M.I., Neveen, A.R., Mohamed, S.A.H. and Khaled, E.Y. Antimicrobial and antiviral activities against newcastle disease virus (NDV) from marine algae isolated from qusier and marsaalam seashore (red sea), Egypt. *African Journal of Biotechnology*. 2012. 11: 8332-8340.
4. Elsayed, K.N.; Radwan, M.M.; Hassan, S.H.; Abdelhameed, M.S.; Ibraheem, I.B.; Ross, S.A. Phytochemical and biological studies on some Egyptian seaweeds. *Nat. Prod. Commun.* 2012, 7, 1209–1210.
5. Al-Saif S.S., Abdel-Raouf N., El-Wazanani H.A., Aref I.A., Antibacterial substances from marine algae isolated from Jeddah coast of Red sea, Saudi Arabia, *Saudi journal of biological sciences*, 21 (2014) 57-64.
6. Abdel-Raouf, N., Al-Enazi, N. M., Al-Homaidan, A. A., Ibraheem, I.B.M., Al-Othman, M. R. and Hatamleh, Antibacterial  $\beta$ -amyryn isolated from *Laurenciamicrocladia*. *Arabian Journal of Chemistry*, 2015, 8, 1, 32–37.
7. Arif JM, Al-Hazzani AA, Kunhi M, Al-Khodairy F: Novel marine compounds: Anticancer or genotoxic? *J Biomed Biotechnol* 2004, 2004(2):93–98.
8. Hediat MH. Salama, NajatMarraiki. (2010). Antimicrobial activity and phytochemical analyses of *Polygonum aviculare* L. (Polygonaceae), naturally growing in Egypt. *Saudi Journal Of Biological Sciences*: 17. 57- 63.
9. Schramm A, Matusik P, Osmenda G, Guzik TJ: Targeting NADPH oxidases in vascular pharmacology. *VascPharmacol* 2012, 56(5–6):216–231. Rosanna DP, Salvatore C: Reactive oxygen species, inflammation, and lung diseases. *Curr Pharm Des* 2012, 18(26):3889–3900.
10. Rosanna DP, Salvatore C: Reactive oxygen species, inflammation, and lung diseases. *Curr Pharm Des* 2012, 18(26):3889–3900.
11. Kim YJ, Kim EH, Hahm KB: Oxidative stress in inflammation-based gastrointestinal tract diseases: challenges and opportunities. *J GastroenterolHepatol* 2012, 27(6):1004–1010.
12. Hulsmans M, Van Dooren E, Holvoet P: Mitochondrial reactive oxygen species and risk of atherosclerosis. *CurrAtheroscler Rep* 2012, 14(3):264–276.
13. Abad MJ, Bedoya LM, Bermejo P: Natural marine anti-inflammatory products. *Mini Rev Med Chem* 2008, 8(8):740–754.
14. Wang W, Wang SX, Guan HS: The antiviral activities and mechanisms of marine polysaccharides: an overview. *Mar Drugs* 2012, 10(12):2795–2816.
15. D'Orazio N, Gammone MA, Gemello E, De Girolamo M, Cusenza S, RiccioniG: Marine bioactives: pharmacological properties and potential applications against inflammatory diseases. *Mar Drugs* 2012, 10(4):812–833.
16. Aleem, A.A. (1978). Contributions to the study of the marine algae of the Red Sea. III- Marine algae from Obhor, in the vicinity of Jeddah, Saudi Arabia. *Bull. Fac. Sci. KAU J.* 2, 99-118.
17. Bold, H.C. (1978). " Introduction to the algae" structure and reproduction. Prentice. Hall. Inc., New-Jersey 07632.

18. Aleem, A.A. (1993). The marine algae of Alexandria. Egypt. Egyptian Books House.
19. Coppejans, E., Leliaert, F., Dargent, O., Gunasekara, K. and Clerck, O. (2009). Srilanka seaweeds. Methodologies and field guide to the dominant species. University of Ruhuna, Dept. of Botany, Matura, Srilanka. 1-265.
20. Rahman A.;Choudhary, M. and thomsen W.(2011). Bioassay Techniques for Drug Development. Hardwood Academic publishers,the Netherlands,pp.16.
21. INTERNATIONAL, A. 2005. Official methods of analysis of AOAC International, AOAC International.
22. KEEGAN, D. 1996. Foundations of distance education, Psychology Press.
23. Tapiero, H., Tew, K.D., Naguyen, G., Math\_e, G., 2002. Polyphenods: do they play a role in the prevention of human pathologies? Review. Biomed. Pharmacother. 56, 200–207.
24. Wang J, Zhang Q, Zhang Z (2008) Antioxidant activity of sulfated polysaccharide fractions extracted from Laminaria japonica. Int J BiolMacromols 42:127–132.
25. Frlich, I., &Riederer, P. (1995). Free radical mechanisms in dementia of Alzheimercetype and the potential for antioxidative treatment. Drug Research, 45, 443–c449.
26. Butterfield, D. A., Castenga, A., Pocernich, C. B., Drake, J., Scapagnini, G., &Calabrese,cV. (2002). Nutritional approaches to combat oxidative stress in Alzheimer’s disease.cJournal of Nutritional Biochemistry, 13, 444–461.