

1 **Assesment of phenology and morphological diversity of 3 species of Asteraceae:**  
2 ***Anacyclus clavatus*, *Chamaemelum fuscatum* and *Leucanthemum parthenium***

3

4

5

6

7

8

9

10

11

12

13

14

15 **Abstract**

16 | **Aim:** ~~3~~Three species of Asteraceae: *Anacyclus clavatus*, *Chamaemelum fuscatum* and  
17 *Leucanthemum parthenium* that have a wide range of uses in medicine and in industry were  
18 characterized by inter-specific variations and phenological activities.

19 **Study Design:** Morphological characterization of these 3 species using 18 quantitative traits and  
20 phenology study: vegetative period, flowering and fruiting time and seed formation for two consecutive  
21 years.

22 **Place and Duration of Study:** Experimental plot at the Faculty of Sciences of Tunis, Tunisia- 2009-  
23 2010.

24 **Methodology:** Measurements of the 18 morphological characters were performed on 3 samples of  
25 *Anacyclus clavatus*, *Chamaemelum fuscatum* and *Leucanthemum parthenium* grown in the Faculty of  
26 Sciences of Tunis, for each species, we have studied 10 individuals. Different phenological stages:  
27 Vegetative period, Flowering and Fruiting of each species are studied.

28 **Results:** The phenological study show that the 3 species studied have distinct phenologies. The  
29 longest phenological cycle is observed for *Leucanthemum parthenium*. Results of morphology study  
30 showed significant differences to highly significant for the majority of the traits studied using variance  
31 analysis. The comparison of means reveals that *Anacyclus clavatus* and *Chamaemelum fuscatum*  
32 form a single group for most of the traits measured, while *Leucanthemum parthenium* is clearly  
33 distinct from these two species. In addition, the principal component analysis confirms the results of  
34 the variance analysis and the comparison of means.

35 **Conclusion:** The results of the phenological cycle's follow-up show that the 3 species studied have  
36 distinct phenologies. The longest phenological cycle is observed for *Leucanthemum parthenium*. The  
37 morphological study reveals that *Anacyclus clavatus* and *Chamaemelum fuscatum* form a single  
38 group while *Leucanthemum parthenium* is clearly distinct from these two species.

39 **Keywords:** *Anacyclus clavatus*; *Chamaemelum fuscatum*; *Leucanthemum parthenium*;  
40 morphological; phenology.

41

## 42 1. Introduction

43 Phenological study is important in plant management and combating afforestation, honey analysis,  
44 floral biology, estimation of reproductivity and regeneration [1]. It is important also in understanding  
45 species interrelations and their interaction with the environment. Variations in phenophases among  
46 individuals of different species have been linked to environmental perturbations [2]. A clear  
47 understanding of phenological behavior on time of anthesis, time and duration of stigma receptivity,  
48 fertilization, mode of pollination, seed development is necessary for breeding programs to obtain

49 better traits [3]. Thus plant phenological study has great significance because it not only provides  
50 knowledge about the plant growth pattern but it also provides the idea on the effect of environment  
51 and selective pressure on flowering and fruiting behavior [4].

52 Evaluation and characterization through morphological parameters of different crop germplasm is  
53 therefore so much important for all plant breeders [5]. Therefore, it is important to make proper  
54 strategies for the collection and evaluation of germplasm sources which are locally used in different  
55 regions of the world and save them from being vanished [6]. To have a variety of better traits of any  
56 crop we need information's about its genetic diversity [7]. Thus, characterization and estimation of  
57 genetic diversity is an important step for the competent and successful maintenance and utilization of  
58 different crop germplasm [8].

59 Genetic diversity is an inherited variation among and between populations, created, activated and  
60 maintained by evolution [9]. Morphological traits provide a simple way of measuring genetic diversity  
61 while studying genotype performance under normal growing conditions, but are influenced by  
62 environmental factors ([10]; [11]). Plants have the potential to response to the changed environments  
63 by changing their morphology and there for, the intra-specific variation in plant characteristics is  
64 usually regarded as the adaptive mechanism to different environments [12].

65 | The Asteraceae is one of the largest families, comprising 250,000 species. It is known for its wide  
66 range of uses not only in medicine but also some plants are grown as ornamental plants such as  
67 chamomile (*Leucanthemum parthenium*), others can provide different products: natural rubber,  
68 colorants, insecticides and spices [13].

69 *A. clavatus* (*Anacyclus clavatus*), belonging to the Asteraceae family, is an herbaceous, annual and  
70 spontaneous plant that is found almost everywhere in the Mediterranean region [14]. It's 20 to 50 cm  
71 tall, hairy, green or whitish-green, with an upright or ascending stem, woolly and rowdy whose  
72 branches are divorced. Leaves are bipinnatized, long to very narrow segments terminated by a small  
73 mucron [15]. The convex or somewhat conical receptacle carries triangular bracts, ovals in the shape  
74 of sequins. The inflorescences have two types of hermaphrodite flowers: the central flowers are  
75 yellow-colored and the peripheral flowers are tongued, long and white. They flourished from March to  
76 June [14].

77 The fruits in the form of achenes are small, very compressed cuneiform and of gray to beige color  
78 [15]. The number of chromosomes of this species is  $2n = 18$  [16]. It's a plant that grows on the edges  
79 of fields and roads and in the wastelands of the entire Mediterranean coast [15]. In Tunisia, it's is  
80 located in the north (Kroumirie, Oued Medjerda and Cap Bon), and in the center. The use of this  
81 species is very limited. The aerial part of *A. clavatus* is used as a powder against stomach and belly  
82 pains. It may also be one of the components of tobacco [17].

83 *C. fuscatum* (*Chamaemelum fuscatum*), belonging to the Asteraceae family, anthemidae tribe,  
84 *Anthemis* genus and Ormenis sub-section, is an annual, herbaceous, glabrous 30 cm rowing,  
85 ascending or upright. The leaves are bipinnated. The heads are heterogeneous with yellow disc and  
86 white ligules; their flowering is very early from November to April. The achene is very small, striated,  
87 tetragonal and brown to yellow in colour. It's a very widespread plant on the banks of the seguias.

88 In Tunisia, *C. fuscatum* is found everywhere: in the north (Ain Drahim, Kef, etc.), in the center  
89 (Sousse, Enfidha, etc.) and in the South (Gabes, etc.). Internationally, It's located in the western  
90 Mediterranean basin of Spain, Greece and North Africa (Tunisia, Morocco and Algeria) [15]. The  
91 number of chromosomes of this species is  $2n = 18$  [18]. It's known for its anti-malaria property and its  
92 protective effect against cell damage [19].

93 *L. parthenium* (*Leucanthemum parthenium*) belonging to the Asteraceae family too, the Anthemidae  
94 tribe and the Asteroidea subfamily [20] and the *Leucanthemum* genus. This chamomile is a very  
95 fragrant, perennial, rooted plant, with flowering stem erect without hair. The leaves are deeply divided  
96 into 4 to 12 toothed segments. The internal tubular flowers are yellow and the ligulate external flowers  
97 are white. They flourish from June to August in European conditions [14] and from July to October in  
98 Iran [21]. The ripe fruits are brown, glandular and surmounted by a very short membranous crown.

99 *L. parthenium* is a medicinal plant used primarily for the prevention and reduction of migraine attacks  
100 frequency, against stomach aches and malaria [22]. It's also known for its properties: antiseptic,  
101 stomachic, antihysterical, vermifuge and insecticide. It's found spontaneously on the edges of roads  
102 and often in the vicinity of dwellings and it can also be grown in gardens as an ornamental plant.  
103 Internationally, *L. parthenium* is found almost all over Europe except the boreal zone and it is also  
104 found in South-Western Asia [14].

105 However, there is little information on the morphological diversity and the phenology of *Anacyclus*  
106 *clavatus*, *Chamaemelum fuscatum* and *Leucanthemum parthenium* and the potential of these species  
107 in breeding programs. The aim of this study is to assess the variations in morphology and phenology  
108 of *A. clavatus*, *C. fuscatum* and *L. parthenium*.

## 109 **2. Materials and methods**

### 110 **2.1. Plant material**

111 Three species of Asteraceae have been studied in this work: *Anacyclus clavatus*, *Chamaemelum*  
112 *fuscatum* and *Leucanthemum parthenium*. These species were grown on an experimental plot at the  
113 Faculty of Sciences of Tunis, Tunisia under uncontrolled conditions. The seeds used originate from  
114 Esbikha for *A. clavatus*, Haouz (Morocco) for *C. fuscatum* whereas the seeds of *L. parthenium* are  
115 available in the laboratory of Genetics and Bioresources of the Faculty of Sciences of Tunis.

### 116 **2.2. Phenological characters**

117 Different phenological stages presented by the individuals of each species are defined:

#### 118 **2.2.1. Vegetative period**

119 This stage spreads from the planting to the beginning of flowering. This is the phase of vegetative  
120 growth.

121

#### 122 **2.2.2. Flowering**

123 This is the period during which the flowers appear. The method of study is based essentially on the  
124 visual observation of the appearance of the flowers.

#### 125 **2.2.3. Fruiting**

126 This phase is characterized by the formation of the fruit. It begins with the formation of the first  
127 seeds and ends with the general ripening of the seeds.

### 128 **2.3. Morphological traits**

129 In order to compare the various species studied, we describe the characters of their vegetative  
130 part: The type of branching, the stem, the structure and color of the leaves, the structure and color of  
131 the inflorescences and flowers, the structure and color of akene and the weight of 100 akene.

132 Measurements of the morphological characters were performed on three samples of *Anacyclus*  
133 *clavatus*, *Chamaemelum fuscatum* and *Leucanthemum parthenium* grown in the Faculty of Sciences  
134 of Tunis, for each species, we have studied 10 individuals. The 18 morphological quantitative traits  
135 were assessed to characterize and estimate genetic diversity among the 3 species studied, the  
136 quantitative traits measured were:

- 137 • Length of main axis in cm: LAP
- 138 • Average length of primary branches in cm: LMRP
- 139 • Average length of branches in cm: LMRS
- 140 • Average length of the tertiary branches in cm: LMRT
- 141 • Length of main root in cm: LRP
- 142 • Number of leaves per plant: NF
- 143 • Average diameter of the receptacle in cm: DMR
- 144 • Average number of leaflets per leaf: NLL
- 145 • Average length of the leaf rachis in cm: LMRF
- 146 • Number of inflorescence per plant: NI
- 147 • Number of primary branches: NRP
- 148 • Number of secondary branches: NRS
- 149 • Number of tertiary branches: NRT
- 150 • Average number of ligules per head: NML
- 151 • Number of ligules of the main axis head: NLCAP
- 152 • Length of the smallest branch in cm: LPR
- 153 • Length of the longest branch in cm: LLR
- 154 • Weight of 100 akene :  $P_{100} A$

#### 155 2.4. Data analysis

156 In order to evaluate morphological diversity and to establish relationships among studied species,  
157 several statistical procedures were conducted. Quantitative data were computed using the software  
158 XLSTAT version 2011 to perform analysis of variance, comparison of mean and to calculate the  
159 Pearson correlation coefficient. Principal component analysis (PCA) was also done using the software  
160 XLSTAT.

### 161 3. Results and discussion


#### 162 3.1. Phenology study

##### 163 3.1.1. Vegetative period

164 The vegetative period is characterized by a strictly herbaceous development and extends from  
165 seedling to full bloom. We divided this phase into 2 stages:

166 **Stage of germination:** it is characterized by the appearance of the primordial leaves. In all three  
167 species, the germination begins after 10 days.


168 **Stage of foliage:** Observation of the phenological spectrum reveals that this stage is the longest of  
169 the phenological cycle. This stage, which is characterized by the growth of the stems in length and by  
170 the formation of the leaves, lasts 6 months for *Chamaemelum fuscatum* (Figure 1) and 7 months for  
171 *Anacyclus clavatus* (Figure 2). *Leucanthemum parthenium* is a perennial herb plant (Figure 3).



172

173


Fig.1. Phenological cycle of *Chamaemelum fuscatum*



174

175

Fig. 2. Phenological cycle of *Anacyclus clavatus*



176

177

Fig.3. Phenological cycle of *Leucanthemum parthenium*

178 **3.1.2. Flowering**

179 Flowering is considered from the formation of the first flower until most flowers have evolved this  
 180 period differs from one species to another: For *Chamaemelum fuscatum*, the flowering period ranges  
 181 from mid-February to the end of April (Figure 1). For *Anacyclus clavatus*, this period extends from the  
 182 end of March to mid-May (Figure 2). For *Leucanthemum parthenium*, the first flower blooms in early  
 183 June and full bloom is observed around mid-July (Figure 3).

184 Flowering appears to be highly favoured during the rainy season for *Anacyclus clavatus* and  
 185 *Chamaemelum fuscatum*, only *Leucanthemum parthenium* flowers during the dry season. We find  
 186 that the species *Chamaemelum fuscatum* characterized by a very early flowering date has a spread



187 flowering period. In addition, the species *Leucanthemum parthenium* characterized by a late flowering  
188 date has a relatively short flowering stage and this to escape the water stress.

### 189 3.1.3. Fruiting

190 It is the formation of fruit in the form of akene. We have noticed that the appearance of the first  
191 akene coincides with the peak of flowering, while the full fructification characterized for the 3 species  
192 by the change of color flowers in tubes from yellow to light gray and the fall of the white ligules is  
193 generally obtained after two weeks of the appearance of the first fruit (Figure 1, 2 and 3).

194 In fact, the study of [23] reveals that achenes of *A. clavatus* that germinated earlier produced plants  
195 with higher biomass and higher reproductive effort. In addition, this work show that the phenology of  
196 *Anacyclus clavatus* achene germination was the main factor affecting postdispersal life-history traits  
197 related to competitive ability and reproductive success.

198 In addition, the study of [24] showed a high phenological diversity for the four phenological patterns  
199 (buds, flowers, fruits and seeds) among fifteen leguminous plant species growing in Amritsar.

## 200 3.2. Morphology study

### 201 3.2.1. Study of vegetative part

202 It allows us to obtain, on the one hand, the resemblances, the objective of the typology of organs in  
203 the sense of a unity of organization and, on the other hand, the differences : the possible variations  
204 around the type : it's the comparative morphology. It seems useful to draw up a comparative table of  
205 the morphological characteristics of the 3 species studied (Table 1).

206 **Table 1:** Main distinctive characteristics of 3 species studied.

Species	NR	Leafs	Flowers	Akenes	P <sub>100</sub> A in mg
<i>Anacyclus clavatus</i>	T+5	Dark green Bipennatized	White ligulated flowers	Beige	45.23
<i>Chamaemelum fuscatum</i>	T+5	Green Bipennatized	Flowers in yellow tubes	Brown to yellow	26.6

<b><i>Leucanthemum parthenium</i></b>	T+3	Greenish-yellowish divided into wide segments.	White ligulated flowers	Brown	0.99
---------------------------------------	-----	------------------------------------------------	-------------------------	-------	------

207

208 **NR:** number of ramifications, **P<sub>100</sub> A:** weight of 100 akenes.

209

#### 210 **The branching**

211 Branching is the development of axillary buds in shoots. The number of branches is counted from  
 212 the principal axis of the stem noted « T ». Two types of branching are found: a branching of type (T +  
 213 5) for the species: *Anacyclus clavatus* and *Chamaemelum fuscatum* at the mature plant stage and a  
 214 Tertiary branching (T + 3) for the species *Leucanthemum parthenium*.

#### 215 **The stem**

216 The main stem of *Anacyclus clavatus* and *Chamaemelum fuscatum* is often orthotropically  
 217 developed. Plagiotropic development is sometimes observed. While their ramification has a  
 218 plagiotropic development. The main stem and branch of *Leucanthemum parthenium* have a strictly  
 219 orthotropic development.

#### 220 **Leaves**

221 The leaves of *Anacyclus clavatus* are alternate, long, of average length equal to 3.19, short  
 222 petiolated, bipennatized, acute at the tip and dark green at maturity. The leaves of *Chamaemelum*  
 223 *fuscatum* are alternate, long, their mean length equal to 2.19, petiolate, bipennatized, containing a  
 224 pointed end and green in color at maturity. The leaves of *Leucanthemum parthenium* are inserted in  
 225 an alternate phyllotaxy. They are long, of average length equal to 4.36, petiolate, divided into narrow  
 226 segments and yellowish-green at maturity.

#### 227 **The inflorescences and the flowers**

228 The inflorescence of *Anacyclus clavatus*, *Chamaemelum fuscatum* and *Leucanthemum parthenium*  
 229 is a flower head containing two types of flowers : yellow flowers tubulated in the center and white  
 230 flowers ligated at the periphery. The flowers of the 3 species have the same floral biology, but show a  
 231 difference in floral structure. Indeed, the liguled flowers of *Chamaemelum fuscatum* are long and

232 beaked at the tip, while those of two other species are similar; they are short and more or less  
233 rounded.

234 The diameter of the receptacle varies from one species to another. It is  $0.65 \pm 0.02$  cm in  
235 *Leucanthemum parthenium*,  $0.67 \pm 0.05$  cm in *Chamaemelum fuscatum* and  $1.56 \pm 0.01$  cm in  
236 *Anacyclus clavatus*.

#### 237 **Fruit**

238 The fruits differ from one species to another. The fruit of *Anacyclus clavatus* (Figure 4) is an indehiscent  
239 akene, beige at maturity, of rectilinear shape to flattened cone. This akene is surrounded by two  
240 membranous wings, clear, very thin, parchment and truncated at the apex. In the case of an akene  
241 without these wings, the fruit appears mottled and has four longitudinal ribs.

242

243 The fruit of *Chamaemelum fuscatum* (Figure 5) is an indehiscent akene, very small, not marginated,  
244 flattened ovoid, raised by 3 ribs weak and finely striated. Their color is brown to yellow at maturity.

245 The fruit of *Leucanthemum parthenium* (Figure 6) is an indehiscent akene, very small, brown at  
246 maturity, glandular and surmounted by a very short membranous crown and crenate.

#### 247 **Weight of 100 akenes**

248 The mean weight of 100 akenes of *A. clavatus* is 45.23 mg, varies from 37.7 to 53.8 mg. For *C.*  
249 *fuscatum*, It is 26.63 mg and varies from 25.3 to 27.9 mg. An average weight of 9.96 mg was  
250 calculated in *L. parthenium*; for this species, the range of variation is 8.5 to 10.9 mg.

#### 251 **3.2.2. Analysis of morphological variability**

252 The evaluation of a collection of genetic resources is commonly based on the simultaneous  
253 examination of many populations for various morphological characters. In this context, data on the  
254 different morphological traits measured were:

- 255 • An analysis of variance with one classification criterion followed by a comparison of means.
- 256 • An estimate of the degrees of association between the different traits studied by the Pearson  
257 correlation coefficient [25].
- 258 • A principal component analysis (PCA) based on the derivation of orthogonal variables [26].

##### 259 **3.2.2.1. Analysis of variance**

260 The analysis of variance with one classification criterion (species effect) showed highly significant  
261 differences between the three species studied (Table 2) for the majority of the quantitative traits  
262 measured such as: Length of the longest branch (LLR), Length of the smallest branch (LPR), number  
263 of secondary branches (NRS), number of primary branches (NRP), mean leaf spine length (LMRF),  
264 average number of leaflets (NLL), mean diameter of the receptacle (DMR), length of the main root  
265 (LRP), mean length of the tertiary branch (LMRT), average length of secondary branch (LMRS),  
266 average length of primary branch (LMRP) and length of the main axis (LAP). This species effect is  
267 only significant for the number of leaves (NF). The difference between the three species is not  
268 significant for: The number of the principal axis head ligules (NLCAP), the average number of ligules  
269 per capitule (NML) and the number of tertiary branches (NRT). This result reflects a phenotypic  
270 heterogeneity between the 3 species studied, taking into account the measured parameters.

271 **Table 2:** Results of the variance analysis of the 17 morphological traits measured.

Characters	ddl/df	Average square	F <sub>obs</sub>	Pr > F
LAP	2	3730,630	68,058	< 0,0001 <b>HS</b>
LMRP	2	982,641	26,382	< 0,0001 <b>HS</b>
LMRS	2	862,412	52,589	< 0,0001 <b>HS</b>
LMRT	2	360,894	26,359	< 0,0001 <b>HS</b>
LRP	2	40,961	11,73	0,000 <b>HS</b>
NF	2	338256,13	5,355	0,011 <b>S</b>
DMR	2	2,701	108,846	< 0,0001 <b>HS</b>
<b>NLL</b>	2	150,633	75,039	< 0,0001 <b>HS</b>
LMRF	2	11,796	36,769	< 0,0001 <b>HS</b>
NI	2	30601,433	2,983	0,068 <b>NS</b>
NRP	2	185,633	14,312	< 0,0001 <b>HS</b>
NRS	2	14770	15,244	< 0,0001 <b>HS</b>
NRT	2	4548,433	0,867	0,432 <b>NS</b>
NML	2	226,9	1,258	0,3 <b>NS</b>
NLCAP	2	0,7	1,086	0,352 <b>NS</b>
LPR	2	15,74	22,619	< 0,0001 <b>HS</b>
LLR	2	935,217	8,415	0,001 <b>HS</b>

272

273 **ddl/df**: degree of freedom; **F<sub>obs</sub>**: F observed; **HS**: highly significant; **S**: significant ( $P < 0.05$ ); **NS**:  
 274 no significant ( $P \geq 0.05$ ).

275

276 **3.2.2.2. Comparison of means**

277 According to the results obtained by analyzing the differences between the means with a 95%  
 278 confidence interval, we distinguish 5 types of groups (Table 3). Comparison of means shows that *A.*  
 279 *clavatus* and *C. fuscatum* are distinctly different from *L. parthenium* for: the length of the main axis  
 280 (LAP), the mean length of the secondary branch (LMRP), the average length of the tertiary branch  
 281 (LMRT), Root length (LR), number of leaves (NF), number of primary branches (NRP) and number of  
 282 secondary branches (NRP).

283 *A. clavatus* is distinguished from *L. parthenium* and *C. fuscatum* for the mean diameter of the  
 284 receptacle (DMR), the length of the smallest branch (LPR) and the length of the longest branch (LLR).

285 In fact, the three species did not differ significantly in the mean diameter of the receptacle (DMR), the  
 286 length of the smallest branch (LPR) and the length of the longest branch (LLR).

287 The parameters discriminating the three species are: the average length of the primary branch  
 288 (LMRP), the mean number of leaflets per leaf (NMf) and the average length of the spine (LMRF). For  
 289 the number of inflorescence per plant (NI), *Anacyclus clavatus* is not significantly different from  
 290 *Chamaemelum fuscatum* or *Leucanthemum parthenium*. Therefore, *Anacyclus clavatus* and  
 291 *Chamaemelum fuscatum* are much alike for more than half the morphological characters studied.  
 292 Most of the highest averages of the morphological traits are observed in *Anacyclus clavatus*, while the  
 293 majority of the lowest averages are observed in *Leucanthemum parthenium* (Table 3).

294  
 295  
 296  
 297

298 **Table 3:** Comparison of means of the 3 species studied.

299

Traits	<i>Anacyclus clavatus</i>	<i>Chamaemelum fuscatum</i>	<i>Leucanthemum parthenium</i> <sup>300</sup>
LAP	19,8 B	20,71 B	53,7 A 301
LMRS	20,6 A	17,91 A	3,39 B
LMRT	12,12 A	12,5 A	1,91 B 302
LR	8,1 B	7,72 B	11,4 A
NF	629,5 A	524,5 A	271,7 B 303
NRP	11,4 B	11,9 B	19,1 A
NRS	39,6 A	29,6 B	100,6 A 304
DMR	1,56 A	0,67 B	0,65 B
LPR	3,21 A	1,4 B	0,8 B 305
LLR	46,69 A	29,97 B	29,91 B
NRT	53,7 A	37,3 A	79,6 A 306
NML	11,7 A	19,9 A	11,6 A
NLACP	13,3 A	13,4 A	12,9A 307
LMRP	36,12 A	24,34 B	16,42 C
NMf	15,6 A	10,9 B	7,9 C 308
LMRF	4,36 A	3,19 B	2,19 C
NI	116,5 A and B	82,4 B	190,6 A 309

310 **3.2.2.3. The Matrix of correlation coefficients**

311 The matrix of correlation coefficients between the characters studied (Table 4) shows: A positive  
312 correlation of the following traits: LMRP and LMRS correlate positively with each other and with all the  
313 parameters of LMRT, NF, DMR, **NLL**, LPR and LLR ; The parameters DMR and NMF correlate  
314 positively with each other and with LPR and LLR ; The character LAP is strongly correlated positively  
315 with the parameters LR, LMRF, NI, NRP and NRT ; LMRT correlates positively with NF and **NLL** ; LR  
316 correlates positively with the parameters **NLL**, LMRF, NI, NRP and NRS ; A positive correlation  
317 between the parameters NF and **NLL**, NRT and LLR ; A highly significant positive correlation between  
318 LMRF with NI, NRP and NRS ; NI correlates strongly with the parameters : NRP, NRS and NRT and  
319 weakly with LLR ; NRP is strongly correlated with NRS and weakly correlated with the characters  
320 NRT and LPR. A strong positive correlation is noted between NRS and NRT. A weak positive  
321 correlation is noted between NRT and LLR.

322 The LAP has a highly significant negative correlation with the parameters (LMRS, LMRT, **NLL**) and  
323 significant with the characters (LMRP, NF, DMR, LPR); LMRS and LMRT correlate negatively with  
324 LR, LMRF, NRP and NRS; **NLL** is significant negatively correlated with NRP and NRS; LR correlates  
325 positively and significantly only with **NLL**; DMR is significant negatively correlated only with NRP  
326 (Table 4). It is important to note that NLCA and NML are not correlated with any of the other  
327 characters and that LMRP is the most positively correlated with the other traits (Table 4).

328 **3.2.2.4. Principal component analysis**


329 The graphical representation of the individuals dispersion of the 3 species studied reveals a more or  
330 less homogeneous grouping of the species studied forming 3 clear groups (Figure 7).

331 Indeed, there is a slight overlap between the two groups: *Anacyclus clavatus* and *Chamaemelum*  
332 *fuscatum*, whereas, *Leucanthemum parthenium* group seems very far from the others. These results  
333 confirm those of the variance analysis which showed a strong resemblance between *Anacyclus*  
334 *clavatus* and *Chamaemelum fuscatum*.

335 It is also observed that the individuals of the species *Chamaemelum fuscatum* occupy a rather  
336 restricted part of the plane and are located entirely in the negative part of the two axes F1 and F2.

337 While, the individuals belonging to *Anacyclus clavatus* are scattered on the two axes (F1 and F2) with  
 338 a trend towards the positive values of the F1 axis (Figure 7).

339 Furthermore, individuals of *Leucanthemum parthenium* are the best dispersed on the 2 axes (F1 and  
 340 F2) with a tendency towards the negative values of F1 axis (Figure 7).



341

342 **Figure 7:** Dispersal of the 3 species individuals in the plan defined by the axis 1 and 2 of the ACP.

343



Table 4: Matrix of correlation coefficients of the different morphological parameters.

Traits	LAP	LMRP	LMRS	LMRT	LR	NF	DMR	NMF	LMRF	NI	NRP	NRS	NRT	NML	NLCAP	LPR	LLR
LAP	1																
LMRP	-0.536	1															
LMRS	-0.810	0.842	1														
LMRT	-0.766	0.707	0.918	1													
LR	0.607	-0.270	-0.572	-0.544	1												
NF	-0.388	0.797	0.763	0.764	-0.281	1											
DMR	-0.451	0.679	0.496	0.315	-0.290	0.271	1										
NMF	-0.670	0.691	0.662	0.522	-0.511	0.423	0.798	1									
LMRF	0.790	-0.266	-0.629	-0.677	0.451	-0.269	-0.048	-0.283	1								
NI	0.532	0.220	-0.123	-0.143	0.417	0.377	-0.176	-0.195	0.523	1							
NRP	0.803	-0.291	-0.579	-0.594	0.575	-0.135	-0.410	-0.572	0.673	0.528	1						
NRS	0.826	-0.119	-0.494	-0.461	0.603	0.014	-0.314	-0.455	0.701	0.872	0.774	1					
NRT	0.410	0.329	-0.007	-0.004	0.303	0.462	-0.104	-0.080	0.373	0.946	0.473	0.798	1				
NML	-0.130	0.052	0.090	0.095	0.014	0.269	-0.160	-0.031	-0.136	0.025	0.171	-0.172	0.048	1			
NLCAP	-0.179	0.282	0.282	0.325	-0.357	0.254	0.153	0.161	-0.267	0.006	-0.058	-0.016	0.075	-0.020	1		
LPR	-0.529	0.576	0.492	0.378	-0.385	0.289	0.762	0.787	-0.142	-0.224	-0.478	-0.387	-0.153	-0.114	0.247	1	
LLR	-0.184	0.868	0.597	0.465	-0.051	0.722	0.541	0.485	0.088	0.495	0.058	0.248	0.526	-0.094	0.194	0.396	1

352 In fact, the morphological study of [27] revealed that the analysis of variance showed variations  
353 among the 33 accessions of *Ricinus communis* L. from Andaman and Nicobar Islands for all the 18  
354 traits studied. This work reveals also that plant height exhibited high significant positive correlations  
355 with the number of nodes on the main stem. In addition, The cluster analysis based on morphological  
356 traits grouped the 33 accessions of *Ricinus communis* L. into two major clusters [27].

357 Furthermore, the study of [28] was found a significant amount of genetic variability for all the twenty  
358 morphological parameters studied among safflower germplasm. In addition, this work reveals that  
359 seed yield plant had high significant and positive correlation with branches plant, capitulum plant,  
360 seeds capitulum and 100 seed weight. Furthermore, the hierarchical cluster analysis based on agro-  
361 morphological parameters divided 121 accessions of safflower into 5 main clusters [28].

362 The morphological study of [29] in rice varieties showed high phenotypic variability ( $P < 0.0001$ ) for  
363 the characters: leaf length and leaf width, primary branching, maturity and grain thickness. In addition,  
364 this work revealed a positive and strong correlation (0.77) between the height at maturity and leaf  
365 length. The cluster analysis of this morphological study based on Euclidian distances between the 98  
366 genotypes of Rice has allowed identifying three major clusters.

#### 367 **4. Conclusion**

368 The results of the phenological study show that the 3 species studied have distinct phenologies. The  
369 longest phenological cycle is observed for *Leucanthemum parthenium*. The variance analysis showed  
370 significant differences to highly significant for the majority of the traits studied. The comparison of  
371 means and the principal component analysis reveals that *Anacyclus clavatus* and *Chamaemelum*  
372 *fuscatum* form a single group for most of the traits measured while *Leucanthemum parthenium* is  
373 clearly distinct from these two species. Furthermore, this study allowed us to validate the  
374 morphological and phenological approach as tools for selection of suitable genotypes.

375

#### 376 **References**

- 377 1. Mulik NG, Bhosale LJ. Flowering phenology of the mangroves from the West Coast of  
378 Maharashtra. J Bom Nat Hist Soc. 1989 ; 86 (3): 355 - 359.

- 379 2. Suresh HS, Sukumar R. Vegetative phenology of tropical montane forests in the Nilgiris, South  
380 India. Jour Natio Sci Founda Srilan. 2011; vol-39 : 333 - 343.
- 381 3. Rout GR, Sahoo DP, Aparajita S. Studies on inter and intrapopulation variability of *Pongamia*  
382 *pinnata*: a bioenergy legume tree. Crop Breed Appl Biotechnol. 2009 ; 9: 268 - 273.
- 383 4. Zhang G, Song Q, Yang D. Phenology of *Ficus racemosa* in Xishungbanna, South West China.  
384 Biotropica. 2006 ; 38: 334 - 341.
- 385 5. Martins SR, Vences FJ, Miera LE, Barrosa MR, Carnide V. RAPD analysis of genetic  
386 diversity among and within Portuguese landraces of common white bean (*Phaseolus vulgaris* L.). Sci  
387 Horti. 2006; 108: 133-142.
- 388 6. Balkaya A, Ergun A. Diversity and use of pinto bean (*Phaseolus vulgaris*) populations from  
389 Samsun, Turkey. N. Z. J. Crop Horti Sci. 2008 ; 36 : 189-197.
- 390 7. Shinwari S, Akbar F, Rabbani MA, Mumtaz AS, Shinwari ZK. Evaluation of genetic diversity  
391 in different genotypes of *Eruca sativa* from Pakistan by SDS-PAGE Analysis. Pak Jour Botan. 2013;  
392 45: 1235 - 1240.
- 393 8. Ghafoor A. Genetic diversity and gene-action in *Vigna mungo* based on morphological and  
394 biochemical markers. Ph D thesis: Quaid-i-Azam University. Islamabad, pp. 192; 1999.
- 395 9. Demol J, Baudoin JP, Louant BP, Maréchal R, Mergeai G, Otoul E. Plant breeding:  
396 application to the main species grown in tropical regions. Ed presses of Gembloux, pp. 583; 2001.
- 397 10. Abdi A, Bekele E, Asfaw Z, Teshome A. Patterns of morphological variation of Sorghum  
398 (*Sorghum bicolor* (L.) Moench) landraces in qualitative characters in North Showa and South Welo,  
399 Ethiopia. Hereditas. 2002; 137: 161-172.
- 400 11. Fufa H, Baenziger PS, Beecher BS, Dweikat I, Graybosch RA, Eskridge KM. Comparison  
401 of phenotypic and molecular marker-based classifications of hard red winter wheat cultivars.  
402 Euphytica. 2005 ; 145: 133-146.

- 403 12. Mal TK, Doust JL. Phenotypic plasticity in vegetative and reproductive traits in an invasive  
404 weed, *Lythrum salicaria* (Lythraceae), in response to soil moisture. Amer Jour Botan. 2005 ; 92 (5) :  
405 819 - 825.
- 406 13. Ducombs G. Lactones sesquiterpéniques et plantes. Rev Fr Allergol. 1999 ; 39 (4) : 295- 298.
- 407 14. Bonnier G. La grande flore, Tome 3. Ed. Berlin, pp. 676; 1990.
- 408 15. Pottier AG. Flore de la Tunisie: Angiospermes-Dicotylédones, Gamopétales. Programme Flore et  
409 Végétation Tunisienne, pp. 651-1189 ; 1981.
- 410 16. Schweizer D, Ehrendorfer F. Giemsa banded Karyotypes, Systematics, and Evolution in  
411 *Anacyclus* (Asteraceae-Anthemideae). Plant Syst Evol. 1976 ; 126: 107- 148.
- 412 17. Le Floc'h E. Contribution à une étude ethnobotanique de la flore Tunisienne. Programme Flore  
413 et Végétation tunisienne : Ministère de l'Enseignement Supérieur et de la recherche Scientifique, pp.  
414 387 ; 1983.
- 415 18. Vogt R, Oberprieler C. Chromosome numbers of North African phanerogamsVIII. More counts  
416 in Compositae. Willdenowia. 2008; 38: 497- 519.
- 417 19. Ameddah S, Menad A, Dendougu H, Meraihi Z, Benayache S, Benayache F. Studies in  
418 Hepatocyte activity and the structure relationships and Luteolin Derivatives from *Chrysanthemum*  
419 *fuscatum*. Ecam. 2007; 4(S1): 55- 58.
- 420 20. Judd WS, Cambell CS, Kellogg EA, Stevens P . Botanique systématique. 1<sup>ère</sup> Edition, pp.  
421 612 ; 1999.
- 422 21. Besharati-Seidani A, Jabbari A, Yamini Y, Saharkiz MJ. Rapid Extraction and analysis of  
423 volatile organic compounds of Iranian feverfew (*Tanacetum parthenium*) using headspace solvent  
424 microextraction (HSME), and gas chromatography / mass spectrometry. Flav Fragr Jour. 2006; 21:  
425 502 - 509.
- 426 22. Castleman M. Les plantes qui guérissent: Le guide le plus complet sur le pouvoir curatif des  
427 remèdes offerts par la nature. Ed. Rodale, pp. 518 ; 1991.

- 428 23. Afonso A, Castro S, Loureiro J, Mota L, De Oliveira JC, Torices R. The effects of achene  
429 type and germination time on plant performance in the heterocarpic *Anacyclus clavatus* (Asteraceae).  
430 Amer Jour Botan. 2014 ; 101 (5) : 1- 7.
- 431 24. Kaur G, Pal Singh B, Kaur Nagpal A. Phenology of Some Phanerogams (Trees and Shrubs) of  
432 Northwestern Punjab, India. Jour Botan. 2013; Article ID 712405: 1- 10.
- 433 25. Clifford HT, Stephenson W. An introduction to numerical classification . Academic Press. New  
434 York, USA; 1975.
- 435 26. Hair JF, Anderson RE, Tatham RL, Black WC. Multivariate data analysis. Macmillan. New  
436 York, USA; 1992.
- 437 27. Kanti M, Anjani K, Usha Kiran B, Vivekananda K. Agro-morphological and Molecular  
438 Diversity in Castor (*Ricinus communis* L.) Germplasm collected from Andaman and Nicobar Islands,  
439 India. Czech J Genet Plant Breed. 2015; 51 (3): 96 - 109.
- 440 28. Arif M, Hussain I, Rabbani MA, Ali S, Ali N, Khan SM, Zaheer Tanoli MT, Raza H.  
441 Assessment of genetic diversity among safflower germplasm through agro-morphological traits. Intern  
442 Jour Biosci. 2016; Vol 9: 1 - 11.
- 443 29. Moukoumbi YD, Sié M, Vodouhe R, Ndri B, Toulou B, Ogunbayo SA, Ahanchede A.  
444 Assessing phenotypic diversity of interspecific rice varieties using agro-morphological  
445 characterization. Jour Plant Breed Crop Sci. 2011 : 74 - 86.
- 446
- 447