

2
3 **Incidence of Root-knot nematodes in Some Selected**
4 **Vegetable Crops in Nukkai Irrigation Field Jalingo,**
5 **Taraba State, Nigeria.**

6 **ABSTRACT**

7 A study to determine the incidence of root-knot nematode (*Melodogyne* spp) in some selected vegetables
8 was conducted in Nukkai Irrigation Field of Jalingo, Nigeria. Samples of Okra (*Hibiscus esculentus*),
9 Spinach (*Amaranthus* spp) and Sorrel (*Hibiscus sabdariffa*) were collected at 2-weeks old and at
10 maturity (flowering stage). Their roots were cut off and nematodes (*Melodogyne* spp) were extracted
11 using the Baermann method and identified using the female perineal pattern manual. The results showed
12 that *Melodogyne* spp are incident in the study area as 248 (34.44%) stands out of the 720 stands studied
13 were found to be infested by root-knot nematode. The results also showed that two *Melodogyne* spp, *M.*
14 *javanica* and *M. incognita* were discovered, and had a total number of 535 individuals. *M. javanica* (345)
15 were predominantly higher than *M. incognita* (190). In all stages of sampling, *Hibiscus esculentus* proved
16 to be more susceptible to infestation, followed by *Amarantus* spp while the least infested was *Hibiscus*
17 *sabdariffa* and *M. javanica* appeared to be more abundant than *M. incognita* in the study area.

18 *Keywords:* *Amaranthus* , *Hibiscus* , Jalingo, *Melodogyne*, *Nigeria*, Nukkai.

19 **INTRODUCTION**

20 Nematodes are soil dwelling organisms that constitute one of the largest animal phyla in the
21 world, with over half a million species known. It has been estimated that four out of every five living
22 animals on this planet are nematodes [1]. Nematodes can live as obligate parasites of plants and animals
23 and can also alternate a parasitic life, with a free living life style or can be strictly free [2]. Despite the
24 diversity in their life style and habitat, all nematodes are morphologically, anatomically and
25 developmentally similar [3].

26 Root-knot nematodes (*Meloidogyne* spp) are minute worm-like animals that are very common in
27 the soil and can invade roots of plants. They have a wide host range, and cause problems in many
28 annual and perennial crops; they occur throughout the world infecting all major crops and causing
29 substantial reduction in yield and quality [4]. There are four important species of the genus; *Meloidogyne*
30 *arenaria*, *Meloidogyne hapla*, *Meloidogyne incognita* and *Meloidogyne javanica* which are considered as

31 the most economically important species responsible for causing more damage to agricultural produce [5,
32 6].

33 Vegetables are much more important commodities needed in high demand due to their nutritive
34 value for balanced nutrition in many areas of the world but because of root-knot nematode (*Meloidogyne*
35 spp), it is increasingly very difficult and sometimes impossible to grow important vegetable in the tropics
36 and semi tropical countries [2]. In Africa, the yield of vegetables is relatively lower, for which there are
37 many constrains including prevalence of disease caused by different pathogens [7]. Among various
38 pathogens responsible for the low yield, root-knot nematodes are of considerable economic importance
39 [8] and can cause an annual loss of 22% in the tropics [9]. In addition, these parasites also interact with
40 other disease causing organisms to produce a disease complex [10]; break down resistance against other
41 pathogens and reduce plant tolerance to environmental stress [11,12].

42 It is against this background that this study was designed to determine the incidence of root-knot
43 nematodes of some vegetables cultivated in Nukkai Irrigation Fields of Jalingo, Taraba State.

44 **MATERIALS AND METHODS**

45 *Study Area*

46 Nukkai Irrigation Field is an irrigation farming field in Jalingo, Taraba State, which has an area of
47 about 20 hectares. Jalingo is located at latitudes 8⁰47' to 9⁰ 01'N and longitude 11⁰ 09' to 11⁰30'E. Nukkai
48 Irrigation Field is located immediately after the Nukkai Bridge south of the Jalingo-Wukari main road in the
49 city of Jalingo (Figure 1).

50 *Sample Collection*

51 All samples were collected using the method adopted by Anwar and McKenry [13]. Samples of
52 Okra (*Hibiscus esculentus*), Spinach (*Amaranthus* spp) and Sorrel (*Hibiscus sabdariffa*) were randomly
53 collected at different growth stages, i.e. seedling stage and matured stage. For each vegetable species,
54 seedling was collected at random when they were aged 2-weeks old and the matured vegetables were
55 also collected immediately they started flowering. All samples collected were washed and cleaned of soil
56 debris. The roots were cut off and kept in the fridge in a plastic container.

57 *Extraction of Nematodes*

58 Nematodes were extracted from roots of plants using the Baermann funnel extraction method as
59 described by Baermann [14]. In the set-up, a funnel was placed in a stand and filled with water until it
60 reached up to 1cm below the rim. A rubber tube was fixed at the mouth of the funnel whose end was
61 tightened using a clip. It was ensured that formation of air bubbles was avoided. The clip ensured it was
62 well closed and the rubber tube did not leak. The sieves with the sample were hung in the funnel so that

63 the sample was totally submerged, without touching the bottom of the funnel. Nematodes crawled out of
64 the sieve into the water and settled. After a period of 16-72 hours, the nematodes suspension was
65 trapped by opening the squeezer clip, regularly tapping and adding water to increase nematodes vitality.

66 *Identification and Counting of Nematodes*

67 Nematodes were identified using the female perineal pattern method manual or guide by [15].
68 During identification, nematodes were counted directly under microscope using the counting dish.

69 *Statistical analysis*


70 The collected data was analyzed using descriptive statistics of means, percentages and chi-
71 square.

72

73

74

75



76
77 **Figure1:** The Map of Jalingo showing the study area (Nukkai irrigation field).

78 **RESULTS**

79 *Incidence of Infestation*

80 The incidence of infestation of plant by root knot nematodes is shown in Table 1. The result
 81 revealed that plant parasitic nematodes are incident in the field of study as they are found in all the
 82 studied vegetable. Out of the 120 stands from the three (3) plots sampled, Okra (*Hibiscus esculentus*)
 83 had the highest infestation of 60 (50.0%), followed by Spinach (*Amaranthus spp*) which had a total
 84 infestation of 52 with an incidence of 43.3% while the least infestation (30) was recorded in Sorrel
 85 (*Hibiscus sabdariffa*) with a total incidence of 25.0%. Chi-square test also revealed that there was a
 86 significant difference ($p>0.05$) between the total number of infested and uninfested stands studied (Table

87 1). Across all plots, plot A showed highest infestation, with an incidence rate of 42.5%, followed by plot B
88 (40.0%), while the least infestation was observed in plot C (35.8%).

89 The incidence of infestation of matured stage of the vegetable plants by root knot nematodes is
90 presented in Table 2. Matured Spinach (*Amaranthus* spp) had the highest total infestation of 54 stands
91 out of the 120 stands from the three (3) plots sampled with an incidence rate of 45.0%, followed by Okra
92 (*Hibiscus esculentus*) which had a total infestation of 51 with an incidence of 42.5% while the least
93 infestation of 38 was recorded in Sorrel (*Hibiscus sabdariffa*) with a total incidence of 31.7%. Chi-square
94 test showed that there is a significant difference ($p>0.05$) between the total number of infested and
95 uninfested stands of all the vegetable plants (Table 2). The results also revealed that plot A had the
96 highest infestation rate (40.0%) followed by plot B (39.2%) while the least infestation was recorded in plot
97 C (35.8%), as shown in Table 2.

98 **Table 1:** Incidence of Plant Parasitic Nematodes at Seedling Stage in Nukkai Irrigation Field, Jalingo (2016)

Plant species	Plot A		Plot B		Plot C		Total	
	+ve	-ve	+ve	-ve	+ve	-ve	+ve	-ve
<i>Amaranthus</i> spp	19(47.5%)	21(52.5%)	13(32.5%)	27(67.5%)	20(50.0%)	20(50.0%)	52(43.3%)	68(55.7%)
<i>Hibiscus esculentus</i>	20(50.0%)	20(50.0%)	25(62.5%)	15(37.5%)	15(37.5%)	25(62.5%)	60(50.0%)	60(50.0%)
<i>Hibiscus sabdariffa</i>	12(30.0%)	28(70.0%)	10(25.0%)	30(75.0%)	08(20.0%)	32(80.0%)	30(25.0%)	90(75.0%)
Total	51(42.5%)	69(57.5%)	48(40.0%)	72(60.0%)	43(35.8%)	87(64.2%)	142(39.4%)*	218(60.6%)*

99 Note: Values with asterix are statistically significant ($\chi^2_{=0.05}$)

100

101 **Table 2:** Incidence of Plant Parasitic Nematodes at Maturated Stage in Nukkai Irrigation Field, Jalingo (2016)

Plant species	Plot A		Plot B		Plot C		Total	
	+ve	-ve	+ve	-ve	+ve	-ve	+ve	-ve
<i>Amaranthus</i> spp	20 (50.0%)	20 (50.0%)	19 (47.5%)	21 (52.5%)	15 (37.5%)	25 (62.5%)	54 (45.0%)	66 (55.0%)
<i>Hibiscus esculentus</i>	18 (45.0%)	22 (55.0%)	16 (40.0%)	24 (60.0%)	17 (42.5%)	23 (57.5%)	51 (42.5%)	69 (57.5%)
<i>Hibiscus sabdariffa</i>	10 (25.0%)	30 (75.0%)	12 (30.0%)	28 (70.0%)	16 (40.0%)	24 (60.0%)	38 (31.7%)	82 (68.3%)
Total	48 (40.0%)	72 (60.0%)	47 (39.2%)	73 (68.8%)	48 (40.0%)	72 (60.0%)	143 (39.7%)	299 (60.3%)

102 Note: Values with asterix are statistically significant ($\chi^2_{=0.05}$)

103 Table 3, showed the abundance of *Melodogyne* spp extracted throughout the study. Out of the
 104 720 stands of vegetables studied across the three plots, 285 stands were found to be positively infested.
 105 A total of 535 *Melodogyne* spp of nematode were extracted, out of which *M. incognita* recorded a total of
 106 190 individuals across the three plots with the highest (78 individuals) in Plot B, while the least in Plot C
 107 (51). *M. javanica* proved to be more abundant in the study area which had a total of 345 individuals
 108 across the three Plots, with Plot A having the highest (115 individuals), followed by Plot C (127
 109 individuals), and the least (103 individuals) was recorded in Plot B (Table 3). The chi-square analysis also
 110 revealed that there was no significant difference ($p>0.05$) between the total number of nematodes
 111 extracted across the three plots (Plots A, B and C) but there was a significant difference ($p>0.05$)
 112 between the total number of *M. incognita* and *M. javanica* extracted.

113 **Table 3:** Abundance of *Melodogyne* spp Collected During the Studies (2016)

Nematode spp	Plot A	Plot B	Plot C	Total
<i>Melodogyne incognita</i>	61	78	51	190*
<i>Melodogyne javanica</i>	115	103	127	345*
Total	176	181	178	535

114 Note: Values with asterisk are statistically significant ($\chi^2_{=0.05}$)

115 DISCUSSION

116 Vascular feeders have become sedentary endoparasites [16]. They usually damage their hosts
 117 by redirecting large amount of energy and nutrients from normal cellular activities into their special
 118 feeding sites and developmental activities [16]. They also alter tissues and then disrupt the vascular
 119 system, which leads to prevention of water transportation and dissolved nutrients upward to the shoots by
 120 the xylem and also the translocation of photosynthesized materials to other parts of the plants by the
 121 phloem [17]. Roots severely galled by these nematodes could predispose the plants to root rots which
 122 could subsequently lead to shorter life span of the plants [18]. Such galled tissues could become
 123 succulent, poorly protected from invasion by other disease pathogens and nutrient rich food substance
 124 that can easily be colonized by fungi [18]. Anwar and Mckenry [19] reported that damages by root-knot
 125 nematodes could lead to poor growth, decline in quality and yield, and reduced resistance to stresses like
 126 drought and disease.

127 As revealed in this study, it is important to note that, there is an increase in number of infestation
 128 in each vegetable crop as it ages. This probably signifies that as the plants aged and increases in size,
 129 more spaces are created for more nematodes to infest and colonizes the vegetable crops. In all stages of
 130 sampling, *Hibiscus esculentus* proved to be more susceptible to infestation, followed by *Amarantus* spp
 131 while the least infested was *Hibiscus sabdariffa*. This probably might be due to selection of host by the
 132 parasites which favored the attack or infestation of *Hibiscus esculentus* than the rest of the vegetables. It

133 could also be that, Sorrel (*H. sabdariffa*) had the least number of infested stands probably because of its
134 sour nature which hinders the nematodes from infesting it more.

135 In terms of abundance, this study identified two species of root-knot nematodes from the study
136 area, *Meloidogyne incognita* and *Meloidogyne javanica*. *M. javanica* appeared to be more abundant than
137 *M. incognita* and this could probably be as a result of the nature of the soil in the study area, which might
138 be more suitable for the survival of *M. javanica* than *M. incognita*. It could also be that *M. javanica* is more
139 resistant to stress and other environmental challenges than *M. incognita* which would generally give *M.*
140 *javanica* reproductive advantage over its cousin (*M. incognita*). This study is in conformity with the works
141 of Haroon and Zylstru [3], Nagesh *et al.*, [20] and Kayami *et al.*, [9] who all reported that root-knot
142 nematodes, *M. javanica* is abundantly available than the other species. The result of this study also
143 agrees with the work of many researchers among which are Khan [21], Verdejo-Lucas *et al.* [22], Anwar
144 and McKenry [13] and Kayami *et al.* [9] who all reported availability or incidence and abundance of root-
145 knot nematode in different regions of the world.

146 REFERENCE

- 147 1. Platt HM. *The Phylogenetic Systematics of free living nematode*. The Ray Society, London, 1994;
148 pp.5.
- 149 2. Babatola OO. A Complete Characterization of the Four Most Common Species of Root-Knot
150 Nematode, *Meloidogyne* with pictorial key. IM Publication Raleigh, N.C.U.S.A, 2003; pp. 20-22.
- 151 3. Haroon SA and Zylstru C. Rapid Identification of Genetic Relationship of *Meloidogyne incognita*
152 Population by Polymerase Chain Reaction RAP Markers. Egyptian Journal of Agronomatology,
153 2003; 32(4): 18-20.
- 154 4. Sasser JN. Economic Importance of *Meloidogyne* in Tropical Countries. In: Lamberti, F. and
155 Taylor C.E. (Eds). *Root-Knot Nematodes (Meloidogyne spp): Systematics Biology and*
156 *Control*. Academic Press, New York, 1987; pp 359–374.
- 157 5. Wilson JA. The Incidence of Root-Knot Nematodes *Meloidogyne arenaria*, *Meloidogyne*
158 *incognita* and *Meloidogyne javanica* on Vegetables. Plant Disease Journal, 2012; 22: 12–18.
- 159 6. Babatola OO and Ogurnunmare OB. Interaction between *Meloidogyne incognita* and *Fusarium*
160 *oxysporum* on Selected Bean Genotypes. Journal of Nematology, 2012; 45 (2): 20-24.
- 161 7. Ahmed R. Root-Knot of Potato in West Pakistan. Plant Disease Reporter, 2012; 57: 505-510.
- 162 8. Hassan SS and Nagesh M. Incidence of Root-Knot Nematode, *Meloidogyne incognita* and
163 *Meloidogyne hapla* in Slovenia. Acta Agriculturae Slovenica, 2011; 451 – 474.
- 164 9. Kayami MZ, Mukhtar T and Hussain MA. Association of Root- Knot Nematodes (*Meloidogyne*
165 spp) With Cucumber in the Pothowar Region of Punjab Province of Pakistan. International
166 Journal of Biotechnology, 2012; 9 (1-2): 23 – 29.
- 167 10. Begun GP. Geographical Distribution and Infestation of Plant Parasitic Nematodes *Fusarium*
168 *oxysporum* on selected Bean Genotypes. Journal of Nematology, 2012; 45(1): 231-233.

- 169 11. Wheeler TA, Hake KD and Dever JK. Survey of *Meloidogyne incognita* and *Thielaviopsis basicola*:
170 Their Impact on Cotton Fruiting and Producers Management Choices in Infested Fields. Journal
171 of Nematology, 2000; 32(4): 576 – 583.
- 172 12. Barbosa DH, Viera SG, Souza RM and Silva CP. Survey of Root-Knot Nematode *Meloidogyne*
173 spp. in Coffee plantations in the state Rio de Janeiro, Brazil. Nematologia Brasileira, 2004; 28(1):
174 43 – 47.
- 175 13. Anwar SA and McKenry MV. Incidence and Population Density of Plant-Parasitic Nematodes
176 Infecting Vegetable Crops and Associated Yield Loss in Punjab, Pakistan. Pakistan Journal of
177 Zoology, 2012; 44(2): 327 – 333.
- 178 14. Baermann G. *Baermann method of root-knot nematode extraction. Methods and Techniques for*
179 *Nematology*, North Carolina State University and state Agency for International Development,
180 Raleigh, 1948; pp 18-19.
- 181 15. Taylor DP Netscher C. An Improved Technique for Preparing Perineal Patterns of *Meloidogyne*
182 spp. Nematologica, 1974; 20(2): 107–270.
- 183 16. Anwar SA. Influence of *Meloidogyne incognita*, *Partrichodorus minor* and *Pratylenicus scribneri* in
184 root-shoot growth and carbohydrate partitioning in tomato. Pakistan Journal of Zoology, 1995; 27:
185 105 – 113.
- 186 17. Hajera H, Feroza N and Shahina F. Effect of Verm and Nematode Interaction on Some
187 Biochemical Parameters of Sunflower. *Pakistan Journal of Nematology*, 2009; 27: 193-201.
- 188 18. Abawi GS and Chen J. Concomitant Pathogen and Pest Interactions. In: Plant and Nematode
189 Interactions (eds. K. R. Barker, G. A. Pederson, and G. I. Windham), Agronomy Mongraph 36.
190 American Society of Agronomy, Madiston, WI,1988; pp 135 – 158.
- 191 19. Anwar SA and McKenry MV. Incidence and reproduction of *Meloidogyne incognita* on vegetable
192 crop genotypes. Pakistan Journal of Zoology, 2010; 42: 135-141.
- 193 20. Nagesh M, Hussaini SS and Chidanandaswamy BS. Incidence of Root-Knot Nematode,
194 *Meloidogyne incognita* on Gherkin (*Cucumis sativus*) and Yield Losses. Indian Journal of Plant
195 Production, 2005; 33:309-311.
- 196 21. Khan RM. Distribution of *Radopholus similis* in India, its Spread in New Regions and Analysis of
197 the Nematofauna of Banana Crop Pathosystem. Nematologia Mediterranea, 1999; 27(2):
198 239-246.
- 199 22. Verdejo-Lucas S, Ornat C, Sorribas FJ and Stcheigel A. Species of Root-Knot Nematodes and
200 Fungal Eggs Parasites Recovered From Vegetables in Almeria and Barcelona, Spain. Journal of
201 Nematology, 2002; 34(4): 405 – 411.