

Original Research Article

1
2
3
4
5
6

**Occurrence of Cassava Mosaic Disease Related to Agro-ecosystem in
Farmer's Fields located in Kongo Central Province, Democratic Republic of
Congo**

7 ABSTRACT

8 **Aim:** To assess the Cassava Mosaic Disease (CMD) pressure by analyzing its incidence,
9 severity and gravity, and to characterize agro-ecosystems where cassava farmers' fields are
10 established.
11

12 **Place and duration:** The study was conducted in three different localities (Mvuazi, Ndembo
13 and Pompage) in Kongo Central province, Democratic Republic of Congo, from June to
14 December 2016.
15

16 **Methodology:** One hundred and fifty farmers' fields randomly selected were investigated
17 during epidemiological survey, with 50 fields in each locality. In each field selected, 30
18 cassava plants randomly selected in a square of 10m x 10m were analyzed. The CMD
19 incidence, severity and gravity were collected, and agronomic and environmental factors
20 relative to cassava fields were analyzed.
21

22 **Results:** In general, CMD was observed in the three localities, with pressure depending
23 upon to localities and fields. The distance between two neighboring fields could vary from 5
24 to 35 or even 50m. Pathological parameters show significant difference ($P = .05$) among
25 fields for the same locality. The lowest pressure was recorded in Mvuazi locality (with 12.8%
26 for incidence, score 2 for severity, and 15% for gravity), while the highest pressure was
27 recorded in Pompage (with 20% for incidence, score 3 for severity, and 32% for gravity).
28 Data recorded on agro-environmental factors show that farmers of the three localities used
29 almost the same agricultural practices. Analysis of data reported suggest that the origin and
30 the type of cassava material cuttings used can play a principal role in the propagation and
31 development of CMD in most of cassava cultivation regions.
32
33

34 **Conclusion:** The results of the present study revealed that CMD was present in different
35 localities surveyed, and its pressure varies among localities, and from one field to another for
36 the same locality. Agricultural practices used by farmers can play an important role in the
37 propagation of CMD in different regions of cassava cultivation.
38

39 **Keywords:** *Cassava Mosaic Disease, Farmers' fields, Agro-Ecosystem, Kongo Central*
40 *province, DR-Congo.*

41 1. Introduction

42

43 Cassava (*Manihot esculenta* Crantz) is an important source of calories for thousands of people living
44 in sub-Saharan Africa [1, 2]. Plant with high potential and adapting to different environments [3],
45 cassava is however subject to the attacks of Cassava Mosaic Disease (CMD) which constitutes a
46 serious and persistent threat to the food security of populations mainly living of that food.

47

48 In Africa, various works of selection and improvement of cassava led to the development of varieties
49 containing acceptable agronomic and qualitative characteristics, and resistant to diseases such as
50 CMD and Cassava Bacterial Blight (CBB) [4, 5]. These varieties have often been introduced and
51 distributed in many regions to control the CMD pandemic. Despite these breeding, improvement and
52 extension efforts, it was observed that the CMD continues to spread with high incidence and severity
53 levels. One of causes that would be the basis for the CMD perpetuation in these regions is the low
54 adoption of improved varieties by farmers. Indeed, farmers felt that these varieties did not meet their
55 expectations and preferences [6, 7], which results in the widespread use of local varieties.

56

57 In the Democratic Republic of Congo (DRC), studies conducted on the evaluation of the CMD
58 pressure indicate that the cassava germplasm is susceptible to this viral disease. For examples, in
59 Yangambi region (Eastern province), Monde [8] found that the majority of local varieties showed
60 severe symptoms of the disease, compared with improved varieties. In Bukavu region (Sud Kivu
61 province), Bisimwa [9] noted that local cassava varieties grown in different agro-ecosystems were all
62 susceptible to various biotic diseases identified on cassava. In Gandajika (Eastern Kasai province),
63 Muengula-Manyi *et al.* [10] observed that local varieties grown by farmers are severely attacked by
64 CMD compared to improved varieties.

65

66 Various other scientific studies have shown that local cassava varieties are severely attacked than
67 improved varieties. Jeremiah & Kulembeka [11] mentioned that all local cassava varieties available
68 are susceptible to CMD. In many countries, CMD would have reached a high severity degree on
69 farmers' fields, which may reduce the yield of cassava tuberous roots. The level of CMD infection
70 varying from agro-ecological systems [12], and poor farming practices and marginal agro-
71 environmental conditions observed in farmers' fields are favorable for CMD development.

72

73 This study aimed to assess the CMD pressure by analyzing the disease incidence, severity and
74 gravity in Mvuazi, Ndembo and Pompage localities (in the Kongo Central province), and to
75 characterize agro-environmental factors where cassava farmers' fields are established.

76

77 **2. Materials and Methods**

78 **2.1. Sites Description and Field Sampling**

79

80 Epidemiological surveys were conducted in Mvuazi, **Ndembo** and Pompage localities (Kongo Central
81 province) in DRC. These regions fall within the Aw4 climate type according to Köppen classification
82 characterized **by** 4 months of dry season coupled 8 months of rainy season. Daily temperature
83 averages 22-24°C and can reach a maximum of 30°C. The average annual rainfall ranges around
84 1,522mm. The surveyed sites were characterized by the presence of savannah dominated by
85 herbaceous species such *Hyparrhenia diplandra*, *Mucuna* sp., *Panicum maximum* and *Pennisetum*
86 *purpureum*. In some places, it **is** observed ragged forest where dominated shrub species such **as**
87 *Lussonia angolensis* and *Hymenocardia acida*. According to Pauwels [13], soils of Kongo Central
88 region are varying types, and revealed the presence of sandy and clay soils.

89

90 Epidemiological surveys were conducted in cassava farmers' fields during the period from June to
91 December 2016. In each locality, 50 fields randomly selected were investigated. In each field
92 selected, 30 cassava plants randomly selected in a square of 10m x 10m were analyzed.

93

94 **2.2. Variables Studied**

95 **2.2.1. Pathological Variables**

96

97 During epidemiological investigations, pathological variables recorded were CMD incidence, severity
98 and gravity. The CMD incidence was **assessed** by the proportion of diseased plants compared to 30
99 plants analyzed. CMD severity symptom was assessed using a scale ranging from 1 to 5 described
100 by Hahn *et al.* [14], where 1 represents an asymptomatic cassava plant (apparently healthy) and 5 a
101 severely infected cassava plant with reduction of leaflets. The CMD gravity was assessed in each
102 diseased plant by the proportion of leaves with typical symptoms of the disease.

103

104 **2.2.2. Agronomic and Environmental Factors**

105

106 For each field surveyed, agronomic and environmental characteristics as described by Muengula-
107 Manyi *et al.* [10] were determined. They include field location, origin and type of cassava material
108 used, age of fields, topography of land, the practice of intercropping, type of crops mixed with
109 cassava, and the topping practice.

110

111

112 2.3. Data Analysis

113

114 Statistical analysis of data recorded was made possible through the R software and Statistix 8.0 (free
115 version). The recorded data were submitted to analysis of variance followed by multiple comparisons
116 by Tukey's HSD, to determine significant differences ($P = .05$) between the surveyed sites. CMD
117 incidence and gravity were previously submitted to a logarithmic transformation to base 10 (\log_{10}).
118 The comparison of means was made using the least significant difference test (LSD) at the 5%
119 probability.

120

121 3. Results

122 3.1. Incidence, Severity and Gravity of Cassava Mosaic Disease

123

124 Results obtained on CMD incidence, severity and gravity recorded in the 3 localities are reported in
125 Table 1.

126

127 Table 1. Incidence, severity and gravity of CMD recorded in Mvuazi, Ndembo and Pompage locality

128

Locality	Pathological variables recorded		
	Incidence (%)	Severity (scale 1 - 5)	Gravity (%)
Mvuazi	12.8 ^b	2	15 ^c
Ndembo	15.2 ^b	3	25 ^b
Pompage	20 ^a	3	32 ^a

129 *In the same column, means followed by the same letter are not significantly different at 5% of*
130 *probability.*

131

132 In general, CMD was present in all sites surveyed with levels of incidence, severity and gravity
133 varying between localities, and from one field to **another** in the same locality. There were significant
134 differences for disease incidence and gravity among fields for the three sites (Table 1). Overall, the
135 mean incidence for all fields surveyed was 16%, severity score was 2.6, and gravity equal to 24%.
136 Details for each locality revealed that the incidence of CMD was 12.8% in Mvuazi, 15.2% in Ndembo
137 and 20% in Pompage. The mean of CMD severity was equal to 2 in Mvuazi, and 3 in Ndembo and
138 Pompage, and the gravity was respectively equal to 15, 25 and 32%.

139

140

141 3.2. Agronomic and Environmental Characteristics of Fields Investigated

142

143 The results of different agronomic and environmental factors analyzed for each cassava field
144 prospected in Mvuazi, Kimpese and Pompage localities are reported in Table 2.

145

146 Table 2. Frequency (%) of cassava fields characteristics in 3 localities investigated in Kongo Central
147 region

Characteristics of fields	Localities		
	Mvuazi	Ndembo	Pompage
Field location			
Secondary forest	40	70	64
Savannah	60	30	36
Site topography			
Flat land	70	24	60
Land with slope	30	76	40
Origin of cassava material used			
Research center	90	-	-
Old field	10	100	100
Type of cassava material used			
Local	14	90	96
Improved	86	10	4
Age of field			
1 to 6 months	8	10	8
7 to 12 months	86	70	72
Older than 12 months	6	20	20
Intercropping practice			
Yes	15	90	85
No	85	10	15
Crop mixed with cassava			
Legume	15	35	45
Cereal	75	25	25
Vegetable crop	10	40	30
Topping practice			
Yes	66	68	70
No	34	32	30

148

149 3.2.1. Field location and site topography

150 Cassava fields investigated were established either in secondary forest or savannah. In the 3
151 localities, cassava crop grown in secondary forest represented 58%, while those established in
152 savannah represented 42%. Farmers' fields were established either on flat land or on land with slope.
153 It was observed that 51.3% of cassava crops were grown on flat lands and 48.6% on lands with slope.
154 Details of cassava fields location and site topography for each locality are described in Table 2.

155 **3.2.2. Origin and type of cassava material used**

156 Analysis of data reported in Table 2 revealed that 30% of farmers used cassava cuttings obtained
 157 from a Research Center, and 70% used cuttings obtained from their previous fields. Farmers used
 158 local or genetically improved cassava varieties. Local cassava varieties were grown in 66.6% of fields,
 159 while improved varieties were planted in 33.3% of fields.

160 **3.2.3. Age of field**

161 According on the date of cassava plantation, fields investigated were classified in 3 groups. The first
 162 group included 1 to 6 months old cassava field, the second group with 7 to 12 months, and the third
 163 group with fields older than 12 months (Table 2). Results obtained revealed that 8.6% of cassava
 164 fields were 1 to 6 months old, 76% were 7 to 12 months old, and 15.3% were older than 12 months.
 165 Details of the three groups for each locality are described in Table 2.

166 **3.2.4. Intercropping practice and type of crops mixed with cassava**

167 The results of this study revealed that cassava was generally grown in association with other crops
 168 such as legumes, cereal or vegetable crops. Analysis of these results indicated that 63.3% of cassava
 169 were mixed with other crops, while in 36.6% of cases, cassava crop was grown alone. In general,
 170 31.6% of cassava stands were grown in association with legumes (soybeans or beans), 41.6% with
 171 cereal (principally maize) and 26.6% with vegetable crops (sweet potatoes). Frequency of
 172 intercropping practice and crops mixed with cassava varied according to localities surveyed (Table 2).

173 **3.2.5. Topping practice**

174 It observed that field topping was generally practiced in the three localities surveyed. This suggest
 175 that cassava leaves are appreciated such an edible legume to meet household needs. Field topping
 176 was practiced in 68% of cassava stands, while no topping was reported in 32% of fields investigated.
 177

178 **4. Discussion**

179

180 This study revealed the presence of cassava mosaic disease (CMD) in different cassava farmers'
 181 fields located in Mvuazi, Ndembo and Pompage localities in Kongo Central province. Overall, CMD
 182 pressure assessed by the analysis of incidence, severity and gravity generally varies among localities,
 183 and from one field to another in the same locality.

184

185 The analysis of pathological variables reported in Table 1 revealed significant difference ($P = .05$)
 186 among localities. In general the CMD pressure was low in Mvuazi, whereas it was higher in Pompage
 187 locality. Results of this study show that CMD pressure is slightly lower compared to data presented in
 188 previous studies. In other regions of DRC, Sseruwagi et al. [15] revealed that the mean incidence of
 189 CMD during the period 2002-2003 was approximately 60%, with severity score equal to 3.1.
 190 According to Ariyo et al. [16] and Ntawuruhunga et al. [17], usually the incidence and severity of CMD
 191 vary according to the year, and from one region to another. Adjata et al. [18] mentioned that the level

192 of CMD incidence probably changes with the pressure of inoculum, which varies from one site to
193 another. Based on our findings and those of previous studies, it is clear that pathological parameters
194 (incidence, severity and gravity) fluctuate depending on several factors such as agronomic,
195 environmental and the pressure of inoculum prevailing in a region, as well as time or period of
196 observations. In addition, Sseruwagi et al. [15] mentioned that in some moderately resistant varieties,
197 symptoms of CMD can be localized or absent in some parts of cassava plant. Muengula-Manyi et al.
198 [10, 19] also observed on a diseased cassava plant that CMD symptoms did not necessarily
199 appeared on all leaves present on the plant. These observations explain the variability of level of
200 gravity recorded on the diseased plants surveyed.

201

202 Results reported in Table 2 indicate in general that farmers use almost the same agricultural practices
203 in the cultivation of cassava. Based on characteristics of fields surveyed, it appeared that 90% of
204 farmers located in Mvuazi use cuttings obtained from the Research center, while all farmers (100%)
205 founded in Ndembo and Pompage localities use cuttings from their previous fields. In addition, in
206 Mvuazi locality, 86% of cassava varieties planted are genetically improved, while 93% of cassava
207 material used in Ndembo and Pompage localities are local varieties (Table 2). These observations
208 may explain the low CMD pressure noted in Mvuazi compared to the two others localities. The results
209 of this study corroborate findings reported by Bisimwa [9] who observed in Bukavu region, that
210 cassava farmers' fields heavily attacked by biotic diseases were planted from local varieties.
211 According to Hillocks & Thresh [20], in some regions the lack of improved varieties orient farmers
212 towards large-scale use of local varieties; and the high frequency of use of local varieties could also
213 be explained by the quest characteristics valued by farmers and by the cost of improved cassava
214 varieties cuttings. In addition, the use of cuttings without health guarantee, taken from previous fields
215 may explain the permanent presence of CMD in some cassava production regions.

216

217 Although the CMD was observed in the three localities, its incidence was overall lower compared to
218 data reported by Sseruwagi et al. [15], while the severity score reported in these two studies was
219 similar. The low level of CMD incidence reported in this study may be due to the use of intercropping
220 practice and the type of crop mixed with cassava. Indeed, there was different crops intercropped with
221 cassava in the three localities surveyed. For example, in Mvuazi, cassava was mixed with cereal in
222 75% of fields investigated, while in Ndembo it was mixed with vegetable crop in 40%, and in
223 Pompage with legumes in 45% (Table 2). Our results corroborate observations made by Monde [8]
224 who observed in the Yangambi region, that incidence and severity of CMD were very lower in fields
225 where cassava was mixed with beans compared to fields where cassava was cultivated without crop
226 mixed.

227

228

229

230 **5. Conclusion**

231

232 The results of the present study revealed that CMD occurs in the three localities surveyed, and its
 233 level pressure varies between localities, and between different fields in the same locality. In general,
 234 results obtained showed that farmers used almost the same agricultural practices to establish their
 235 cassava fields. Origin and type of cassava material used **indeed** play a significant role in the spread
 236 and development of CMD. In the region where improved varieties were used, CMD pressure **was**
 237 **lower than where local varieties were used.** **The** low level of incidence, severity and gravity of CMD
 238 can be attributed to the use of intercropping practice and the type of crop mixed with cassava.

239

240 **References**

241

- 242 [1] Alabi OJ, Ogbe FO, Bandyopadhyay R, Lava Kumar P, Dixon AGO, Hughes Jd'A, Naidu RA.
 243 Alternate hosts of African cassava mosaic virus and East African cassava mosaic Cameroon
 244 virus in Nigeria. Arch. Virol. 2008; 153:1743-1747.
- 245 [2] FAOSTAT. FAO database. Food and Agriculture Organization of the United Nations, Rome, Italy,
 246 2009. <http://FAOSTAT.fao.org/site/339/default.aspx>.
- 247 [3] Kawano K, Daza P, Aruya A, Rios M, Gonzales WMF. Evaluation of cassava germplasm for
 248 productivity. Crop Sci. 1978; 18:372-380.
- 249 [4] Cach NT, Pérez JC, Lenis JI, Calle F, Morante F, Ceballos H. Epistasis in the expression of
 250 relevant traits in cassava (*Manihot esculenta* Crantz) for sub-humid conditions. J. of Heredity.
 251 2005; 96:586-592.
- 252 [5] Pérez JC, Ceballos H, Jaramillo, G, Morante N, Calle F, Arias B, Bellotti AC. Epistasis in cassava
 253 (*Manihot esculenta* Crantz) adapted to mild-altitude valley environment. Crop Sci. 2005;
 254 45:1491-1496.
- 255 [6] Nweke FI. Farm level practices relevant to cassava plant protection. Afr. Crop Sc. J. 1994; 2:563-
 256 582.
- 257 [7] Benesi IRM. Characterization of Malawian cassava germplasm for diversity, starch extraction and
 258 its native and modified properties. PhD Thesis. Department of Plant Sciences: Plant Breeding,
 259 Faculty of Natural and Agriculture Sciences, University of the Free State, South Africa; 2005.
- 260 [8] Monde G. Epidémiologie, diversité génétique et phylogéographie des virus de la mosaïque
 261 africaine du manioc dans la région de Yangambi en République Démocratique du Congo.

- 262 Thèse de doctorat, Université catholique de Louvain, Louvain-la-Neuve, Belgique; 2010.
263 French.
- 264 [9] Bisimwa E. Epidémiologie, diversité génétique, distribution et contrôle des virus de la mosaïque
265 africaine du manioc et de son vecteur (*Bemisia tabaci*) dans la région du Sud-Kivu en
266 République Démocratique du Congo. Thèse de doctorat, Université catholique de Louvain,
267 Louvain-la-Neuve, Belgique; 2011. French.
- 268 [10] Muengula-Manyi M, Nkongolo KK, Bragard C, Tshilenge-Djim P, Winter S, Kalonji-Mbuyi A.
269 Incidence, severity and gravity of cassava mosaic disease in savannah agro-ecological region
270 of DR-Congo: analysis of agro-environmental factors. *Am. J. of Plant Sci.* 2012; 3:512-519.
- 271 [11] Jeremiah SC, Kulembeka HP. Screening of local cassava varieties against cassava mosaic
272 disease and cassava green mite. In: Proceedings of the 13th ISTRC Symposium, Arusha,
273 Tanzania; 2007.
- 274 [12] Balyejusa KE, Bua A, Fregene M, Egwang T, Gullberg U, Westerbergh A. The effect of cassava
275 mosaic disease on the genetic diversity of cassava in Uganda. *Euphytica.* 2005; 146:45-54.
- 276 [13] Pauwels L. Catalogue des plantes cultivées au Jardin Botanique de Kisantu; 1977.
- 277 [14] Hahn SK, Terry ER, Leuschner ER. Breeding cassava for resistance to cassava mosaic disease.
278 *Euphytica.* 1980; 29:673-683.
- 279 [15] Sseruwagi P, Sserubombwe WS, Legg JP, Ndunguru J, Thresh JM. Methods of surveying the
280 incidence and severity of cassava mosaic disease and whitefly vector populations on cassava
281 in Africa: a review. *Virus Research.* 2004; 100:129-142.
- 282 [16] Ariyo AO, Dixon AGO, Atiri GI. Whitefly *Bemisia tabaci* (*Homoptera: Aleyrodidae*) infestation on
283 cassava genotypes grown at different Ecozones in Nigeria. *J. of Econ. Entom.* 2005; 98:611-
284 617.
- 285 [17] Ntawuruhunga P, Okao-Okuja G, Bembe A, Obambi M, Mvila AJC, Legg JP. Incidence and
286 severity of cassava mosaic disease in Republic of Congo. *Afr. Crop Sc. J.* 2007; 15(1):1-9.
- 287 [18] Adjata KD, Muller E, Peterschmitt M, Aziadekey M, Gumedzoe YMD. Incidence of cassava viral
288 diseases and first identification of East cassava mosaic virus and Indian cassava mosaic virus
289 by PCR in cassava (*Manihot esculenta* Crantz) fields in Togo. *Am. J. of Plant Physiol.* 2008;
290 3:73-80.

- 291 [19] Muengula-Manyi M, Mukwa L, Nkongolo KK, Tshilenge-Djim P, Winter S, Bragard C, Kalonji-
292 Mbuyi A. Assessing reactions of genetically improved and local cassava varieties to cassava
293 mosaic disease (CMD) infection in a savannah region of the DR-Congo. *Am. J. of Plant Sci.*
294 2013; 4:824-837.
- 295 [20] Hillocks RJ, Thresh JM. Cassava mosaic and cassava brown streak virus diseases in Africa: A
296 comparative guide to symptoms and etiologies. *Roots*. 2000; 7(1):1-8.