

**Iron Overload in the Root Environment of Rice (*Oryza sativa- L*)  
with a Miserable Nutrients Specification.**

**Abstract:**

In waterlogged soils under low pH, Fe<sup>2+</sup> availability increases and may reach toxic levels. The conditions of iron toxicity are quite well established over the World. The physiological effects of Fe<sup>2+</sup> within plant with subsequent plants' nutrients status are well documented in many literatures. Despite our current knowledge of the processes and mechanisms involved, iron toxicity, a function of growth conditions and the cultivar types remains as an important constraint to rice production, together with nutrients deficiency in the regional levels. To screen Fe tolerant cultivars and thus to evaluate the mechanisms involved in response to excess Fe, experiment was carried out with rice cultivars – Ranjit, Siyal Sali and Mahsuri, grown by developing artificial Fe toxic conditions in the soils of experimental pots applying different Fe<sup>2+</sup> concentrations (control- normal soil iron from rice field, +100, +200 and +300 ppm respectively). The study of plants' biochemical parameters confirmed the resistance of Mahsuri plants to Fe excess. With steady recovery of neutral pH and better chlorophyll contents, the root and shoot nutrients of Mahsuri were found to be higher compared to the plants of other two varieties when exposed to excess Fe. Except Fe and N in roots and shoots, the excess of Fe caused a negative impact on other nutrients in these vulnerable cultivars. Plants of Ranjit and Siyal Sali seem to be affected directly by Fe toxicity and also by the pseudo Fe toxicity, whereas Mahsuri seems to make use of the exclusion /and or avoidance mechanism to Fe overload.

Key Words: toxicity, nutrients, *Oryza sativa- L*, investigation, vulnerable

**Introduction:**

Iron is essential for plant growth and development<sup>1</sup>. In anaerobic acid soils, however, high concentrations of ferrous (Fe<sup>2+</sup>) ions may lead to Fe toxicity due to excessive Fe uptake<sup>2</sup>, which can result in yield reductions from 12 to 100 percent<sup>3</sup>. Excess Fe can be extremely toxic, as it reacts with oxygen and catalyses the production of free radical species. In waterlogged soil iron toxicity may disrupts or over expresses a number of metabolic routes can bring about nutrient disorder in rice cultivars. The expression of iron-toxicity symptom requires the excessive uptake of Fe<sup>2+</sup> by roots and its acropetal translocation via xylem flow into the leaves.

In North East India, a major portion of the rice is grown under lowland conditions<sup>4</sup>, and Assam is the highest rice producing state, where all rice is grown in waterlogged soils. Use of tolerant rice cultivars retaining better nutrients level is the best alternative and inexpensive technologies for rice production on Fe toxic soils of this area<sup>4,5</sup>.

Although several research work have been conducted worldwide to identify adaptive responses of different rice genotypes still rate of nutrients absorption (ionic competition for absorption) and their availability (in favourable oxidation states) under higher iron concentrations is a matter of debates. Under anaerobic conditions O<sub>2</sub> release from rice roots, oxidise Fe<sup>2+</sup> to polymeric oxy-hydroxide which coats on roots surface preventing the uptake of Fe<sup>2+</sup>, Mn<sup>2+</sup> and also acts as P reservoir<sup>6</sup>. Silveira et al (2007)<sup>6</sup> had also cited that except for Mn, no other nutrients seemed to have impaired uptake due to Fe toxicity in the vulnerable cultivar (I409 plants) and not in the resistant one (E108 plants).

The Fe<sup>2+</sup> concentrations in the soil solution that reportedly affect lowland-rice yields can range from 10 to >2000 mg per liter<sup>7</sup>. Iron-induced yield reduction is frequently associated

50 with a poor nutrient status of the soil<sup>8</sup>. Hence, many workers suggest that excess Fe<sup>2+</sup> may  
51 result in lower uptake of other essential nutrients due to chemical interactions in soil  
52 (ZnFe<sub>2</sub>O<sub>3</sub>, K-Fe complex). Sahrawat (2004, 2010)<sup>3,9</sup> has reported the possibility of “pseudo  
53 Fe toxicity” (Fe toxicity symptoms induced by nutrients deficiency) and “true Fe toxicity”  
54 (caused by excessive Fe<sup>2+</sup> uptake) in rice grown at higher iron concentration.

55 Plant’s tolerance to excess Fe might be the effect of Fe avoidance and/or tolerance to  
56 high internal Fe concentration. Such avoidance and/or tolerance capacity to Fe overload is a  
57 genotypic function<sup>6,10</sup>. To sum-up, the conditions of iron toxicity are quite well established all  
58 over the World. The physiological effects of Fe<sup>2+</sup> within the plant with subsequent plants  
59 nutrients status are well documented in many literatures. In spite of our current knowledge of  
60 the processes and mechanisms involved, iron toxicity remains an important constraint to rice  
61 production in regional level where selection of cultivars having the ability to maintain high  
62 levels of essential micro and macro nutrients under Fe toxic condition is a successful  
63 approach for lowland rice cultivation in acid soil. To screen Fe tolerant cultivars and thus to  
64 evaluate the mechanisms involved in response to excess Fe, experiment was carried out with  
65 rice cultivars – Ranjit, Siyal Sali and Mahsuri, grown by developing artificial Fe toxic  
66 conditions in the soils of experimental pots applying different Fe<sup>2+</sup> concentrations. In this  
67 work we studied the differential responses of three rice cultivars to iron excess by evaluating  
68 the influence of Fe nutrition on other nutrients uptake, their elemental concentrations in rice  
69 roots and shoots, to help the investigation of mechanisms involved in resistance to Fe  
70 toxicity.

71 **Materials and methods:**

72 An artificial Fe toxic conditions in the experimental pots were developed with soils  
73 collected from a rice field located at Titabor of state Assam, India (soil type-sandy clay loam,  
74 total soil iron 345ppm, pH 5.4, available phosphorus 18.1kg.ha<sup>-1</sup>, nitrogen 460kg.ha<sup>-1</sup>, potash  
75 127kg ha<sup>-1</sup> and organic carbon 1.2%). The experiments was conducted with three rice (*Oryza*  
76 *sativa*-L) varieties viz. Mahsuri, Ranjit (high yielding varieties) and Siyal Sali (traditional tall  
77 variety) and four different levels of Fe<sup>2+</sup> like control (normal soil without adding external  
78 Fe<sup>2+</sup>), +100 ppm, +200 ppm and +300 ppm in the form of FeSO<sub>4</sub>.7H<sub>2</sub>O were applied to the  
79 experimental pots. Treatments were replicated four times in a randomized block design  
80 (Fisher and Yates, 1957). Thirty days old seedlings of uniform vigour were transplanted at  
81 the rate of three seedlings per pot. 100 ml of Fe<sup>2+</sup> solutions of the said concentrations were  
82 added to the pots once a week after transplanting at an interval of seven days till panicle  
83 initiation stage. A uniform waterlogged environment was maintained with distilled water in  
84 the pots throughout the experimental period.

85 **pH record:** pH of soil solutions were recorded (from each pot) *in-situ* at an interval of seven  
86 days from days after transplanting (DAT) (n=5 for each pot) with the help of a digital pH  
87 meter.

88 **Total Leaf chlorophyll:** Extraction and estimation was done by spectrophotometric  
89 method<sup>12</sup>.

90 **Nutrients analysis:** The K content was determined flame photo-metrically from mineral  
91 solution obtained after tri-acid digestion<sup>13</sup>. P was estimated from mineral solution converting  
92 phosphate to phosphomolybdic acid and finally reducing with hydroquinone. The blue colour  
93 developed was measured in a spectrophotometer (Systronics UV-VIS Spectrophotometer  
94 118) at 660nm<sup>13</sup>. The total nitrogen in roots and shoots were determined by Micro-Kjeldahl’s  
95 method<sup>14</sup> with 0.5 g powdered sample after digesting with concentrated H<sub>2</sub>SO<sub>4</sub> and catalyst  
96 mixture. N content was determined by titrating the distillate with 0.1N HCl.

97 Mineral solution was prepared by digesting 1g dry samples in tri-acid mixture and  
98 extracted with concentrated nitric acid. Fe, Mn, Zn & Cu were determined by Atomic

99 Absorption Spectrophotometer (Chemito, AA 203D) from mineral solution using separate  
100 primary standard for each micro-nutrient<sup>14</sup>.

101 Statistical analyses of experimental data were carried out by using SPSS software.  
102 Analysis of variance was carried out to test the significance of treatment effect. F-test,  
103 coefficient of variance and critical difference were calculated by standard method<sup>15</sup>.

#### 104 **Results and Discussions:**

105 The pH of the growth medium has significant impact on the properties of soils and  
106 consequently on the nutrient uptake by crop plants. The pH of soil solution is thought to be  
107 best for plant growth when kept between 5.5 and 6.5. Plant growth in acid soils may be  
108 limited by pH-induced Fe<sup>2+</sup> toxicity as acidity increases the solubility of Mn and Fe in acid  
109 soils<sup>7</sup>. In such adverse pH condition plants suffer from ionic imbalance through a competition  
110 between the similarly charged ions for binding and carrier sites. Although acidic injury is  
111 negligible in a medium at a pH above 4, lower pH in acid soils is one of the factors  
112 responsible for growth retardation, impairs mineral nutrients in plants<sup>16</sup>. In our investigation  
113 we detected an interesting relation between soil pH vs varieties and also these variables with  
114 Treatments (Figure 1). The initial pH rested in between 5 to 5.5 irrespective of treatments.  
115 Here Mahsuri considered being efficient variety which showed recovery of pH after sharp  
116 drop in the initial period. Similar improvement of pH was not observed in varieties Ranjit and  
117 Siyal Sali. The varieties Ranjit and Siyal Sali could not recover the initial pH (pH=5.2) up  
118 to 70 days after transplanting (DAT), rather a decreasing trend was detected for Ranjit at 300  
119 ppm Fe<sup>2+</sup> in the medium (Figure 1D). Of course Mahsuri showed a differential behaviour in  
120 the change of pH and showed better recovery at different growth stages. At maximum  
121 tillering stage (MTS) and panicle initiation stage (PIS) we observed a sharp increase in soil  
122 pH (pH =6) and a quick revival after 70 DAT (as straight line) at 300 ppm Fe<sup>2+</sup> (Figure 1D).  
123 A sound recovery in the pH in the efficient plants might lower the reduction of Fe<sup>3+</sup> to Fe<sup>2+</sup>  
124 on root surface through the release of other reductants and the plants sustain better  
125 physiological, biochemical activities. At nearly neutral pH solubility of Fe in the rooting  
126 medium is reduced by the fast oxidation of ferrous-Fe, favours the formation of iron plaque  
127 and hence iron immobilization occurs in the roots. Although we conducted the experiment  
128 with similar soil environment, different pH curves were documented by the three varieties  
129 which signify that pH variation in water saturated soil is also a varietal function.

130 A marked reduction in chlorophyll contents were observed at 300 ppm and 200 ppm Fe<sup>2+</sup>  
131 in the growth medium (Figure 2). Maximum chlorophyll content was recorded in the plants  
132 grown in control soil iron. Total leaf chlorophyll content was found to be reduced in the  
133 varieties Ranjit and Siyal Sali grown at higher level of iron (Figure 2). Interactions between  
134 levels of Fe<sup>2+</sup> and varieties were also found to be significant on the total chlorophyll content.  
135 Mahsuri recorded relatively higher chlorophyll content irrespective of the treatments  
136 compared to Ranjit and Siyal Sali. Mahsuri might have been able to maintain higher  
137 chlorophyll content through chloroplast development<sup>4</sup>. Ranjit recorded lower chlorophyll at  
138 200ppm and 300ppm Fe followed by Siyal Sali. Our findings also revealed the variation of  
139 total chlorophyll in different growth stages. Mahsuri sustained stable leaf chlorophyll content  
140 at different growth stages, quite reverse to the other two varieties (Figure 3). Perhaps, severe  
141 damage in cell structural components in early growing stages due to Fe<sup>2+</sup> mediated ROS,  
142 might be the reason of rapid reduction of chlorophyll content in these varieties.

143 The detrimental effect of Fe<sup>2+</sup> became more pronounced when its concentrations increase  
144 in waterlogged soil. In waterlogged soil, excess uptake of Fe<sup>2+</sup> by the roots and its acropetal  
145 translocation into the leaves must have catalyzed the generation of active O<sub>2</sub> species or free  
146 radicals which could render the peroxidation of chloroplast membranes, damage cell  
147 structural components and impair the plants' physiological processes and subsequently lead  
148 to a decrease in chlorophyll content in the sensitive varieties.

149 As expected, plants grown in higher soil Fe<sup>2+</sup> had higher Fe concentrations than those  
150 grown under control Fe levels, both in roots and shoots (Figure 4). Of course higher Fe  
151 concentrations were found in roots than in shoots. At 200 & 300 ppm Fe<sup>2+</sup>, higher  
152 concentrations of total Fe were detected in the plants of Ranjit and Siyal Sali, more  
153 susceptible to Fe toxicity, both in roots and shoot; with shoot concentrations nearly 2 times  
154 higher than in Mahsuri plants.

155 Expression of some plant ferritin isoforms can be induced by Fe overload<sup>17</sup> and iron  
156 storage inside ferritin could be related to Fe overload tolerance in some rice cultivars<sup>6</sup>.  
157 Surprisingly, lower shoot Fe in the present investigation could not define the tolerance  
158 capacity of the cultivars to ferritin expression. Audebert and Sahrawat (2000)<sup>18</sup> reported that  
159 Fe tolerant cultivar absorbed less Fe or translocated less Fe from root to shoot, a mechanism  
160 involved in cultivar differences in Fe toxicity tolerance. Here we suggest that Mahsuri plants  
161 are more resistant to excess Fe due to the possible induction of avoidance and / or exclusion  
162 mechanisms, allowing the plant to keep lower Fe amounts in its tissues and reducing Fe  
163 translocation to shoots. Moreover a large concentration of root Fe compared to shoot Fe  
164 concentrations might also be attributed to the formation of root plaque in the form of  
165 *Compound B* (goethite and lepidocrocite) as stated by Silvaira et al. (2007)<sup>6</sup>.

166 The root Zn concentrations for the three cultivars were found higher up to 1.5 times than  
167 shoot (Figure 5 A and D). Marked treatment effects were predicted in root and shoot Zn  
168 concentrations. Here Zn concentration decreases both in roots and shoots of the tree cultivars  
169 with the increment of Fe<sup>2+</sup> treatments. An apparent difference observed in Zn concentration  
170 under Fe excess was a higher Zn concentration in shoots of Mahsuri plants than in Ranjit and  
171 Siyal Sali plants. Shoot Zn concentration in Mahsuri was about 6 times higher than Ranjit and  
172 also above 2 times compared to Siyal Sali when the plants exposed to 300 ppm Fe<sup>2+</sup>.

173 Iron (III)-oxides are known to have a strong zinc-binding tendency. In waterlogged soil  
174 environment, Zn becomes available in the process of iron oxide reduction<sup>19</sup>. At the same time  
175 the plaque formation resulting from Fe re-oxidation around the rice root can reduce the  
176 concentration of soluble Zn in the rhizosphere by forming sparingly soluble ZnFe<sub>2</sub>O<sub>4</sub>  
177 complex<sup>20</sup>. Moreover reduced Fe can also exert a direct antagonistic effect on Zn uptake<sup>20</sup>.  
178 Sometimes it might also happened that the Fe plaque can lead to higher or lower Zn  
179 concentrations in shoots, depending on the size of plaque layer<sup>21</sup>. In the present work, the  
180 lower shoot Zn content compared to root in excess Fe<sup>2+</sup> may be referred to the root Fe plaque  
181 formation that seem to be acting either as a Zn reservoir or preventing Zn uptake. On the  
182 other hand a better shoot Zn content in Mahsuri indicates its tolerant capacity to higher Fe<sup>2+</sup>  
183 levels than the other two cultivars which may attributes to up regulation of some *ZIP* genes in  
184 Mahsuri plants<sup>22, 23</sup>.

185 Mn concentration in shoots were higher than roots in all the three cultivars, but a  
186 considerable reductions were observed in both roots and shoots subjected to higher Fe levels,  
187 with Ranjit reaching lowest levels of Mn concentration in roots and shoots (Figure 5 B and  
188 E).Of course, shoot Mn content in Mahsuri was significantly higher than other two cultivars  
189 at 300 ppm Fe<sup>2+</sup>. Precipitation of Mn in the Fe plaque may have resulted in its lower  
190 absorption by the cultivars where highest Fe concentrations were found. Such negative  
191 interactions between Fe and Mn have also been reported in plants<sup>24</sup>.

192 Except Mahsuri, lower Cu concentrations were recorded in roots and shoots of Ranjit  
193 and Siyal Sali when submitted to Fe excess. It has been suggested that the Fe plaque could act  
194 as a Cu reservoir in plants, increasing Cu absorption<sup>25</sup>. But in our experiment, reduction of  
195 Cu content in roots and shoots of Ranjit and Siyal Sali (Figure 5 C and F) might be due to  
196 formation of Fe plaque, being able to act as a barrier to Cu absorption<sup>6</sup> or preferential uptake  
197 of Fe<sup>2+</sup> on Fe overload, supported by highest shoot Fe concentrations. Mahsuri, on the other  
198 hand only the variety that could sustain better Cu concentrations by active absorption through

199 roots and dynamic translocation to shoots even 300 ppm Fe supplementation. The varietal  
200 differences in shoots Cu and Zn concentrations may also be attributed to higher Cu/Zn SOD  
201 activities in tolerant plants in Fe overload<sup>5</sup>.

202 The nitrogen concentrations for the plants grown in higher Fe were high than grown  
203 under control Fe levels, both in roots and shoots (Figure 6 C and F). Of course rate of  
204 increment in shoots were higher than roots. The percentage increase of shoots nitrogen  
205 concentration in Mahsuri plants was higher than the other two varieties. The variations  
206 pattern of nitrogen concentrations were similar to that of root and shoot Fe concentrations in  
207 the cultivars. Since in water saturated acidic soil  $\text{Fe}^{3+}$  and  $\text{NO}_3^-$  act as electron acceptors, a  
208 strong ionic competition of  $\text{Fe}^{2+}$  and  $\text{NO}_2^-$  might developed around roots' periphery and  
209 accelerated the uptake of nitrogen adduct along with Fe. Earlier studies have also  
210 demonstrated a direct relation in the uptake of Fe and N in wheat plant or seed and external  
211 supply of N at different phenological stages<sup>26,27,28</sup>.

212 Moreover, the uptake and transport of metals in plant is regulated by some special N  
213 loaded transporter proteins situated in different tissues of root, stem, leaf and reproductive  
214 parts. Many of them, like the proteins of *NRAMPs* and *ZIP* family are specific in transporting  
215 iron<sup>29,30</sup>. Thus the rice plants grown in higher soil  $\text{Fe}^{2+}$ , the superior uptake rate of Fe from  
216 soil and their translocation to leaf and to grain is facilitated by transporter proteins, which  
217 might be considered as the possible mechanism of higher N supplement to plants.

218 Phosphorus concentration decreased considerably in roots and shoots of Ranjit and Siyal  
219 Sali plants submitted to excess Fe but not in roots of Mahsuri plants (Figure 6 A and D). A  
220 decreasing trend of phosphorus concentration also observed in the shoots of Mahsuri plants  
221 when exposed 200 & 300 ppm  $\text{Fe}^{2+}$ , was suggesting the limited P translocation to the shoots  
222 of all the cultivars.

223 The concentration of phosphorus in the soil solution depends mainly on soil pH, and a  
224 decrease in pH can reduce P concentration by causing precipitation of amorphous Fe-  
225 phosphate polynuclear complexes with high surface area. In the present investigation we  
226 proposed that at low soil pH, higher amounts of Fe(III) oxides may be accumulated in the  
227 roots that can absorb anions such as phosphate and control the uptake of apoplast P into the  
228 simplast<sup>31,6</sup>. Our shoot data seems to agree with Howeler (1973)<sup>32</sup>, who states that, the root's  
229 apoplastic precipitation results in lower P absorption by the plant.

230 Potassium is a common macronutrient in plants that activates many enzymes involved in  
231 photosynthesis, respiration and plays important roles such as starch and protein synthesis, cell  
232 expansion, and stress alleviation<sup>33,34</sup>. However, higher Fe concentration in the medium plays  
233 an antagonistic role in plants' K uptake. Mehraban et al, (2008)<sup>34</sup> reported lower root and  
234 shoot K concentration under high Fe nutrition. In the present experiment K concentration  
235 decreased in the roots and shoots of Ranijt and Siyal Sali under excess  $\text{Fe}^{2+}$  in comparison to  
236 the control treatment (Figure 6 B and E) which may be considered as the consequence higher  
237 Fe nutrition and formation of Fe—K complex in soil solution. In contrast the plants of  
238 Mahsuri sustained stable K concentrations in roots and shoots, where it would be expected  
239 due to its higher sustainability to excess Fe toxicity.

#### 240 CONCLUSION:

241 The variability observed in the results of soil pH, leaf chlorophyll contents and all the  
242 major nutrients in roots and shoots under excess  $\text{Fe}^{2+}$  indicate the differential tolerance  
243 capacity among the cultivars. Although, root and shoot Fe and N concentrations showed  
244 positive correlation among the cultivars, the remarkable shoot Fe concentrations with  
245 simultaneous reduction in leaf chlorophyll contents explains the oxidative damage in the  
246 plants of Ranjit and Siyal Sali due to  $\text{Fe}^{2+}$  induced reactive oxygen species,  $\text{OH}^-$  radicals  
247 through Fentons' reactions. On the other hand, variety Mahsuri probably with its tolerable  
248 shoots Fe concentration and radical pH recovery, sustained ionic balance around root surface


249 and thereby showed a positive respond to Fe overload. This variety recorded superior  
250 nutrients status even at 300ppm and may be conspired as Fe tolerant cultivars. With deferred  
251 pH recovery, low leaf chlorophyll and reduced root and shoot nutrients level, the plants of  
252 Ranjit and Siyal Sali exhibited Fe susceptible nature when grown in Fe<sup>2+</sup> excess medium.  
253 Moreover, except Fe and N, all other nutrients seemed to have impaired uptake due to Fe  
254 toxicity in these susceptible cultivars compared to Mahsuri. Thus the plants of these cultivars  
255 appear to be affected by direct Fe toxicity as well as by pseudo Fe toxicity”-- Fe toxicity  
256 symptoms induced by nutrients deficiency. The Mahsuri cultivar seems to keep up mostly on  
257 avoidance and/or exclusion of Fe uptake into the plant and decreased translocation to shoots,  
258 being able to maintain higher nutrients levels in roots and shoots.

#### 259 **References:**


- 260 1. Curie C, Briat J F (2003) Iron transport and signaling in plants. *Annu. Rev. Plant Biol.*  
261 54:183-206
- 262 2. Bienfait H F, Van den Briel W, Mesland-Mul NT (1985) Free space iron pools in roots -  
263 generation and mobilization. *Plant Physiol.* 78:596-600
- 264 3. Sahrawat KL (2004) Iron toxicity in wetland rice and the role of other nutrients. *J. Plant*  
265 *Nutr.* 27:1471-1504
- 266 4. Baruah, K.K., Das, S., Das, K. (2007). Physiological disorder of rice associated with high  
267 levels of iron in growth medium. *J. Plant Nutr.* 30, 1871-1883
- 268 5. Saikia T. and Baruah K.K (2012) Iron toxicity tolerance in rice (*Oryza sativa*) and its  
269 association with anti-oxidative enzyme activity. *J.of Crop Sci.* 3, (3), 90-94.
- 270 6. Silveira, V. C.; Oliveira, A. P.; Sperotto, R. A.; Espindola, L. S.; Amaral, L.; Dias, J. F.;  
271 Cunha, J. B.; Fett, J. P. (2007) Influence of iron on mineral status of two rice (*Oryza sativa* L)  
272 cultivars. *Braz. J. Plant Physiol.* 19,127-139
- 273 7. Backer, M. Asch, F.( 2005), Iron toxicity in rice- conditions and management concepts. *J.*  
274 *Plant Nutr. Soil Sci.* 168, 558—573
- 275 8. Benckiser, G., Ottow, J. C. G., Santiago, S., Watanabe, I. (1983): Iron toxicity – Effect of  
276 phosphorus, potassium, calcium and magnesium fertilization on rhizoflora, redox potential  
277 and iron uptake by different rice varieties (*Oryza sativa* L.). *Landw. Forschung* 36, 285–299.
- 278 9. Sahrawat, K.L. (2010): Reducing iron toxicity in lowland rice with tolerant genotypes and  
279 plant nutrition Plant Stress. Global Science Books
- 280 10. Wu L. B., Shhadi M. Y., Gregorio G., Matthus E., Becker M. and Frei M. (2014), Genetic  
281 and physiological analysis of tolerance to acute iron toxicity in rice. *Rice*, 7:8
- 282 11. Fisher R. A., and Yates F. (1957), *Statistical Tables for Biological, Agricultural and*  
283 *Medical Research.* Fifth Edition, Published by Oliver and Boyd, London,
- 284 12. Anderson, J.M. and Boardman, N.K (1964). Studies on the greening of dark grown bean  
285 plants. II.Development of photochemical activity. *Aust. J.Biol. Sci.*, 17: 93-101.
- 286 13. Jackson, M.L. (1973) *Soil Chemical Analysis.* Prentice Hall of India Pvt. Ltd., New Delhi.
- 287 14. A.O.A.C. (1970) *Official Method of Analysis.* Association of Official Analytical  
288 Chemists. 11<sup>th</sup> Ed. Washington D.C. USA.
- 289 15. Cochran,W.G.;Cox, G.M. *Experimental Design.* John Wiley and Sons Inc., New York.  
290 1963.
- 291 16. Arnon D.I., Fratzke W.E., Johnson C.M. (1942) Hydrogen ion concentration in relation to  
292 absorption of inorganic nutrient by higher plants. *Plant Physiol* 17:515.
- 293 17. Petit J-M, Briat JF, Lobreaux S (2001) Structure and differential expression of the four  
294 members of the Arabidopsis thaliana ferritin gene family. *Biochem. J.* 359:575-582.
- 295 18. Audebert A, Sahrawat KL (2000) Mechanisms for iron toxicity tolerance in lowland rice.  
296 *J Plant Nutr* 23(11–12):1877–1885

- 297 19. Mandal B. Chatterjee J Harza G C and Mandal L N. (1992) Effect of preflooding on  
298 transformation of applied zinc and its uptake by rice in lateritic soils. *Soil Science* 153: 250-  
299 257
- 300 20. Sajwan, K. S., Lindsay, W. L. (1988) Effects of redox on zinc deficiency in paddy rice.  
301 *Soil Sci. Soc. Am. J.* 50, 1264–1269.
- 302 21. Zhang, X., Zhang, F., Mao, D. (1998) Effect of iron plaque outside roots on nutrient  
303 uptake by rice (*Oryza sativa* L.): II. Zinc uptake by Fe-deficient rice. *Plant Soil* 202, 33–39
- 304 22. Lee S, An G (2009a) Over-expression of OsIRT1 leads to increased iron and zinc  
305 accumulations in rice. *Plant Cell Environ* 32:408–416
- 306 23. Krithika S. and Balachandar D. (2016) Expression of Zinc Transporter Genes in Rice as  
307 Influenced by Zinc-Solubilizing *Enterobacter cloacae* Strain ZSB14. *Frontiers in Plant*  
308 *Science.* 7 (446)
- 309 24. Fageria N. K. and Moreira A., (2011). The Role of Mineral Nutrition on Root Growth of  
310 Crop Plants. In Donald L. Sparks, editor: *Advances in Agronomy*, Vol. 110, Burlington:  
311 Academic Press, pp. 251-331.
- 312 25. Ye Z H, Cheung K C, Wong M H (2001) Copper uptake in *Typha latifolia* as affected by  
313 iron and manganese plaque on the root surface. *Can. J. Bot.* 79:314-320
- 314 26. Kutman U. B., Yildiz B., Ozturk L., Cakmak I. (2010). Biofortification of durum wheat  
315 with zinc through soil and foliar applications of nitrogen. *Cereal Chem.* 87, 1–  
316 910.1094/CCHEM-87-1-0001
- 317 27. Shi R., Zhang Y., Chen X., Sun Q., Zhang F., Römheld V., Zou C. (2010). Influence of  
318 long-term nitrogen fertilization on micronutrient density in grain of winter wheat (*Triticum*  
319 *aestivum* L.). *J. Cereal Sci.* 51, 165–170.10.1016/j.jcs.2009.11.008
- 320 28. Cakmak I, Pfeiffer W. and McClafferty B. (2010) Bio fortification of durum wheat with  
321 zinc and iron. *Cereal Chem.* 87:10-20
- 322 29. Sperotto R A, Vasconcelos M W, Grusak M A and Fett J P. (2012) Effects of different  
323 Fe supplies on mineral partitioning and remobilization during the reproductive development  
324 of rice (*Oryza sativa* L.). *Rice*, 5:27
- 325 30. Masuda H, Aung M S and Nishizawa N K,(2013) Iron bio fortification of rice using  
326 different transgenic approaches. *Rice*, 6:40
- 327 31. Zhang X, Zhang F, Mao D (1999) Effect of iron plaque outside roots on nutrient uptake  
328 by rice (*Oryza sativa* L. ): Phosphorus uptake. *Plant Soil* 209:187-192
- 329 32. Howeler R H (1973) Iron-induced orange disease of rice in relation to physico-chemical  
330 changes in a flooded oxisol. *Soil Sci. Soc. Am. Proc.* 37:898-903
- 331 33. Marschner H., (1995). Mineral nutrition of higher plants. Academic Press, London, pp:  
332 313—323.
- 333 34. Mehraban P, Zadeh A A, Sadeghipour H R (2008) Iron toxicity in rice (*Oryza sativa* L.)  
334 under different potassium nutrition. *Asian J Plant Sci* 7:251–259.
- 335 **Figure and Table Captions:**
- 336 Figure 1: Varietal impacts of iron treatments on pH change in the soil solutions at different  
337 growth stages.
- 338 Figure 2: Varietal impacts of iron treatments on leaf chlorophyll contents at different growth  
339 stages (in mg g<sup>-1</sup> FW). The vertical bars represent the standard errors.
- 340 Figure 3: Varietal change on leaf chlorophyll contents at different growth stages (in mg g<sup>-1</sup>  
341 FW). The vertical bars represent the standard errors.
- 342 Figure 4: Varietal impacts of iron treatments on roots and shoots Fe. The vertical bars  
343 represent the standard errors.
- 344
- 345 Figure 5: Varietal impacts of iron treatments on roots and shoots Zn (A, D), Mn (B, E) and  
346 Cu (C, F) (in µg g<sup>-1</sup> DW). The vertical bars represent the standard errors.

347 Figure 6: Varietal impacts of Fe treatments on root and shoot P (A, D), K (B, E), and N (C, F)  
 348 (in  $\mu\text{g g}^{-1}$  DW). The vertical bars represent the standard errors.  
 349  
 350  
 351  
 352




353  
 354 Figure 1: Varietal impacts of iron treatments on pH change in the soil solutions at different  
 355 growth stages.  
 356




357




358 Figure 2: Varietal impacts of iron treatments on leaf chlorophyll contents at different growth  
 359 stages (in  $\text{mg g}^{-1}$  FW). The vertical bars represent the standard errors.  
 360



361 Figure 3: Varietal change on leaf chlorophyll contents at different growth stages (in  $\text{mg g}^{-1}$   
 362 FW). The vertical bars represent the standard errors.  
 363  
 364  
 365




366 Figure 4: Varietal impacts of iron treatments on roots and shoots Fe. The vertical bars  
 367 represent the standard errors.  
 368  
 369



370  
371  
372  
373

Figure 5: Varietal impacts of iron treatments on roots and shoots Zn (A, D), Mn (B, E) and Cu (C, F) (in  $\mu\text{g g}^{-1}$  DW). The vertical bars represent the standard errors.



374  
 375 Figure 6: Varietal impacts of iron treatments on roots and shoots P (A, D), K (B, E), and N  
 376 (C, F) (in  $\mu\text{g g}^{-1} \text{DW}$ ). The vertical bars represent the standard errors.