

Effect of cotton seed meal on the performance traits and meat composition in commercial broilers

Shilpi Islam^{1*}, Mohammad Nazrul Islam², A. K. M. Ahsan Kabir³

¹Department of Animal Science and Nutrition, Bangabandhu Sheikh Mujibur Rahman Agricultural University, Salna, Gazipur-1706, Bangladesh.

²Director, Phenix Group, House No. 39A, Road No. 8, Dhanmondi, Dhaka, Bangladesh.

³Department of Animal Science, Bangladesh Agricultural University, Mymensingh- 2202, Bangladesh.

ABSTRACT

Aims: To evaluate the effect of different levels of cotton seed meal (CSM) on performance traits and meat composition in commercial broilers.

Place and Duration of the study: The experiment was carried out at Bangabandhu Sheikh Mujibur Rahman Agricultural University, Salna, Gazipur during the period from 26 February, 2018 to 10 April, 2018.

Study design and methodology: Two hundred fifty two Cobb-500 day-old broiler chicks with good health were randomly allotted to six dietary treatments in three replications with fourteen birds per replication in a complete randomized design for 35 days period. The dietary treatments were: T0, soyabean meal (SBM) based diet; T1, 10% CSM protein with 90% SBM protein; T2, 20% CSM protein with 80% SBM protein; T3, 30% CSM protein with 70% SBM protein; T4, 40% CSM protein with 60% SBM protein and T5, 50% CSM protein with 50% SBM protein. The mash feed was supplied *ad libitum* basis.

Results: Average feed intake (g/d) was increased ($P=0.001$) in higher amount of CSM group. Dressing percentage was tended to higher ($P = 0.089$) in T0 and lower value was showed in T5. Crude protein (CP) content of breast meat significantly ($P < 0.01$) affected among the treatments. The highest CP content was observed in T5 (22.57%) and lowest CP content was in T1 (21.12%). Crude fiber (CF) content was significantly increased ($P < 0.01$) in the diet contained higher amount of CSM (0.35%, 0.32%, 0.31%, 0.22%, 0.13% for T5, T4, T3, T2 and T1; respectively) and the lowest CF was observed for T0 (0.11%). Ether extract (EE) of breast muscle was also significantly increased ($P < 0.01$) in the diet contained higher amount of CSM (1.27%, 1.15%, 1.12%, 1.09%, 1.05% for T5, T4, T3, T2 and T1; respectively) and lower EE was observed in T0 group (1.01%). Ash content was higher ($P < 0.05$) in T0 (1.49%), T2 (1.48%) and T3 (1.45%) group compare to others. The second higher value was observed for T1 (1.4%) diet and the lowest ash content was observed in T4 (1.25%) and T5 (1.32%).

Conclusion: It would be concluded that CSM can be a substitute of soyabean meal in broiler ration and up to 40% CSM protein can be incorporated in broiler chicken diet without any adverse effects.

Keywords: Cotton seed meal; soyabean meal; broiler; carcass weight; dressing percentage; breast muscle.

21 1. INTRODUCTION

22

23 Broilers play an important role in human nutrition, national income, employment and income
24 generation in Bangladesh. As an important **sub-sector** of livestock production, the poultry
25 industry in Bangladesh plays a vital role in economic growth and simultaneously creates
26 numerous employment opportunities. Poultry industry is a fundamental part of animal
27 production, is committed to the nation for supplying a cheap source of good quality nutritious
28 animal protein in terms of meat and eggs [1]. It was recorded that poultry meat alone
29 contributes 37% of the total meat production in Bangladesh [2]. Poultry contributes about 22-
30 27% of the total animal protein supply in the country. So, to cope with market demand for
31 animal meat protein, modern broilers are reaching market age sooner each year. Therefore,
32 advances in nutrition will be the fundamental for securing this rapid growth achievement and
33 maintaining sustainable broiler production. Soyabean meal is generally **recognized** as an
34 effective and high-quality vegetable protein feed-stuff [3, 4]. Recently in Bangladesh, high
35 demand of soyabean meal has been observed but its availability is not sufficient round the
36 year and the prices are also higher in off-season. Therefore, it is very important to improve
37 the scientific knowledge for utilizing low cost locally available agro-industrial by-products in
38 broiler feed in order to reduce the feed cost and to substitute as an effective protein source.
39 **Cottonseed meal** is one of them. **Cotton seed meal** is a by-product of cotton seed that is
40 used for animal feeding because it is rich in oil and protein [5]. CSM is a fairly good source of
41 protein (222.0 to 560.2 g per kg); [6, 7] and metabolizable energy (7.4 to 11.99MJ per kg);
42 [7]. Another researcher reported that cottonseed cake has been used as a cheaper
43 alternative to soybean cake in livestock feeding and a good source of dietary protein [8]. So,
44 CSM is very useful in livestock feeding in the cotton growing areas. Although CSM is an
45 inexpensive source of protein with high protein content [9], its nutrient bioavailability in
46 poultry diets is low due to the presence of anti-nutritional factors, such as free gossypol,
47 Cycloproponic fatty acids and crude fibre [10], which may cause negative effects on growth,
48 reproductive performance and organ abnormalities [9, 11]. But cottonseed products offer a
49 safe alternative feed when fed at recommended levels [12, 13, 14]. If carefully incorporated,
50 cotton seed meal can reduce feed costs while maintaining or increasing the level of bird's
51 performance. **In Bangladesh context, there is very little research work on inclusion of CSM
52 as an alternative protein source in broiler diets.** Therefore, the purpose of this study was to
53 evaluate the effect of different levels of cotton seed meal on performance traits and meat
54 composition in commercial broilers.

55

56 2. MATERIAL AND METHODS

57

58 2.1 Animal, experimental design and management

59 The experiment was carried out at Bangabandhu Sheikh Mujibur Rahman Agricultural
60 University Poultry Farm, Salna, Gazipur, Bangladesh. Two hundred fifty two (252) healthy
61 day-old Cobb-500 broiler chicks were weighed and randomly allocated to six dietary
62 treatments replicated three times with fourteen birds per replicate in a Complete
63 Randomized Design (CRD). **The dietary treatments were: T0, SBM based diet; T1, 10% CSM
64 protein with 90% SBM protein; T2, 20% CSM protein with 80% SBM protein; T3, 30% CSM
65 protein with 70% SBM protein; T4, 40% CSM protein with 60% SBM protein and T5, 50%
66 CSM protein with 50% SBM protein.** A strict bio-security program was maintained inside and
67 outside of the research shed. The birds were vaccinated against Infectious Bursal Disease
68 (IBD) and Newcastle Disease (ND). The management practices were identical for all dietary
69 groups. Electric light was provided for 24 hours and the brooding temperature was almost
70 maintained at 33±2 °C for first week. In course of the trial, the temperature was gradually
71 reduced to 25±2 °C at the end of the experiment. Fresh and dried saw dust was used at a
72 depth of about 3 cm for bedding material. The birds were critically observed twice a day for
73 clinical sign if any (slow movement, infrequent sitting, lack of appetite, significant changes of

74 feathering, paralysis etc.) and for monitoring other activities. **Feeders were** cleaned in each
75 week and **drinkers were** washed twice daily.

76 **2.2 Preparation of experimental diet and feeding**

77 The experimental diets were formulated by replacing soyabean meal with CSM according to
78 the recommendation of NRC [15] in three phases namely starter (1 to 14 days), grower (15
79 to 28 days) and finisher (29 to 35 days). All feed ingredients were weighed separately and
80 soyabean oil was incorporated into soyabean meal first and then mixed thoroughly with other
81 macro ingredients. Micro ingredients were mixed thoroughly with the ground maize and then
82 mixed with the other macro ingredients. Diet for each treatment was prepared properly as
83 per recommendation. The ingredients and nutritional composition of different diets (starter,
84 grower and finisher) are presented in Table 1, Table 2 and Table 3; respectively. All diets
85 were free **from** antibiotics. The broiler mash feed was supplied three times daily on an ad
86 libitum basis.
87

88 **2.3 Slaughtering and sample collection of broilers**

89 **At the 35th day** of the experiment, three (3) birds from each replicate were randomly
90 selected from each pen and each broiler chicken was weighed. Birds were sacrificed and
91 hanged until complete bleeding. After complete bleeding the birds feathers were removed by
92 hand and pinning was done manually. Viscera and giblet were removed from the carcass.
93 Legs, head, neck and shank were separated from the body parts. Live bird, slaughtered bird
94 (after complete bleeding), skin, viscera, giblet, legs, head, neck, shank and carcass were
95 weighed individually. Breast muscles were collected randomly from each replicate.
96

97 **2.4 Parameters measured**

98 The feed intake of each replication was determined by subtracting the amount of left over
99 from the amount of supplied feed on the previous day. Live weight of each bird was recorded
100 as the average weight of all birds of each replicate. Carcass weight and dressing percent
101 were calculated accordingly by considering the live weight of broilers for each replication.
102

103 **2.5 Chemical analysis**

104 Samples of breast meat were analyzed to determine dry matter (DM), crude protein (CP),
105 ether extract (EE), crude fibre (CF), nitrogen free extract (NFE) and total ash were
106 determined according to the methods of Association of Official Analytical Chemists [16].
107

108 **2.6 Statistical Analysis**

109 The data were analyzed by using the statistical program (SPSS 16.0) to compute analysis of
110 variance (ANOVA) for a completely randomized design (CRD) and Duncan's multiple range
111 test (DMRT) was done to differentiate among the treatment means at 5% level of significant.
112
113
114
115
116
117
118
119
120
121
122
123
124
125

126
127

Table 1: Ingredients composition and nutrient content of broiler starter diet

Items	Treatments					
	T0	T1	T2	T3	T4	T5
Ingredients (% as fed basis)						
Corn	54.73	51.38	47.88	44.28	40.53	36.35
Cotton seed meal	0	5.13	10.5	16.04	21.75	28.19
Soyabean meal	29	26.7	24.29	21.8	19.25	16.34
Soyabean oil	1.25	1.77	2.31	2.86	3.45	4.1
Distillers Dried Grains with Solubles (DDGs)	6	6	6	6	6	6
Protein concentrate	6	6	6	6	6	6
Lime stone	1.4	1.4	1.4	1.4	1.4	1.4
Di calcium phosphate	0.6	0.6	0.6	0.6	0.6	0.6
^a Vitamin–Mineral Premix	0.25	0.25	0.25	0.25	0.25	0.25
Threonine	0.05	0.05	0.05	0.05	0.05	0.05
L- Lysine	0.1	0.1	0.1	0.1	0.1	0.1
DL-Methionine	0.25	0.25	0.25	0.25	0.25	0.25
Salt	0.3	0.3	0.3	0.3	0.3	0.3
Enzyme	0.04	0.04	0.04	0.04	0.04	0.04
Phytase	0.01	0.01	0.01	0.01	0.01	0.01
Anti-Oxidant	0.02	0.02	0.02	0.02	0.02	0.02
Total	100	100	100	100	100	100
Calculated nutrient analysis						
ME (Kcal/Kg)	2951.08	2951.17	2951.04	2951.14	2951.17	2951.31
Crude Protein (%)	23.02	23.02	23.02	23.02	23.03	23.02
Linoleic acid (%)	1.15	1.08	1.00	0.93	0.84	0.75
Ca (%)	1.12	1.13	1.14	1.15	1.15	1.16
P (Total) (%)	0.68	0.69	0.70	0.70	0.71	0.72
P (non-phy) (%)	0.47	0.47	0.47	0.46	0.46	0.46
Na (%)	0.16	0.16	0.15	0.15	0.15	0.14
Cl (%)	0.22	0.21	0.21	0.21	0.21	0.20
K (%)	1.76	1.69	1.62	1.54	1.46	1.36
Lysine (%)	1.24	1.22	1.21	1.20	1.18	1.17
Methionine (%)	0.64	0.64	0.65	0.65	0.65	0.65
Cystine (%)	0.31	0.32	0.32	0.32	0.32	0.32
Methionine +cystine (%)	0.96	0.96	0.96	0.97	0.97	0.97
Threonine (%)	0.72	0.72	0.71	0.71	0.70	0.69
Tryptophan (%)	0.28	0.28	0.27	0.27	0.26	0.26
Feed cost/kg (Tk.)	37.61	37.45	37.27	37.09	36.92	36.71

128
129
130
131
132
133
134
135
136
137

^aVitamin–Mineral Premix provided the following per kilo gram of diet: Vitamin A, 5.0 MU; Vitamin D, 1.0 MU; Vitamin E, 10.0 g; Vitamin K, 1.6 g; Vitamin B1, 0.6 g; Vitamin B2, 2.0 g; Vitamin B6, 1.6 g; Vitamin B12, 4.0 mg; Biotin, 20.0 mg; Pantothenic Acid, 4.0 g; Folic Acid, 0.2 g; Nicotinic Acid, 12.0 g; Copper, 2.4 g; Iron, 9.6 g; Zinc, 160 g; Manganese, 19.2g; Selenium, 0.05 g; Cobalt, 0.12 g; Iodine, 0.24 g

138
139
140
141

Table 2: Ingredients composition and nutrient content of broiler grower diet

Items	Treatment					
	T0	T1	T2	T3	T4	T5
Ingredients (% as fed basis)						
Corn	54.48	51.14	47.64	44.05	40.29	36.11
Cotton seed meal	0	5.13	10.5	16.03	21.78	28.19
Soyabean meal	29.01	26.7	24.29	21.8	19.22	16.34
Soyabean oil	3.8	4.32	4.86	5.41	6	6.65
Distillers Dried Grains with Solubles (DDGs)	6	6	6	6	6	6
Protein concentrate	3.7	3.7	3.7	3.7	3.7	3.7
Lime stone	1.4	1.4	1.4	1.4	1.4	1.4
Di calcium phosphate	0.6	0.6	0.6	0.6	0.6	0.6
^a Vitamin –Mineral Premix	0.25	0.25	0.25	0.25	0.25	0.25
Threonine	0.05	0.05	0.05	0.05	0.05	0.05
L- Lysine	0.1	0.1	0.1	0.1	0.1	0.1
DL-Methionine	0.25	0.25	0.25	0.25	0.25	0.25
Salt	0.3	0.3	0.3	0.3	0.3	0.3
Enzyme	0.04	0.04	0.04	0.04	0.04	0.04
Phytase	0.01	0.01	0.01	0.01	0.01	0.01
Anti-Oxidant	0.015	0.015	0.015	0.015	0.015	0.015
Total	100	100	100	100	100	100
Calculated nutrient analysis						
ME (Kcal/Kg)	3101.42	3101.59	3101.46	3101.05	3101.56	3101.73
Crude Protein (%)	21.55	21.55	21.55	21.55	21.55	21.55
Linoleic acid (%)	1.15	1.07	1.00	0.92	0.84	0.75
Ca (%)	0.98	0.98	0.99	1.00	1.01	1.01
P (Total) (%)	0.61	0.61	0.62	0.63	0.64	0.64
P (non-phy) (%)	0.39	0.39	0.39	0.39	0.39	0.38
Na (%)	0.16	0.16	0.15	0.15	0.15	0.14
Cl (%)	0.22	0.21	0.21	0.21	0.21	0.20
K (%)	1.76	1.69	1.61	1.54	1.45	1.36
Lysine (%)	1.15	1.13	1.12	1.11	1.09	1.08
Methionine (%)	0.60	0.60	0.60	0.60	0.61	0.61
Cystine (%)	0.28	0.28	0.29	0.29	0.29	0.29
Met+cys (%)	0.88	0.88	0.89	0.89	0.90	0.90
Threonine (%)	0.72	0.72	0.71	0.71	0.70	0.69
Tryptophan (%)	0.27	0.27	0.26	0.26	0.25	0.25
Feed cost/kg (Tk.)	37.17	37.01	36.83	36.65	36.47	36.27

142 ^aVitamin–Mineral Premix provided the following per kilo gram of diet: Vitamin A, 5.0 MU; Vitamin D, 1.0
143 MU; Vitamin E, 10.0 g; Vitamin K, 1.6 g; Vitamin B1, 0.6 g; Vitamin B2, 2.0 g; Vitamin B6, 1.6 g;
144 Vitamin B12, 4.0 mg; Biotin, 20.0 mg; Pantothenic Acid, 4.0 g; Folic Acid, 0.2 g; Nicotinic Acid, 12.0 g;
145 Copper, 2.4 g; Iron, 9.6 g; Zinc, 160 g; Manganese, 19.2g; Selenium, 0.05 g; Cobalt, 0.12 g; Iodine,
146 0.24 g

147
148
149
150

151
152
153
154

Table 3: Ingredients composition and nutrient content of broiler finisher diet

Items	Treatment					
	T0	T1	T2	T3	T4	T5
Ingredients (% as fed basis)						
Corn	64.08	62.32	60.7	58.62	56.67	54.44
Cotton seed meal	0	2.7	5.2	8.4	11.4	14.8
Soyabean meal	16.01	14.8	13.67	12.23	10.88	9.36
Soyabean oil	2.5	2.77	3.02	3.34	3.64	3.99
Distillers Dried Grains with Solubles (DDGs)	5	5	5	5	5	5
Protein concentrate	9.5	9.5	9.5	9.5	9.5	9.5
Lime stone	1.3	1.3	1.3	1.3	1.3	1.3
Di calcium phosphate	0.6	0.6	0.6	0.6	0.6	0.6
^a Vitamin-Mineral Premix	0.25	0.25	0.25	0.25	0.25	0.25
Threonine	0.05	0.05	0.05	0.05	0.05	0.05
L- Lysine	0.1	0.1	0.1	0.1	0.1	0.1
DL-Methionine	0.25	0.25	0.25	0.25	0.25	0.25
Salt	0.3	0.3	0.3	0.3	0.3	0.3
Enzyme	0.04	0.04	0.04	0.04	0.04	0.04
Phytase	0.01	0.01	0.01	0.01	0.01	0.01
Anti-Oxidant	0.015	0.015	0.015	0.015	0.015	0.015
Total	100	100	100	100	100	100
Calculated nutrient analysis						
ME (Kcal/Kg)	3121.64	3121.48	3121.41	3121.27	3121.13	3121.45
C.Protein (%)	20.05	20.05	20.05	20.05	20.05	20.05
Linoleic acid (%)	1.28	1.24	1.20	1.16	1.12	1.07
Ca (%)	1.28	1.28	1.28	1.29	1.29	1.30
P (Total) (%)	0.73	0.73	0.74	0.74	0.74	0.75
P (non-phy) (%)	0.55	0.55	0.55	0.55	0.55	0.55
Na (%)	0.16	0.16	0.15	0.15	0.15	0.15
Cl (%)	0.21	0.21	0.21	0.21	0.21	0.21
K (%)	1.46	1.42	1.39	1.34	1.30	1.25
Lysine (%)	1.03	1.03	1.02	1.01	1.01	1.00
Methionine (%)	0.64	0.64	0.64	0.65	0.65	0.65
Cystine (%)	0.29	0.29	0.29	0.29	0.29	0.29
Met+cys (%)	0.93	0.93	0.93	0.93	0.94	0.94
Threonine (%)	0.53	0.53	0.53	0.52	0.52	0.52
Tryptophan (%)	0.21	0.21	0.21	0.20	0.20	0.20
Feed cost/kg (Tk.)	39.33	39.24	39.16	39.05	38.95	38.85

155 ^aVitamin–Mineral Premix provided the following per kilo gram of diet: Vitamin A, 5.0 MU; Vitamin D, 1.0
156 MU; Vitamin E, 10.0 g; Vitamin K, 1.6 g; Vitamin B1, 0.6 g; Vitamin B2, 2.0 g; Vitamin B6, 1.6 g;
157 Vitamin B12, 4.0 mg; Biotin, 20.0 mg; Pantothenic Acid, 4.0 g; Folic Acid, 0.2 g; Nicotinic Acid, 12.0 g;
158 Copper, 2.4 g; Iron, 9.6 g; Zinc, 160 g; Manganese, 19.2g; Selenium, 0.05 g; Cobalt, 0.12 g; Iodine,
159 0.24 g
160

161 3. RESULTS AND DISCUSSION

162 3.1 Performance traits

163 Performance traits of broilers fed different experimental diets are presented in Table 4.
164 Average feed intake was significantly higher (P < 0.01) in the diets containing higher amount
165

166 of CSM. This result is consistent with the observation of other researchers [17, 8] who
 167 reported that CSM influence higher feed intake and at moderate levels (20-30%) of CSM
 168 incorporation feed intake can be increased, which impairs feed efficiency [18]. In this study,
 169 there was no significant difference ($P > 0.05$) for average live weight gain when broilers fed
 170 different levels of CSM, which were also consistent with the findings of previous studies [10,
 171 19, 9]. Although, the birds fed on diet T2, T3 and T4 had their weights numerically tended to
 172 improved, but the birds with diet T2 showed superiority in weights over other diets. These
 173 results showed consonance with earlier researcher report [17], who concluded that feeding
 174 cotton seed cake up to 50% had no significant effect on performance of broiler chickens.
 175 However, decreased efficiency of CSM utilization was also observed when the level of CSM
 176 was increased in the diet [23, 24]. On the other hand, another researchers [25] finding were
 177 disagreed with the previous results on live weight and feed conversion ratio and reported
 178 that, there were no adverse effect of CSM at the level of 30%. Similar results were also
 179 observed in this study which was the fully agreed above statement. Live weight and carcass
 180 weight did not show any significant difference among the treatments. But dressing
 181 percentage was tended to significant ($P = 0.089$) among the treatments. The higher value
 182 was observed in control (0% CSM) group and the lower value was for T5 group where
 183 broilers received 50% CSM protein. However, after receiving of CSM diet (up to 15%)
 184 dressing percentage value were 64.8 to 66.8% [14], which was more or less similar to the
 185 present observations. No significant difference was observed in feed cost per kg live weight
 186 gain. However, some researchers [17, 26] reported that feed cost was numerically
 187 decreased with increasing levels of CSM in the diet. In this study similar trend was also
 188 observed because CSM is relatively cheaper compared to soyabean meal in the market. But
 189 higher percent of CSM level influence the higher amount of feed intake. Accordingly, the cost
 190 per kg live weight gain was similar to all diets. The substitution of soyabean meal with CSM
 191 might have lowered the actual energy content [27] and digestible lysine content [28, 29, 14]
 192 of the diets. Supplementation of lysine can help to alleviate the negative effects of
 193 cottonseed meal [20, 21, 22]. In this study, 100g L-lysine was added to all of the diets. Due
 194 to addition of same amount of lysine in all treatments, the beneficial effect of lysine
 195 supplementation on free gossypol was not prominent in this study. As a result the average
 196 growth rate was more or less similar in all of the treatments.

197
198
199 **Table 4. Performance traits of broilers fed different experimental diets**
200

Parameters	Dietary treatment					SEM	P-value	
	T0	T1	T2	T3	T4			T5
Average feed intake (g)	91.99 ^a	91.46 ^a	93.37 ^b	94.65 ^b	94.16 ^b	93.51 ^b	1.24	0.001
Average live weight gain (g/d)	48.59	48.61	50.45	48.99	48.85	48.17	0.79	0.616
Carcass traits								
Live weight (g)	1876.22	1950.67	1851.00	1896.00	1916.11	1830.89	15.39	0.737
Carcass weight (g)	1259.84	1304.22	1223.12	1222.76	1250.56	1179.47	13.19	0.525
Dressing percentage (%)	67.12 ^b	66.74 ^{ab}	66.03 ^{ab}	64.41 ^a	65.18 ^{ab}	64.43 ^a	0.49	0.089
Feed cost/kg live weight gain (Tk.)	72.29	71.46	69.91	72.75	72.21	72.51	0.450	0.698

201 ^{a,b} Means with different letters in rows differ significantly ($P \leq 0.05$)
202

203 3.2 Nutrient composition of meat

204 Nutrient compositions of breast meat of broilers of different treatments are shown in the
 205 Table 5. No significant difference was found for the DM content of broilers breast meat
 206 ranged due to the treatments. CP content of breast meat was significantly ($P < 0.01$) differed
 207 among the treatments. The highest CP content was observed in T5 and the lowest CP

208 content was in T1. Second lowest value was showed by T4. However, T0, T2 and T3 did not
 209 show significant difference among them. Little information is available about the effects of
 210 CSM on the meat compositions of broiler chickens. It was reported that the CP content of
 211 breast muscle was 22.57 to 23.08% for day 42 and day 52 Cobb broiler chickens [30] and
 212 was 19.7% for day 45 Cobb broiler chickens [31]. In this study, similar values were also
 213 found for the CP content of breast muscle of Cobb-500 broiler chickens at 35th day. Higher
 214 level of CSM influenced the higher fibre content in breast meat. The CF content of breast
 215 muscle was significantly ($P < 0.01$) higher in T5 diet and significantly lower value was
 216 observed in T0 and T1 diets. The CF content of breast muscle was increased with increasing
 217 the CSM in diets. Higher amount of CSM may influence the higher amount of CF in breast
 218 muscle. Cotton seed meal contained higher amount of EE compared to soyabean meal
 219 which may influenced ($P < 0.01$) the higher intramuscular EE content of breast muscle in
 220 higher CSM receiving groups (T5) compared to small amount of CSM contained diets
 221 receiving group (T1) and the lower EE value was observed for control group (T0). The
 222 increased EE in breast muscle were observed when broiler fed higher percentage of CSM
 223 containing diets, which might be attributed to the enhanced anabolism of intramuscular fat
 224 [9]. However, others observed that the EE content of breast muscle was 2.22 to 2.55% [30]
 225 and 3.6% [31] which value was higher with compare to this research. Ash content was
 226 higher ($P < 0.05$) in T0, T2 and T3 diets compare to the other treatment diets. But T0, T2
 227 and T3 diets did not show any significant difference among the diets. The second higher
 228 value was observed for T1 diet but T0, T1, T2 and T3 did not show any significant difference
 229 among the treatments. However, the lowest ash content was observed in T4 but T4 and T5
 230 did not differ significantly between the diets for the ash content of breast muscle. This
 231 observation was more or less similar (1.13% to 1.17% and 1.4%) with the result that was
 232 reported by others [30, 31] for meat composition of Cobb broilers. Mortality was only 0.5%
 233 and no health problems were detected, need for prolonged feeding trial to assess safety and
 234 productivity of the use of CSM is clear warranted.

235
236
237

Table 5: Nutrient composition of breast meat for different experimental diets

Parameters	Dietary treatment						SEM	P-value
	T0	T1	T2	T3	T4	T5		
DM%	24.82	23.78	25.09	25.25	24.41	25.59	1.00	0.688
<i>Nutrient composition (% DM basis)</i>								
CP%	22.11 ^c	21.12 ^a	22.22 ^c	22.18 ^c	21.61 ^b	22.57 ^d	0.84	0.000
CF%	0.11 ^a	0.13 ^a	0.22 ^b	0.31 ^c	0.32 ^{cd}	0.35 ^d	0.10	0.000
EE%	1.01 ^a	1.05 ^{ab}	1.09 ^{abc}	1.12 ^{bc}	1.15 ^c	1.27 ^d	0.10	0.001
Ash%	1.49 ^c	1.4 ^{bc}	1.48 ^c	1.45 ^c	1.25 ^a	1.32 ^{ab}	0.10	0.002

238 ^{a,b,c}Means with different letters in rows differ significantly ($P \leq 0.01$)

239

240 4. CONCLUSION

241

242 From the results of this study it would be concluded that, CSM protein can be incorporated
 243 (up to 40%) in broiler chicken diet without any adverse effects on feed quality and birds
 244 performance.

245

246 ACKNOWLEDGEMENTS

247

248 The authors gratefully acknowledges to the Research Management Committee (RMC) of
 249 Sheikh Mujibur Rahman Agricultural University (BSMRAU), Salna, Gazipur for providing the
 250 financial support to conduct this research work.

251

252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303

COMPETING INTERESTS

“Authors have declared that no competing interests exist”.

AUTHORS' CONTRIBUTIONS

Author SI designed the study, wrote the protocol and performed the experiment. Author AKM AK performed the statistical analysis. Author SI and MNI managed the literature searches and wrote the first draft of the manuscript. Author MNI will give the financial support for publication charges. All authors read and approved the final manuscript.

REFERENCES

1. Akter A, Uddin S. Bangladesh Poultry Industry. *Journal of Business and Technology (Dhaka)*. 2009; 4(2): 97-112.
2. DLS. Bangladesh delta plan 2100 formulation project (Livestock). General economics division planning commission Government of Bangladesh. 2015.
3. Azarm HM, Lee SM. Effect of partial substitution of dietary fish meal by fermented soyabean meal on growth performance, amino acid and biochemical parameters of juvenile black sea bream *acanthopagrus schlegeli*. *Aquaculture Research*. 2014; 45: 994-1003.
4. Kim S W. Identification of a second major antigenic epitope in the α -subunit of soy α -conglycinin. *Food and Agricultural Immunology*. 2014; 25: 311-321.
5. Gadelha ICN, Fonseca NBS, Oloris SCS, Melo MM, Soto-Blanco B. Gossypol toxicity from cottonseed products. *The Scientific World Journal*. 2014, 231635.
6. Anonymous. Cottonseed meal. *Pakistan Poultry*. 1996; 7.
7. Nagalakshmi, D, Rao SVR, Panda AK, Sastry VRB. Cottonseed meal in poultry diets: a review. *Poultry Science*. 2007; 44: 119-134.
8. Adeymo GO, Longe OG. Effects of graded levels of cottonseed cake on performance, haematological and carcass characteristics of broiler fed from day old to 8 weeks of age. *African Journal of Biotechnology*. 2007; 6(8): 1064-1071.
9. Nie C, Zhang W, Wenxia GE, Wang Y, Liu Y, Liu J. Effect of fermented cotton seed meal on growth performance, apparent digestibility, carcass traits and meat composition in yellow-feathered broilers. *Turkish Journal of Veterinary and Animal Science*. 2015; 39: 350-356.
10. Tang JW, Sun H, Yao XH, Wu YF, Wang X, Feng J. Effects of replacement of soybean meal by fermented cottonseed meal on growth performance, serum biochemical parameters and immune function of yellow-feathered broilers. *Asian-Australasian Journal of Animal Sciences*. 2012; 25(3): 393-400.
11. S. ŚWIĄTKIEWICZ. The use of cotton seed meal as a protein source for poultry: An update review. *Worlds Poultry Science Journal*. 2016; 72: 473-484.
12. Sterling KG, Costa EF, Henry MH, Pesti GM, Bakalli RI. Responses of broiler chickens to cottonseed and soybean meal-based diets at several protein levels. *Poultry Science*. 2002; 81(2): 217-26.
13. Perez-Maldonado RA. Canola meal and Cottonseed meal in broiler and layer diets. A report for the Australian Egg Corporation Limited. 2003.
14. Mishra A, Ray S, Sarkar SK, Haldar S. Cotton seed meal as a partial replacement for soyabean meal in cob 400 broiler ration. *Indian Journal of Animal Nutrition*. 2015; 32(1): 69-74.

- 304 15. NRC (National Research Council). Nutrient requirements of domestic animals No. 1
305 Nutrient Requirement of Poultry. 9th rev. ed. National Academy of Science, Washington
306 D.C. 1994.
- 307 16. AOAC. Official methods of analysis. 16th Edn. Association of official Analytical
308 Chemistry, Washington, DC. 1995.
- 309 17. Ojewola GS, Ukachukwu SN, Okulonye EI. Cottonseed meal as substitute for soyabean
310 meal in broiler ration. *International Journal of Poultry Science*. 2006; 5 (4): 360-364.
- 311 18. Batonon-Alavo DI, Faruk MU, Lescoat P, Weber GM, Bastianelli D. Inclusion of
312 sorghum, millet and cottonseed meal in broiler diets: a meta-analysis of effects on
313 performance. *Animal*. 2015; 9(7): 1120-1130.
- 314 19. Sun H, Tang JW, Fang CL, Yao XH, Wu YF, Wang X, Feng J. Molecular analysis of
315 intestinal bacterial microbiota of broiler chickens fed diets containing fermented cotton
316 seed meal. *Poultry Science*. 2013; 92: 392-401.
- 317 20. Henry MH, Pesti GM, Bakalli R, Lee J, Toledo RT, Eitenmiller RR, Phillips RD. The
318 performance of broiler chicks fed diets containing extruded cottonseed meal
319 supplemented with lysine. *Poultry Science*. 2001; 80(6): 762-768.
- 320 21. Sekhar Reddy P, Sudhakar Reddy P, Satyanarayana Reddy PVV, Srinivasa Rao D.
321 Influence of cottonseed cake on the performance of broilers. *Indian Journal of Animal
322 Nutrition*. 1998; 15:188-193.
- 323 22. Ryan JR, Kratzer FH, Grau CR, Vohra P. Glandless cottonseed meal for laying and
324 breeding hens and broiler chicks. *Poultry Science*. 1986; 65(5): 949-955.
- 325 23. Fafiolu AO, Oso AO, Bangbose, AM, Omodia RE, Sediq RM. Performance of weaner
326 rabbits fed cottonseed cake as replacer for soyabean meal. In: Proc. 31st Annual
327 Conference of the Nigerian Society for Animal Production. 2006 (12-15th March); 376-
328 379.
- 329 24. Mahmood F, Khan MZ, Khan A, Muhammad G, Javed I. Lysine induced modulation of
330 toxicopathological effects of cottonseed meal in broiler breeder males. *Pakistan Journal
331 of Zoology*. 2011; 43(2): 357-365.
- 332 25. Abdulrashid, M, Joseph ZO, Mohammed A, Adamu HY. Response of broiler chickens
333 fed cottonseed meal based diets. *International Journal of Agricultural Research*. 2013, 1:
334 62-65.
- 335 26. Attanayaka PMGSK, Pathirana, APDG, Priyankarage, N, Silva SSP, Nayananjalie WAD.
336 Effect of substitution of soyabean meal with cottonseed meal on the performances of
337 broiler chicken. *International Journal of Livestock. Research*. 2016; 6(3): 24-30.
- 338 27. Nzekwe NM, Olomu JM. Cottonseed as a substitute for groundnut meal in the rations of
339 laying chicken and growing turkeys. *Journal of Anim. Prodction Research*. 1984; 4: 57-71.
- 340 28. Heidarinia A, Malakian M. Nutritional evaluation of cottonseed meal with and without
341 ferrous sulfate for broiler chickens. *Research Journal of Poult Science*. 2011; 4: 14-17.
- 342 29. Zaboli GR, Miri A. Effect of dietary lysine to crude protein ratio in diets containing corn,
343 cottonseed meal and soyabean meal on broiler performance during starter period. *Life
344 Science Journal*. 2013; 10: 454-458.
- 345 30. Suchý P, Jelíek P, Straková E, Hucl J. Chemical composition of muscles of hybrid
346 broiler chickens during prolonged feeding. *Czech Journal of Animal Science*. 2002;
347 47(12): 511-518.
- 348 31. Oliveira Jde, Avanco SV, Garcia-Neto M, Ponsano EHG. Composition of broilers meat.
349 *Journal of Applied Poultry Research*. 2016; 25: 173-181.
- 350