

# Comparison of Jacobi and Gauss-Seidel Iterative Methods for the Solution of Systems of Linear Equations

## ABSTRACT

In this research work two iterative methods of solving system of linear equation has been compared, the iterative methods are used for solving sparse and dense system of linear equation and the methods were being considered are: Jacobi method and gauss-seidel method. The results show that gauss-seidel method is more efficient than Jacobi method by considering maximum number of iteration required to converge and accuracy.

*Keywords: Iterative methods; Linear equations problem; Convergence; square matrix.*

## 1. INTRODUCTION

The development of numerical methods on a daily basis is to find the right solution techniques for solving problems in the field of applied science and pure science, such as weather forecasts, population, the spread of the disease, chemical reactions, physics, optics and others.

Many problems (Kendall E 2007) in applied mathematics involve solving systems of linear equations, with the linear system occurring naturally in some cases and as a part of the solution process in other cases

Collections of linear equations is called linear systems of equations. They involve same set of variables. Various methods have been introduced to solve systems of linear equations (N. A. Saeed and A.Bhatti 2008). There is no single method that is best for all situations. These methods should be determined according to speed and accuracy. Speed is an important factor in solving large systems of equations because the volume of computations involved is huge.(Noreen Jamil 2015)

Systems of linear equations (Kalambi 2008) arise in a large number of areas both directly in modeling physical situations and indirectly in the numerical solutions of the other mathematical models. These application occur in virtually all areas of the physical, biological and social science. A linear equation in the variable  $x_1, x_2, \dots, x_n$  is any equation of the form  $a_1x_1 + a_2x_2 + \dots + a_nx_n = b_1$  (1)

It is fast and simple to use when coefficient matrix is sparse. There are many approaches of solving system of linear equations i.e. direct methods and Indirect (iterative) methods. The direct methods give the exact solution in which there is no error except the round off error due to the machine, whereas iterative methods give the approximate solutions in which there is some error (Iascar A. H. and Samira Behera, 2014).

## 2. JACOBI METHOD

The first iterative technique is called the Jacobi method, after Carl Gustav Jacob Jacobi (1804–1851) a German mathematicians who was one of the famous algorist formulate an iterative method of solving system of linear equations. This method makes two assumptions:

- (1) That the system given by has a unique solution and
- (2) That the coefficient matrix A has no zeros on its main diagonal.

If any of the diagonal entries are zero, then rows or columns must be interchanged to obtain a coefficient matrix that has nonzero entries on the main diagonal.

## 3. GAUSS-SIEDEL METHOD

The next method is called Gauss-Seidel method, which the modification of Jacobi method, named after Carl Friedrich Gauss (1777–1855) and Philipp L. Seidel (1821–1896). This modification is no more difficult to use than the Jacobi method, and it often requires fewer iterations to produce the same degree of accuracy. With the Jacobi method, the values of obtained in the nth approximation remain unchanged until the entire nth approximation has been calculated. With the Gauss- Seidel method, on the other hand, we use the new values of each as soon as they are known. That is, once we have determined from the first equation, its value is

then used in the second equation to obtain the new values. Similarly, the new value and the first value are used in the third equation to obtain the new and so on

#### 4. ANALYSIS METHOD 1

The Jacobi method was obtain by solving the  $i$ th equation in  $Ax = b$ , to obtain  $x_i$

(Provided  $a_{ii} \neq 0$ ) i. e. given a system of linear equation

$$\begin{aligned}
 a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \dots + a_{1n}x_n &= b_1 \\
 a_{21}x_1 + a_{22}x_2 + a_{23}x_3 + \dots + a_{2n}x_n &= b_2 \\
 a_{31}x_1 + a_{32}x_2 + a_{33}x_3 + \dots + a_{3n}x_n &= b_3 \\
 \cdot & \cdot \cdot \cdot \cdot \cdot \cdot \\
 \cdot & \cdot \cdot \cdot \cdot \cdot \cdot \\
 \cdot & \cdot \cdot \cdot \cdot \cdot \cdot \\
 a_{n1}x_1 + a_{n2}x_2 + a_{n3}x_3 + \dots + a_{nn}x_n &= b_n
 \end{aligned} \tag{2}$$

To begin, solve first equation for  $x_1$ , second equation for  $x_2$ , third equation for  $x_3$  and so on to obtain

$$\begin{aligned}
 x_1^{k+1} &= \frac{1}{a_{11}}(b_1 - a_{12}x_2^k - a_{13}x_3^k - \dots - a_{1n}x_n^k) \\
 x_2^{k+1} &= \frac{1}{a_{22}}(b_2 - a_{21}x_1^k - a_{23}x_3^k - \dots - a_{2n}x_n^k) \\
 x_3^{k+1} &= \frac{1}{a_{33}}(b_3 - a_{31}x_1^k - a_{32}x_2^k - \dots - a_{3n}x_n^k) \\
 & \cdot \\
 & \cdot \\
 & \cdot \\
 x_n^{k+1} &= \frac{1}{a_{nn}}(b_n - a_{n1}x_1^k - a_{n2}x_2^k - \dots - a_{nn-1}x_{n-1}^k)
 \end{aligned} \tag{3}$$

Then make initial guess (zero iteration) for the solution  $x^{(0)} = (x_1^{(0)}, x_2^{(0)}, x_3^{(0)} \dots x_n^{(0)})$  substitute these value into the right hand side of (3). This constitute first iteration

$x^1 = (x_1^{(1)}, x_2^{(1)}, x_3^{(1)} \dots x_n^{(1)})$ . Second iteration is obtained by substituting first iteration into

the left hand side of (3.2), that is  $x^2 = (x_1^{(2)}, x_2^{(2)}, x_3^{(2)}, \dots x_n^{(2)})$ . And so on. The Jacobi method

can be generalize as for each  $k \geq 0$  we can generate the component  $x_i^{k+1}$  of  $x^{k+1}$  from  $x^k$  by

$$x_i^{k+1} = \frac{1}{a_{ii}} \left[ \sum_{j=1, j \neq i}^n (-a_{ij}x_j^k) + b_i \right] \quad \text{For } i = 1, 2, \dots, n$$

The Jacobi method in matrix form can be found by considering an  $n \times n$  system of linear equation

$Ax = b$  where

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ a_{31} & a_{32} & a_{33} & \dots & a_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & a_{n3} & \dots & a_{nn} \end{pmatrix}, \quad x = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_n \end{pmatrix} \quad \text{and } b = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \\ \vdots \\ b_n \end{pmatrix}. \quad \text{We split matrix}$$

$$A = \begin{pmatrix} a_{11} & 0 & 0 & \dots & 0 \\ 0 & a_{22} & 0 & \dots & 0 \\ 0 & 0 & a_{33} & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & a_{nn} \end{pmatrix} - \begin{pmatrix} 0 & 0 & \dots & 0 & 0 \\ -a_{21} & 0 & \dots & 0 & 0 \\ -a_{31} & -a_{32} & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ -a_{n1} & -a_{n2} & \dots & -a_{n,n-1} & 0 \end{pmatrix} -$$

$$\begin{pmatrix} -a_{11} & -a_{12} & -a_{13} & \dots & -a_{1n} \\ 0 & -a_{22} & -a_{23} & \dots & -a_{2n} \\ 0 & 0 & 0 & \dots & \vdots \\ \vdots & \vdots & \vdots & \ddots & -a_{n-1,n} \\ 0 & 0 & 0 & \dots & 0 \end{pmatrix} = D - L - U.$$

Therefore the matrix  $Ax = b$  can be transform into  $(D - L - U) x = b$ , this implies that

$$Dx = (L + U)x + b.$$

## 5. ANALYSIS OF METHOD 2

With Jacobi method, the value of  $x^{k+1}$  was obtain in (k+1)th iteration remain unchanged until the entire (k+1)th iteration has been calculated. With gauss-seidel method we use the value of  $x_i^{k+1}$  as soon as they are known. That is

$$\begin{aligned} x_1^{k+1} &= \frac{1}{a_{11}} (b_1 - a_{12}x_2^k - a_{13}x_3^k - \dots - a_{1n}x_n^k) \\ x_2^{k+1} &= \frac{1}{a_{22}} (b_2 - a_{21}x_1^{k+1} - a_{23}x_3^k - \dots - a_{2n}x_n^k) \\ x_3^{k+1} &= \frac{1}{a_{33}} (b_3 - a_{31}x_1^{k+1} - a_{32}x_2^{k+1} - \dots - a_{3n}x_n^k) \\ &\vdots \\ &\vdots \\ x_n^{k+1} &= \frac{1}{a_{nn}} (b_n - a_{n1}x_1^{k+1} - a_{n2}x_2^{k+1} - \dots - a_{nn-1}x_{n-1}^{k+1}) \end{aligned} \tag{4}$$

Then make initial guess (zero iteration) for the solution  $x^{(0)} = (x_1^{(0)}, x_2^{(0)}, x_3^{(0)} \dots x_n^{(0)})$  substitute these value into the right hand side of (3.3). This constitute first iteration,  $x^1 = (x_1^{(1)}, x_2^{(1)}, x_3^{(1)} \dots x_n^{(1)})$ . Second iteration is obtained by substituting first iteration into the left hand side of (3.3),  $x^2 = (x_1^{(2)}, x_2^{(2)}, x_3^{(2)}, \dots x_n^{(2)})$ . And so on. This method can be generalize as for each  $k \geq 0$  we can generate the component  $x_i^{k+1}$  of  $x^{k+1}$  from  $x^k$  by

$$x_i^{k+1} = \frac{1}{a_{ii}} [-\sum_{j=1}^{i-1} (a_{ij}x_j^k) - \sum_{j=i+1}^n (a_{ij}x_j^{k+1}) + b_i] \quad \text{For } i = 1, 2 \dots n$$

The gauss-seidel method in a matrix form is given by;

$$(D - L)x^{k+1} = Ux^k + b$$

$$x^{k+1} = (D - L)^{-1}Ux^k + (D - L)^{-1}b$$

## 6. CONVERGENCE OF ITERATIVE METHOD

The rate of convergence of iterative methods determine how fast the error  $|x^k - x|$  goes to zero as  $k$ , the number of iteration increases. The sufficient condition of convergence for iterative methods define as  $x^{k+1} = Bx^k + c$  to converge is that  $\rho(B) = \max_{1 \leq i < n} |\lambda_i(B)| < 1$  where  $\rho(B)$  is the spectral radius of  $B$ . This condition is fulfilled for both Jacobi and Gauss-seidel methods if the coefficient matrix is diagonally dominant for any choice of initial approximation.

**THEOREM (6.1):** For any  $x^0$  in  $R^n$  the sequence  $\{x^{k+1}\}_{k=0}^{\infty}$  define by  $x^{k+1} = Tx^k + c$ , for each  $k \geq 0$  converge to a unique solution of  $x = Tx + c$  iff  $\rho(T) < 1$

**6.2 Absolute error** = |True value – approximate value|

**6.3 Relative error:** =  $\frac{\text{Absolute error}}{|\text{True value}|}$

**6.4 percentage relative error** = Relative error  $\times$  100%

## 7. NUMERICAL EXPERIMENTS

The problem in system of linear equations shall be considered

### Problem 7.1

Solve the equation using Jacobi's method.

$$4x_1 - x_2 - x_4 = 0$$

$$-x_1 + 4x_2 - x_3 - x_5 = 5$$

$$-x_2 + 4x_3 - x_6 = 0$$

$$-x_1 + x_4 - x_5 = 6$$

$$-x_2 - x_4 + 4x_5 - x_6 = -2$$

$$-x_3 - x_5 + 4x_6 = 6$$

Taking the initial approximation  $x_1^k = x_2^k = x_3^k = x_4^k = x_5^k = x_6^k = 0$  starting with these value and continuous to iterate we obtain the solution in the table below

**TABLE 1: Iteration Result for Jacobi Method**

Iteration	X <sub>1</sub>	X <sub>2</sub>	X <sub>3</sub>	X <sub>4</sub>	X <sub>5</sub>	X <sub>6</sub>
0	0.000000	0.000000	0.000000	0.000000	0.000000	0.000000
1	0.000000	1.250000	0.000000	1.500000	-0.500000	1.500000
2	0.687500	1.125000	0.687500	1.375000	0.562500	1.375000

3	0.625000	1.73445	0.625000	1.812575	0.46875	1.812575
4	0.886756	1.679688	0.886756	1.773438	0.839900	1.773438
5	0.823282	1.898554	0.823282	1.926865	0.806641	1.926865
6	0.956355	1.883301	0.956355	1.917481	0.938668	1.917481
7	0.950196	1.962695	0.950196	1.973606	0.929566	1.973606
8	0.984075	1.957490	0.984075	1.969941	0.977477	1.969941
9	0.981858	1.986497	0.981858	1.990388	0.974343	1.990388
10	0.994199	1.984515	0.994199	1.989050	0.991796	1.989050
11	0.993391	1.995041	0.993391	1.996499	0.990684	1.996499
12	0.997887	1.994354	0.997887	1.996011	0.997011	1.996011
13	0.997593	1.99816	0.997593	1.998725	0.996595	1.998728
14	0.999230	1.997945	0.999230	1.998547	0.998912	1.998547
15	0.999123	1.999343	0.999123	1.999536	0.998760	1.999536
16	0.999720	1.999252	0.999720	1.999471	0.999604	1.999471
17	0.999681	1.999761	0.999681	1.999831	0.999549	1.999831
18	0.999898	1.999728	0.999898	1.999808	0.999856	1.999808

This complete the table of the solution to the system of linear equation given above and the values of  $x_i$  are  $(x_1, x_2, x_3, x_4, x_5, x_6) = (0.999898, 1.999728, 0.999898, 1.999808, 0.999856, 1.999808)$  respectively.



**Problem 7.2**

Solve the equation using gauss-seidel method.

$$4x_1 - x_2 - x_4 = 0$$

$$-x_1 + 4x_2 - x_3 - x_5 = 5$$

$$-x_2 + 4x_3 - x_6 = 0$$

$$-x_1 + x_4 - x_5 = 6$$

$$-x_2 - x_4 + 4x_5 - x_6 = -2$$

$$-x_3 - x_5 + 4x_6 = 6$$

Taking the initial approximation  $x_1^k = x_2^k = x_3^k = x_4^k = x_5^k = x_6^k = 0$

**TABLE 2: : Iteration Result for Gauss-Seidel Method**

Iteration	X <sub>1</sub>	X <sub>2</sub>	X <sub>3</sub>	X <sub>4</sub>	X <sub>5</sub>	X <sub>6</sub>
0	0.000000	0.000000	0.000000	0.000000	0.000000	0.000000
1	0.000000	1.250000	0.312500	1.500000	0.187500	1.625000
2	0.687500	1.546875	0.79296	1.718750	0.722656	1.878906
3	0.816406	1.833008	0.927979	1.884766	0.899188	1.956792
4	0.929444	1.939153	0.973986	1.957158	0.963276	1.984316
5	0.974078	1.977835	0.990538	1.984339	0.986623	1.994290

6	0.990544	1.991926	0.996554	1.994292	0.995127	1.997920
7	0.996555	1.997059	0.998745	1.997921	0.998225	1.999243
8	0.998745	1.998929	0.999543	1.999243	0.999354	1.999724
9	0.999543	1.999610	0.999834	1.999724	0.999765	1.999900
10	0.999834	1.999858	0.999940	1.999900	0.999915	1.999964

Hence the solution is obtain after ten successive iteration we have  $x_1 = 0.999834$ ,  $x_2 = 1.999858$ ,  $x_3 = 0.999940$ ,  $x_4 = 1.999900$ ,  $x_5 = 0.999915$ ,  $x_6 = 1.999964$

### 7.2 Error analysis of Jacobi method

The true value are  $(x_1, x_2, x_3, x_4, x_5, x_6) = (1, 2, 1, 2, 1, 2)$  while the computed value are  $(0.999898, 1.999728, 0.999898, 1.999808, 0.999856, 1.999808)$ . Hence we determine the error as follows by using  $x_2$

Absolute error = |True value – approximate value| =  $|2 - 1.999728| = 0.000272$

Percentage relative error =  $\frac{\text{Absolute error}}{|\text{True value}|} = 0.000136 * 100\% = 0.0136\%$

### 7.3 Error analysis of gauss-seidel method

The true value are  $(x_1, x_2, x_3, x_4, x_5, x_6) = (1, 2, 1, 2, 1, 2)$  while the computed value are,  $(0.999834, 1.999858, 0.999940, 1.999900, 0.999915, 1.999964)$ . Hence we determine the error as follows by using  $x_2$

Absolute error = |True value – approximate value| = |2 - 1.999834| = 0.000166

Percentage relative error =  $\frac{\text{Absolute error}}{|\text{True value}|} = 0.000166 * 100\% = 0.0083\%$

#### 7.4 COMPARISON OF TWO ITERATIVE METHODS USED IN THE EVALUATION

**TABLE 3 showing percentage error of both methods**

METHODS	NO OF ITERATION	ERROR%
JACOBI METHOD	18	0.0136
GAUSS-SIEDAL METHOD	10	0.0083

#### 8 Conclusion

There are different methods of solving system of linear equation, some are direct methods while some are numerical method (iterative method). In this research work two iterative methods of solving system of linear equations has been presented where the gauss-seidel method proved to be the best and effective in the sense that it converge very fast. From the practical example, we observe that the required solution were obtained with very little iteration without much problem relating to the starting condition.

#### REFERENCE

- Agmon. S. The relaxation method for linear inequalities. Canadian Journal Mathematics, pages 382–392, 1954.
- Anita H.M. (2002). Numerical-Methods for Scientist and Engineers. Birkhauser-verlag.

- Censor and T.Y. Elfving. New methods for linear inequalities, linear algebra and its applications. *Linear Algebra and Its Applications*, 64:243–253, 1985.00
- Censor. Y. Row-action methods for huge and sparse systems and their applications. pages 444–466, 1981.
- De Pierro A. and A. Iuseam (1985). A simultaneous projection method for linear inequalities. *Linear Algebra and Its Applications*, 64:243–253.
- Kalambi, I. B. (December, 2008).” A Comparison of three Iterative Methods for the Solution of Linear Equations” *J. Appl. Sci. Environ. Manage.* December, 2008 Vol. 12 (4), 53 – 55
- Kalambi, I.B. (1998). ‘Solutions of Simultaneous Equations by Iterative Methods’. Postgraduate Diploma in Computer Science Project. Abubakar Tafawa Balewa University, Bauchi.
- Kendall E Atkinson (June 2007) *Numerical method, quantitative analysis*
- Motzkin and Schoenberg. The relaxation method for linear inequalities. *Canadian Journal Mathematics*, pages 393–404, 1954.
- Noreen Jamil (2012) “A Comparison of Direct and Indirect Solvers for Linear Systems of Equations” *Int. J. Emerg. Sci.*
- PARLETT B. N. (1980). A new look at the lanczos algorithm for solving symmetric systems of linear equations. *Lin. Alg. Appl*, pages 323–346.
- Rajasekaran. S. *Numerical methods in Science and Engineering*. Wheelerand Co. Ltd Allahabad, 1992. 11
- SAAD Y. and M. H. SCHULTZ. A generalized minimal residual algorithm for solving non-symmetric linear systems. 7, 1986.

SAAD.Y. Practical use of some krylov subspace methods for solving indefinite and unsymmetrical linear systems. pages 203–228, 1984.

Saeed, N. A. A.Bhatti. Numerical Analysis. Shahryar, 2008.

Turner. P. Guide to Numerical Analysis. Macmillan Education Ltd. Hong Kong, 1989.