

## Original Research Article

# Effect of Nitrogen Rates on Growth, Carbon Assimilation and Quality of Water Spinach (*Ipomea aquatica*)

### ABSTRACT

**Aims:** This study was conducted to investigate the impact of the nitrogen fertilization on the growth, leaf gas exchange and bio-metabolite accumulation in *Ipomea aquatica*.

**Treatment and Study experimental design:** *Ipomea aquatica* plants were exposed to four different rates of nitrogen (0, 30, 60 and 90 N kg/ha) using NPK green fertilizer as a nitrogen source using. The experiment was laid out in Complete Randomize Design (CRD).

**Place and Duration of Study:** Department of Biology, Faculty of Science, Universiti Putra Malaysia between September to November 2016.

**Methodology:** There were four nitrogen rates were applied (0, 30, 60 and 90 N Kg/ha) using NPK green fertilizer as a nitrogen source. The growth data collections were conducted once a week after the application of the treatments for the plant growth parameter. The total chlorophyll content in the leaves was measured using a SPAD chlorophyll meter. The leaf gas exchange was determined using a LI-6400XT portable photosynthesis system. Total phenolics and flavonoid was determined using Folin-Ciocalteu reagent.

**Results:** It was found that the growth parameters which are plant height, leaf numbers, branches numbers, total biomass and chlorophyll content recorded the highest measurement at 90 kg N/ha and the lowest at 0 kg N/ha. As for the leaf gas exchange, the positive effect of nitrogen fertilization on kangkung was shown by the increased in photosynthesis rate (A) and stomatal conductance (gs) where the highest measurement recorded at 90 kg N/ha, while the lowest at 0 kg N/ha. However, the water use efficiency (WUE) decreased as the nitrogen rates increased. At lower rates of nitrogen fertilization (30 kg N/ha) produced the highest production of secondary metabolites, where the total phenolics and flavonoids production were enhanced compared to other nitrogen treatments.

**Conclusion:** In conclusion, as the nitrogen rates increased, the growth and leaf gas exchange properties was enhanced however the production of total phenolics and flavonoids was reduced and get the highest accumulation at 30 kg N/ha.

**Keywords:** [Nitrogen, *Ipomea aquatica*, growth, leaf gas exchange, biometabolites production]

## 1. INTRODUCTION

In Malaysia, agriculture sector has contributed about 8.5% to Gross Domestic Products (GDP). About 39% of the contributions originated from the production of food crops, fruits and vegetables. It is estimated that there are about 44, 000 hectares of the total area in Malaysia was used for vegetable cultivation [1]. According to Department of Agriculture Malaysia in 2011, *Ipomea Aquatica* is one out of ten types of vegetables that occupied consumed the largest highest area for vegetable production. This plant is among the most consumed vegetable in Asia. This is because of its low price compared to other types of vegetable. Kangkung air or its scientific name, *Ipomea aquatica*, is a widely known leafy vegetable especially in Asian country. The plant is also commonly known with different local names, such as water spinach, swamp cabbage, or water convolvulus. From its scientific classification, kangkung has been classified under the a family of Convolvulaceae [2]. According to [3], Convolvulaceae family consists of primarily 1650 of tropical species. Moreover, the genus of kangkung which is *Ipomea* has about 500 to 600 different species and it has been the most number of containing species in Convolvulaceae family [4]. This species of family can nicely be grown at almost anywhere at the higher or even the lower land. *Ipomea aquatica* is one of the species that is cultivated on the higher land. Besides easy to be grown, kangkung often be the favorable plant to be cultivated because it does not take long time to mature and harvest. It can easily adapt towards its

**Comment [A1]:** There is no coherency in the whole abstract. Author(s) need to re-write the whole abstract to be more scientific and to conform with the journal format

**Comment [A2]:** Write in full since it is appearing for the first time

31 grow environment and usually unsusceptible to disease. Almost all parts of kangkung plant are edible  
32 [3].  
33

34 According to Susila et al. [5], nitrogen is the primary nutrient that involve in producing a high yield of  
35 vegetables. Nitrogen is one of the macro-nutrients that is very crucial especially for a plant to have a  
36 proper growth and development [6] such as that required in constructing the matter of the plant cell  
37 and tissue [7]. The amount of nitrogen in the soil could be insufficient for the plant to grow. Therefore,  
38 the source of nitrogen for plant especially in agriculture field often be found in the form of a fertilizer.  
39 Both organic and inorganic nitrogen fertilizer are widely used in agriculture especially in cultivating  
40 green crops to keep the source of nutrients for plant being supply [6]. Practically, an appropriate and  
41 suitable amount of nitrogen to be given to plant will affect it's crop yield. Nitrogen also is very  
42 important especially to promote the growth of the plant leaf [8]. Nitrogen is a crucial element not only  
43 to promote the growth and plant development, also increase yield and quality in vegetable crops.  
44 Increasing level of nitrogen resulting in a number of leaves, leaf length and plant body [8]. Nitrogen  
45 also enhanced the size of fruits and vegetables where at optimum application of N will resulting in a  
46 better size. The metabolic process which stimulated by N by enhance the vegetative and also the  
47 reproductive growth in plant. Besides, high plant biomass can be obtained when there is high N  
48 accumulated in shoot, along with the increasing of root growth in plant if there is sufficient amount of  
49 N supply [9]. However, the lack of N in plant would caused the reduced in plant development and  
50 eventually will lower the crop yield.  
51

52 Nitrogen had been proven to have a strong relationship with photosynthesis process in plant.  
53 Increasing N level lead to higher N content in leaf. N also enhances the leaf chlorophyll and CO<sub>2</sub>  
54 assimilation which increase in the Rubisco activity [10]. Therefore, increase in rate of photosynthesis  
55 **photosynthesis** is the most vital biochemical process in plants [11]. According to [12,13], **rate of the**  
56 **photosynthesis rate** (A) depends on the growth development of the plant's leaf. The leaf development  
57 includes the increase in leaf area, leaf thickness, the surface volume of mesophyll cells, and leaf  
58 chloroplast. The photosynthesis rate will be increased as the leaf development also increased [14].  
59 Nitrogen is an element that has a significance role in photosynthesis which involve in the opening of  
60 the stomata. The stomatal vent will decrease following the nitrogen deficit which then will decrease  
61 the transpiration rate [15].  
62

63 Secondary metabolites such as phenolic in plant **is** usually associated with the plant survival and  
64 health benefits for those who consume the plant. Low nitrogen level in plant has been reported to  
65 have more secondary metabolites compare to plant that has high N level [16]. Application of more N  
66 level resulting in decrease of phenolic concentrations based on carbon/nutrients balance (CNB)  
67 hypothesis [17]. Flavanoids also a secondary metabolite which is widely distributed with different  
68 functions in plants. The biological functions of flavonoids include defense against UV-B radiation,  
69 pathogen infection, nodulation and pollen fertility [18]. A study done by [8] on leaf mustard where the  
70 total phenolics concentration was observed to be decreased as the level increased. It is well known  
71 that nitrogen application can directly affect the morphological growth and yield of this plant, however,  
72 little work has been carried out to look on the impact of nitrogen of the leaf gas exchange properties  
73 and previous work have not comprehensively considered the production of secondary metabolites of  
74 *I. aquatica* under nitrogen fertilization. there was no study about the production of secondary  
75 metabolites and leaf gas exchange properties under different nitrogen rates. The main aim of the  
76 research was to investigate the effect of nitrogen fertilization on the growth, leaf gas exchange and  
77 production of secondary metabolites of *I. aquatica* and to determine the best nitrogen rates for growth  
78 and development of *I. aquatica*. This research will provide the important information for vegetable  
79 growers that involved in cultivation of vegetables in Malaysia.  
80

## 81 2. MATERIAL AND METHODS

### 82 83 2.1. Plant material and maintenance

84  
85 The experiment was conducted at **the** Department of Biology, Universiti Putra Malaysia, Serdang  
86 (UPM), Selangor **where the area of the experiment was located at the site where the direct**  
87 **sunlight is always sufficient and available**. These seeds were **pre-germinated in the nursery**  
88 **allowed to germinate** for two weeks. **After which they all the seeds has been germinated, all the**  
89 **plants then** were transplanted into the polybags filled with a mixture of topsoil, organic matter and  
90 sand with the ratio of 3:2:1. The nitrogen sources used were from the NPK green fertilizer (15:15:15).

91 The polybags were arranged according to Completely Randomized Design (CRD) with five  
92 replications. There were four nitrogen rates were applied (0, 30, 60 and 90 Kg N/ha) with overall 160  
93 of *I. aquatica* plants were used. The growth data collections were conducted once a week for four  
94 weeks after the application of the treatments for the plant growth parameter. Whereas the destructive  
95 analysis and leaf gas exchange of the experiment was conducted at the end of the experiment.

## 96 **2.2. Plant height, leaf and branch numbers**

97  
98  
99 As for plant height, it was measured starting from the stem that was at the soil surface up until the  
100 highest shoot grow or at tip using measuring tape. The leaf and branches number were counted  
101 manually per plant basis

## 102 **2.3 Plant total dry weight measurement**

103  
104  
105 The plants were first removed from the soil carefully and the dirt from the soil were washed with tap  
106 water. After that, the shoot and the root parts were separated. All the plants were dried in an oven for  
107 48 hours at temperature of 60°C until constant weight reached.

## 108 **2.4 Total Chlorophyll content**

109  
110  
111 SPAD chlorophyll meter was used to measure the total chlorophyll content of the leaves. Three  
112 readings were taken at three spot on a leaf of each plant and the average readings were recorded.  
113 Time interval between 9.00 a.m and 12.00 p.m was used to measure the chlorophyll content.

## 114 **2.5 Leaf gas exchange measurement**

115  
116  
117 The leaf gas exchange measurement was obtained after week 4 the treatment was given. The result  
118 then was obtained by using the Portable Photosynthesis System machine (LICOR 6400 XT). The  
119 IRGA was firstly warm up for at least 30 minutes before the leaf gas exchange was collected with  
120 Zero IRGA mode. The optimal condition was set to 400  $\mu\text{mol mol}^{-1}$  carbon dioxide ( $\text{CO}_2$ ), 30 °C  
121 cuvette temperature, 60% relative humidity with air flow rate set at 500  $\text{cm}^3 \text{min}^{-1}$ , and 800  $\mu\text{molm}^{-2}\text{s}^{-1}$   
122 of cuvette condition of photosynthetic photon flux density (PPFD). The time for the measurement were  
123 done at the morning of a day. The measurement of photosynthesis rate were taken from the first  
124 kangkung leaves starting from the plant apex. The data then were recorded and stored in a console of  
125 the system and analyse with Photosyn Assistant Software. The photosynthesis (A), transpiration rate  
126 (E), stomata conductance (gs) and water use efficiency (WUE) data was recorded during the  
127 measurement.

## 128 **2.7 Total phenolics and flavonoids quantification**

129  
130  
131 The methods used for extraction and quantification of total phenolics and flavonoids contents followed  
132 that described in Ibrahim et al. [19]. A fixed amount of ground tissue samples (0.1 g) was extracted  
133 with 80% ethanol (10 mL) on an orbital shaker for 120 min at 50 °C. The mixture was subsequently  
134 filtered (Whatman™ No.1), and the filtrate was used for the quantification of total phenolics and total  
135 flavonoids. Folin–Ciocalteu reagent (diluted 10-fold) was used to determine total phenolics content of  
136 the leaf samples. The sample extract at 200 $\mu\text{L}$  was mixed with Folin–Ciocalteu reagent (1.5 mL) and  
137 allowed to stand at 22 °C for 5 min before adding  $\text{NaNO}_3$  solution (1.5 mL, 60  $\text{g L}^{-1}$ ). After two hours  
138 at 22 °C, absorbance was measured at 725 nm. The results were expressed as  $\text{mg g}^{-1}$  gallic acid  
139 equivalent ( $\text{mg GAE g}^{-1}$  dry sample). For total flavonoids determination, samples (1 mL) were mixed  
140 with  $\text{NaNO}_3$  (0.3 mL) in a test tube covered with aluminium foil, and left for 5 min. Then 10%  $\text{AlCl}_3$  (0.3  
141 mL) was added followed by addition of 1 M NaOH (2 mL). The absorbance was measured at 510 nm  
142 using a spectrophotometer with rutin as a standard (results expressed as  $\text{mg/g}$  rutin dry sample).

## 143 **2.8 Statistical analysis**


144  
145  
146 Data were analysed using the analysis of variance procedure in SAS version 17. Means separation  
147 between treatments was performed using Duncan multiple range test and the standard error of  
148 differences between means was calculated with the assumption that data were normally distributed  
149 and equally replicated.

150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207

### 3. RESULTS AND DISCUSSION

#### 3.1 Plant height


Figure 1 shows the plant height of *I. aquatica* as influenced by differing nitrogen treatments. **As can be seen**, the plant height of kangkung were mostly affected by different rates of nitrogen treatment in all week of measurement ( $P \leq 0.05$ ). In view of the result obtained, as nitrogen levels increased from 0 to 90 kg N/ha the plant height was enhanced in all weeks of measurement. In 4 weeks after treatment (4 WAT), plant at 0 kg/ha have the average of 31.02 cm compared to 32.17 cm by 30 kg/ha, 35.61 cm by 60 kg/ha and 37.24 cm in 90 kg/ha. Clearly, as expected, applying higher rates of nitrogen levels would enhance the plant height of *I. aquatica*. The positive effects on plant height cause by the increase of nitrogen rates application may be due to the natural role of nitrogen on vegetative growth performance of plants [6]. The increase in plant height under nitrogen fertilization might be due to well-developed stem under high nitrogen fertilization that resulted in taller plant [20]. Besides that, increase in plant height might be associated with the increased of number and length of the internodes by nitrogen [21]. The result obtained agreed with the previous work carried out by [4] and [6] where the increment of nitrogen fertilization rates applied towards *I. aquatica* had significantly increased the plant height at end of the harvesting period. It can be concluded, that in the present study, that high application of nitrogen has shown to enhance the height of *I. aquatica*.



**Fig 1. The impact of different nitrogen rates on the height of *Ipomea aquatica*. Mean with the same letter indicates that all of the groups were not significantly different according to Duncan multiple range test ( $P \geq 0.05$ ) N=10.**

#### 3.2 Leaves numbers


The variation of leaf numbers with different nitrogen fertilization in *I. aquatica* is depicted in Figure 2. Generally, leaf number of *I. aquatica* plant was found to be influenced by the different rates of nitrogen treatments (0, 30, 60 and 90 kg/ha;  $P \leq 0.05$ ). Based on Figure 2, it shows that there were significant effects of nitrogen fertilization rates on the number of leaves in every week of measurements. Overall at 90 N kg/ha as the highest treatments of nitrogen applied, lead to the drastic production in number of leaves from 1 to 4 WAT. An increase in number of leafage in plants indicate better plant growth and development. Eventually, the plant production also will increase. Similar trends were observed in [6] and [20] where they found that as the rate of nitrogen increases the *I. aquatica* leaf numbers were also enhances. The increase in leaf number in *I. aquatica* might be due to increase in internodes number with high application of nitrogen [21]. The high application of nitrogen usually would reduce the apical dominance and stimulated the development of lateral buds that eventually increase the production of plant leaf and simultaneously enhanced the leaf numbers [22].



208  
209 **Fig 2. The impact of different nitrogen rates on the leaves numbers of *Ipomea aquatica*. Mean**  
210 **with the same letter indicates that all of the groups were not significantly different according**  
211 **to Duncan multiple range test ( $P \geq 0.05$ ) N=10.**  
212  
213

### 214 3.3 Plant Total dry biomass

215  
216 Nitrogen application had significantly influenced on the total plant dry weight of *I. aquatica* plant as  
217 shown in Figure 3. The graph pattern show increased in production in total biomass with higher  
218 application of nitrogen fertilization rates. At end of the treatments, It was observed that the highest  
219 total biomass of kangkung was obtained in 90 kg N/ha, followed by 60 kg N/ha and 30 N kg N/ha that  
220 recorded at 3.7g and 3.26g respectively. The lowest total biomass was recorded in control treatment 0  
221 kg N/ha that just recorded 3.13g. The increase of total plant biomass with increasing nitrogen levels  
222 can be explained by the increase in plant sink strength with increasing nitrogen levels. As nitrogen  
223 uptake increased, more of accumulation of dry biomass will be expected due to increase in plant sink  
224 strength that can accommodate initiation of new plant sink There were no significant different  
225 occurred in between 0 and 30 N kg/ha treatment ( $p \geq 0.05$ ). The result of the present study was in  
226 agreement with the research conducted by [23] where, they found that the dry weight of shoot  
227 increased with the increase of nitrogen supplied in *Ipomea aquatica*. This justify that high availability  
228 of nitrogen was important in increasing the dry biomass of *I. aquatica* that was observed in the  
229 present study [24,25].  
230  
231




234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253

**Fig. 3. The impact of different nitrogen rates on total biomass of *Ipomea aquatica*. Mean with the same letter indicates that all of the groups were not significantly different according to Duncan multiple range test ( $p \geq 0.05$ ) N=10.**

### 3.4 Number of branches

The Figure 4 below shows the branches number of kangkung plant as affected by nitrogen treatments in all four weeks of treatment. As the higher rate of nitrogen treatments, the branching of plants was enhanced. At the first 2 weeks after the treatments were applied, the number of branches at 60 N kg/ha were higher than plants that were applied with 90 N kg/ha. But then, at week 3 and 4, the opposite results were obtained where the highest number of branches occurred at 90 N kg/ha. The study was in agreement with findings by Nashrin et al. [6] on *Ipomea aquatica*, where the highest branching was obtained under highest nitrogen fertilization. Also, Osman and Abo Hassan [26], observed increased branching of Mangrove as nitrogen rate was increased to 100 kg N/ha. The increased in branching of the plant under high nitrogen fertilization might be due to increase in apical branches with higher nitrogen fertilization. This was due to enhanced vegetative growth under high nitrogen fertilization that enhanced the branching abilities of the plant [27].



254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278

**Fig. 4. The impact of different nitrogen rates on the branch number of *Ipomea aquatica*. Mean with the same letter indicates that all of the groups were not significantly different according to Duncan multiple range test ( $p \geq 0.05$ ) N=10.**

### 3.5 Total Chlorophyll Content

Figure 5 showed the impact of nitrogen fertilization on total chlorophyll content (TCC) of *I. aquatica* in 4 weeks of treatments. There were significant differences was observed for TCC in every weeks of measurement. The chlorophyll content increased after week 1 and reached it's maximum WAT content at week 3 as shown in Figure 5. In 1 WAT to 4 WAT, As the rate increased from 0 to 90 kg/ha, The TCC was steadily enhanced with the increasing nitrogen rates. In 2 -4 WAT there were no significant difference observed between 60 and 90 kg/ha in TCC. The study was in agreement with findings of According to Bojović and Marković [28] where higher application of nitrogen increased the TCC in wheat, where establishes a linear relationship between the rates of nitrogen and the chlorophyll content in plants. The plant that has been treated under high N level will resulted in higher chlorophyll content where this might be due to the immediate absorbance of nitrogen in plant [29]. Since N is important for the structural element of chlorophyll and protein molecules, low N level will affect the formation of chloroplasts and the accumulation of chlorophyll in the plant [22]. Furthermore, as the plant age increased or getting mature, the N level tend to decrease and get mobilized to other part of the plant [29]. It can be concluded that in the present study, the higher rates of nitrogen application have increases the TCC in *I. aquatica*.

279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320





Fig. 5. The impact of different nitrogen rates on the total chlorophyll content of *Ipomea aquatica*. Mean with the same letter indicates that all of the groups were not significantly different according to Duncan multiple range test ( $P \geq 0.05$ )  $N=10$ .

### 3.6 Photosynthesis rate

The photosynthesis rate of *I. aquatica* was affected by four different nitrogen treatments. It is clearly observed that from the graph pattern, as the nitrogen rate fertilization become higher ( $0 > 90$  kg/ha), the rate of photosynthesis also enhances (Figure 6). The highest A was observed in 90 kg/ha nitrogen, followed by 60 and 30 kg/ha, with the means of 3.91, 3.42, and 2.69  $\mu\text{mol}/\text{m}^2/\text{s}$  respectively. The lowest A was observed in 0 kg/ha where it just recorded 2.31  $\mu\text{mol}/\text{m}^2/\text{s}$ . The increase in A under high nitrogen level might be due to increases in leaf area that correspondingly enhanced photosynthetic activity per plant [30]. The result was also in agreement with Boussadia et al. [31] where higher nitrogen content have shown to enhanced the photosynthesis rate in olive plants. The nitrogen and photosynthesis activity is linked together because of the Calvin Cycle protein which represent the nitrogen in leaf [32]. At lower N level, the rate of photosynthesis was low. This might be due to the greater resistance and low biochemical of chloroplast [33]. According to Makino et al. [34] the increase in rate of nitrogen leads to a greater N allocation to Rubisco. Rubisco is the primary  $\text{CO}_2$  for enzyme fixation where the amount of this enzyme can drastically affect the photosynthesis rate. Besides, high N is needed in Rubisco protein due to the low rate of catalysis in Rubisco. It can be concluded that, enhanced application of nitrogen would enhance rubisco production that enhanced the net photosynthesis of *Ipomea aquatica* that was observed in the present study.




321

322 **Fig. 6. The impact of different nitrogen rates on the photosynthesis rate of *Ipomea***  
323 ***aquatica*. Mean with the same letter indicates that all of the groups were not**  
324 **significantly different according to Duncan multiple range test ( $P \geq 0.05$ ) N=10.**  
325  
326  
327

### 328 3.7 Stomatal conductance

329  
330 Stomatal conductance can be defined as the rate of carbon dioxide uptake and the water loss through  
331 stomatal leaves [35]. Based on Figure 7 below, it is distinctly observed that different rates of nitrogen  
332 had greatly affected the measurement of stomatal conductance. The higher the treatment  
333 concentrations (0,30,60,90 kg/ha), the rate of stomatal conductance have shown to increased. The  
334 stomatal conductance measurement was the highest at 90 N kg/ha ( $1.15 \text{ mmol m}^{-2} \text{ s}^{-1}$ ), while the  
335 lowest rate of stomatal conductance was measured at 0 kg/ha nitrogen treatment that recorded  $0.33$   
336  $\text{mmol m}^{-2} \text{ s}^{-1}$ . The present result was in agreement with the findings of [36], where they found that the  
337 increase in photosynthesis rate and stomatal conductance are correlated to increase in nitrogen  
338 application to the plants. Despite nitrogen, the size of the leaf can be important for certain plant  
339 species as it helps for greater conductance through the high number opening of the stomata [37]. This  
340 indicate that stomata conductance was enhanced with high levels of nitrogen applied to *I. aquatica*.  
341  
342  
343  
344




345  
346  
347  
348 **Fig. 7. The impact of different nitrogen rates on the stomatal conductance of *Ipomea aquatica*.**  
349 **Mean with the same letter indicates that all of the groups were not significantly different**  
350 **according to Duncan multiple range test ( $P \geq 0.05$ ) N=10.**  
351  
352

### 353 3.8 Water use efficiency (WUE)

354  
355 Water use efficiency (WUE) was illustrated in Figure 8 as it was influenced by the nitrogen treatments  
356 ( $P \leq 0.05$ ). Plant with the highest concentration of nitrogen (90 kg/ha) has the lowest measurement  
357 recorded in water use efficiency with the mean of  $1.46 \mu\text{mol CO}_2/\text{H}_2\text{O}$  transpired. While the highest  
358 measurement in water use efficiency was recorded in plant that was applied with 90 Kg/Ha nitrogen  
359 with mean of  $1.97 \mu\text{mol CO}_2/\text{H}_2\text{O}$  transpired. The current result was contradicting with the findings of  
360 Stewart [38] in cotton where the highest nitrogen application has shown to enhanced the WUE in the  
361 plant. The increased of WUE is usually, attributed to the increase of the transpiration rate and showed  
362 plant under water stress condition. The current result showed that higher application of nitrogen rates  
363 in *I. aquatica* can reduce the plant stress by having lower WUE. [22]. A similar result was obtained by  
364 Artur et al. [39] where the increase of N has reduced the WUE in Marandu grass that showed high  
365 application of nitrogen can reduce stress in *I. aquatica*.  
366  
367






368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390

**Fig. 8.** The impact of different nitrogen rates on the water use efficiency of *Ipomea aquatica*. Mean with the same letter indicates that all of the groups were not significantly different according to Duncan multiple range test ( $P \geq 0.05$ )  $N=10$ .

### 3.9 Total phenolics

Total plant phenolics contents was influenced with nitrogen fertilization ( $P \leq 0.05$ ; Figure 9). As levels of nitrogen enhanced, the total phenolics content was seemed to be reduced. Total phenolics was 203%, 41% and 13% higher in 30 kg/ha, 60 kg/ha and 0 kg/ha respectively compared to 90 kg/ha treatments. Previous study had showed that when the level of nitrogen decreased, the phenolic compound increased in Broccoli [40]. Another result obtained by Stewart et al. [41], also prove that the phenolic content increased as the plant faced deficiency in nitrogen level. The result obtained in this study suggested that at lower nitrogen fertilization i.e. 30 kg N/ha the production of total phenolics in *Ipomea aquatica* was enhanced. According [42], when a plant undergo N deficiency, the process of distributing carbon-based secondary compounds will increase, thus, decreasing the synthesis of nitrogen-based secondary compounds. Besides, Ibrahim et al. [19] stated that the increased in total phenolics production under low N level also might be due to the increase of total carbohydrate structural production that enhanced the production of carbon based secondary metabolites.




391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403

**Fig. 9.** The impact of different nitrogen rates on total phenolics of *Ipomea aquatica*. Mean with the same letter indicates that all of the groups were not significantly different according to Duncan multiple range test ( $P \geq 0.05$ )  $N=4$ .

### 3.10 Total flavanoids

The total flavanoids of *Ipomea aquatica* was observed to be affected by the different rates of nitrogen treatments (Figure 10;  $P \leq 0.05$ ). The production of total flavanoids has the same trends with total phenolics production content where plants which applied with 30 N kg/ha treatments has the highest total flavanoids content (1.05 mg Rutin/g dry weight) compared to 90 kg/ha that only recorded 0.27

404 mg rutin/ g dry weight. The same observation was obtained by [43] (2012) in Yaupon where the  
 405 flavonoid content reduces when applied with high N rate. According to [44] the flavonoids content in  
 406 plant tissues can be increased when having lower nitrogen content in the plant tissues. The increases  
 407 in synthesis of flavonoid at lower nitrogen level might be due to increases in phenylalanine availability  
 408 that enhances the pheyllaline lyase activity that simultaneously enhanced the production of secondary  
 409 metabolites [45]. It can be concluded in the present study, that under high nitrogen level the  
 410 production of total phenolics and flavonoids was reduced in *I. aquatica*.  
 411  
 412



413  
 414  
 415  
 416 **Fig.10.The impact of different nitrogen rates on total flavonoids of *Ipomea***  
 417 ***aquatica*. Mean with the same letter indicates that all of the groups were not**  
 418 **significantly different according to Duncan multiple range test ( $P \geq 0.05$ ) N=4.**  
 419  
 420

421 **4. CONCLUSION**

422  
 423 In this work, four levels of nitrogen rates (0, 30, 60 and 90 kg/ha) was applied to *I. aquatica* to assess  
 424 the growth, leaf gas exchange and production of secondary metabolites characteristics. It was found  
 425 that as the nitrogen rates increased, the growth and leaf gas exchange properties of *I. aquatica* was  
 426 enhanced. However, the production of phenolics and flavonoids of kangkung was reduced with high  
 427 levels of nitrogen application as both total phenolics and flavonoid reached highest content at 30 kg N  
 428 /ha. This work gives support that high nitrogen fertilization to *I. aquatica* can reduces the production of  
 429 secondary metabolites although the growth parameters was enhanced with high nitrogen fertilization.  
 430

431 **REFERENCES**

432  
 433 1. Malaysia Department of Agriculture. Ringkasan Maklumat Perangkaan  
 434 Agromakanan. www.moa.gov.my.2011  
 435  
 436 2. Ismail A, Marjan ZM and Foong CW. Total antioxidant activity and phenolic contents in  
 437 selected vegetables. Food Chem. 2004;87: 581–6.  
 438  
 439 3. Edie EE and Ho BWC. Ipomoea aquatica as a vegetable crop in Hong Kong. Econ. Bot. 1969;  
 440 23:32–6.  
 441  
 442 4. Nashrin S, Farooque A, Siddiqua M, Rahman M, and Khanam M. Effect of nitrogen and  
 443 spacing on the growth and yield of gimakalmi, *Ipomea reptans* poir. J. Biol. Sci. 2002;76: 170-  
 174.  
 444  
 445 5. Candlish I K, Gourley L and Lee HP. Dietary fiber and starch contents of some Southeast  
 Asian vegetables. J. Agri. Food Chem. 1987; 35: 319–21.

- 444 6. Susila A, Prasetyo T, and Palada M. Optimum Fertilizer rate for kangkong (*Ipomoea reptans*  
445 L.) Production in Ultisols Nanggung.2008;34:12-34.
- 446 7. Ayodele O, Alabi E, and Aluko M. Nitrogen fertilizer effect on growth, yield, and chemical  
447 composition of hot pepper (*rodo*). *Int. Agri. Crop Sci.* 2015;8(5): 666-673.
- 448 8. Li J, Zhu Z, and Gerendas J. Effects of nitrogen and sulfur on total phenolics and antioxidant  
449 activity in two genotypes of leaf mustard. *J. Plant Nutr.* 2008; 31: 1642-1655.
- 450 9. Mokhele B, Zhan X, Yang G, and Zhang X. Nitrogen assimilation in crop plants and its  
451 affecting factors. *Canadian Journal of Plant Science.* 2011;17: 231-245.
- 452 10. Zaman MS, Hashem MA, Jahiruddin M, and Rahim MA. Effect of nitrogen for yield  
453 maximization of garlic in old Brahmaputra flood plain soil. *Bangla. J. Agri. Res.*2011; 36(2):  
454 357-367.
- 455 11. Mansour MMF. Nitrogen containing compounds and adaptation of plants to salinity stress.  
456 *Biol. Plantarum.*2000; 43: 491–500.
- 457 12. Grusak MA, Della PD and Welch RM. Physiological process affecting the content and  
458 distribution of phytonutrients in plants. *Nut. Rev.* 1999; 57(9): S27–S33.
- 459 13. Campostrini E, Yamanishi O, and Martinez E. Leaf Gas Exchange Characteristics of Four  
460 Papaya Genotypes During Different Stages of Development. *Revista Brasileira de*  
461 *Fruticultura.* 2001; 23(3): 522-525.
- 462 14. Hajiboland R and Beiramzadeh N. Growth, gas exchange and function of antioxidant defense  
463 system in two contrasting rice genotypes under Zn and Fe deficiency and hypoxia. *Acta Biol.*  
464 *Szegediensis* 2008; 52: 283–294.
- 465 15. Stiller V, Lafitte HR, Sperry JS. Hydraulic properties of rice and the response of gas exchange  
466 to water stress. *Plant Physiol.* 2003; 132: 1698–1706.
- 467 16. Dingkuhn M; Cruz RT, O'Toole JC, Turner NC, Doerffling K. Responses of seven diverse rice  
468 cultivars to water deficits. III. Accumulation of abscisic acid and proline in relation to leaf  
469 water-potential and osmotic adjustment. *Field Crops Res.* 1991; 27: 103–117.
- 470 17. Dixon RA and Paiva NL. Stress-induced phenylpropanoid metabolism. *Plant Cell.* 1995; 7: 1  
471 085–1097.
- 472 18. Due BM, Humphries D, Mai LTB, Dao AH, Co TM, Nga HH and Kim PT. Iron and vitamin C  
473 content of commonly consumed foods in Vietnam. *Asia-Pacific Journal of Clinical Nutrition.*  
474 1999; 8: 36–38.
- 475  
476 19. Ibrahim MH, Jaafar HZE, Rahmant A, and Rahman Z. Effects of Nitrogen fertilization on  
477 synthesis of primary and secondary metabolites in three varieties of kacip Fatimah (*Labisia*  
478 *pumila blume*). *Int. J. Mol Sci.* .2011; 12(8): 5238-5254.
- 479 20. Taheri E, Soleymani A, and Javanmard HR. Effect of different levels of nitrogen on  
480 morphological traits of two cultivars of rapeseed in isfahan region. *Int. J. Agri Crop Sci.*  
481 2012;35: 1587-1590.
- 482  
483 21. Amin MEM. Effect of different nitrogen sources on growth, yield and quality of fodder maize  
484 (*Zea mays L.*). *J. Saudi Soc. Agri. Sci.* 2010; 67: 17-23.
- 485  
486 22. Sarkar R, Jana J and Datta, S. Effect of cutting frequencies and nitrogen levels yield and  
487 quality of water spinach (*Ipomoea reptans Poir.*). *Journal of Applied and Natural Science,*  
488 2014;76: 545-551.
- 489

- 490 23. Phimmasan H, Kongvongxay S, Chhayty P and Preston T R. Water spinach (*Ipomoea*  
491 *aquatica*) and Stylo 184 (*Stylosanthesguianesis* CIAT 184) as basal diets for growing rabbits.  
492 *Livestock Res. for Rural Development*. 2004;16: 46–59.  
493
- 494 24. Hare PD, Cress WA and Staden VJ. Proline synthesis and degradation: a model system for  
495 elucidating stress-related signal transduction. *J. Exp. Bot.* 1999; 50: 413–34.  
496
- 497 25. Prasad NK, Divakar S, Shivamurthy GR and Aradhya SM. Isolation of a free radical  
498 scavenging antioxidant from water spinach (*Ipomoea aquatic* Forsk.). *J. Sci. Food Agri.* 2005;  
499 85: 1 461–8.  
500
- 501 26. Osman H, and Abu HA. Effect of NPK Fertilization on growth and dry matter accumulation in  
502 mangrove [*Avicennia marina* (Forssk) vierh] Grown in Western Saudi Arabia. *Meteorol.*  
503 *Environ. Arid Land Agri.*2010;56: 57-70.  
504
- 505 27. Koo HM and Suhaila M. Flavonoid (Myricetin, quercetin, kaempferol, luteolin and apegenin)  
506 content of edible tropical plants. *J. Agri. Food Chem.* 2011; 49: 3 106– 12.  
507
- 508 28. Bojovic B and Markoviae A. Correlation Between Nitrogen and Chlorophyll Content In Wheat  
509 (*Triticum aestivum* L.). *Kragujevac J. Sci.*2009; 3: 69-74.  
510
- 511 29. Moreno N, Barrios A, Leal R, Franco A, Rodriguez A, and Hernandez L. Effect of Nitrogen  
512 Deficiency and Toxicity in Two Varieties of Tomato (*Lycopersicum esculentum* L.). *Agri.*  
513 *Sci.*2014; 5: 1361-1368.  
514
- 515 30. Mansour MMF. Nitrogen containing compounds and adaptation of plants to salinity stress.  
516 *Biologia Plantarum.* 2000; 43: 491–500.  
517
- 518 31. Boussadia K, Steppe K, Labeke MC, Lemeur R and Braham M. Effects of Nitrogen deficiency  
519 on Leaf Chlorophyll Fluorescence Parameters in Two Olive Tree Cultivars Meski' and  
520 'Koroneiki'. *J. Plant Nutr.*2015;76; 2230-2246.  
521
- 522 32. Baque MA, Karim MA, Hamid A, Tetsushi H. Effects of fertilizer potassium on growth, yield  
523 nutrient uptake of wheat (*Triticum aestivum*) under water stress conditions. *South Pac.*  
524 *Studies* 2006; 27: 29–35.  
525
- 526 33. Grzebisz W, Gransee A, Szczepaniak W, Diatta J. The effects of potassium fertilization on  
527 water-use efficiency in crop plants. *J. Plant Nutr. Soil Sci.* 2013: 176: 355–374.  
528
- 529 34. Makino, A. Photosynthesis, Grain Yield, and Nitrogen Utilization in Rice and Wheat. *Plant*  
530 *Physiology.* 2011; 155: 125-129.  
531
- 532 35. Pankovica D, Plesnicar M, Arsenijevicâ MI, Petrovica N, Sakaci Z, and Kastori R. Effects Of  
533 Nitrogen nutrition on photosynthesis in cd-treated sunflower plants. *Annals of Bot.*,2000; 86:  
534 841-847.  
535
- 536 36. Nori M, Bayat F, and Esmaili A. Changes of vegetative growth indices and yield of garlic  
537 (*Allium sativum* L.) in different sources and levels of nitrogen fertilizer. *Int. J. Agri. Crop*  
538 *Sci.*2012;67; 1394-1400.  
539
- 540 37. Schulze ED, Kelliher F, Korner C, Lloyd J, and Leuning R. Relationships among maximum  
541 stomatal conductance, ecosystem surface conductance, carbon assimilation rate, and plant  
542 nitrogen nutrition: a global ecology scaling exercise. *Ann. Rev. of Ecol. Systematics.*1994; 25:  
543 629-660.  
544
- 545 38. Stewart W. Balanced fertilization increases water use efficiency. Atlanta, Georgia: Potash &  
546 Phosphate Institute (PPI) and Potash & Phosphate Institute of Canada (PPIC).2001.  
547
- 548 39. Artur A, Garcez T, and Monteiro F. Water use efficiency of marandu palisadegrass as  
549 affected by nitrogen and sulphur rates. *Artigo Cientifico.* 2014; 45(1): 10-17.

- 550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572
40. Naguib, AM. Enhancement of phenolics, flavonoids and glucosinolates of Broccoli (*Brassica oleracea*, var. *Italica*) as antioxidants in response to organic and bio-organic fertilizers. *Journal of the Saudi Society of Agricultural Sciences*.2013;78: 135-142.
  41. Stewart A, Chapman W, Jenkins G, Graham I, Martin T, and Crozier A. The effect of nitrogen and phosphorus deficiency on flavonol accumulation in plant tissues. *Plant Cell Environ*.2011; 24: 1189-1197.
  42. Orphanides A, Goulas V and Gekas V. Effect of drying method on the phenolic content and antioxidant capacity of spearmint. *Czech J. Food Sci*.2013; 31(5): 509-513.
  43. Palumbo, M., Putz, F., & Talcott, S. Nitrogen fertilizer and gender effects on the secondary metabolism of yaupon, a caffeine-containing north American holly. *Oecologia*.2006; 151(1): 1-9 .
  44. Meyer S, Cerovic ZG, Goulas Y, Montpied P, Demotes S, Bidet LPR, Moya I and Dreyer E. Relationship between assessed polyphenols and chlorophyll contents and leaf mass per area ratio in woody plants. *Plant Cell Environ*. 2006;29: 1338-1348.
  45. Margna U, Margna E and Vainjarv T. Influence of nitrogen nutrition on the utilization of L-phenylalanine for building flavonoids in buckwheat seedling tissue. *J. Plant Physiol*. 1989;134:697-702.