

Original Research Article

Enalapril confers protective effect on isoproterenol-induced myocardial infarction in rats through downregulation of cardiac troponin, C-reactive protein, upregulation of IL-10 β as well as anti-oxidant and anti-inflammatory activities

Abstract

Myocardial infarction is an irreversible death of heart muscle secondary due to prolonged lack of oxygen supply. The present study was designed to evaluate the protective effect of enalapril in isoproterenol-induced myocardial infarction in rats using changes in haemodynamic, biochemical, histopathological and immunohistochemistry parameters. Twenty-one male Wistar rats divided into three groups were used where the control (group A) was administered for normal saline which continued for 7 days, group B animals received normal saline for 7 days and thereafter isoproterenol (ISO) at 85 mg/kg on day 8 and 9. Group C animals were pretreated with enalapril (10mg/kg) for 7 days and thereafter received ISO on day 8 and 9. On day 10, the blood pressure change in the animals were measured and thereafter sacrificed by cervical dislocation. The heart of each rat was removed, homogenized and used to assay for some oxidative stress markers and some antioxidant parameters. In this study, ISO caused myocardial infarction as seen by significant decrease in systolic, diastolic and mean arterial pressure but was corrected by enalapril. Enalapril caused significant increase in the levels of SOD, GPx, GST and GSH but significant decrease in MDA content and H₂O₂ generation. But reverse was the case for group B animals. Immunohistochemistry showed that ISO caused higher expressions of cardiac C-reactive protein (CRP) and cardiac troponins 1 (CTn1) and decrease in IL-10 β but vice-versa for enalapril. No histopathological changes were recorded for enalapril. The study thus showed that enalapril significantly exhibits cardioprotective effects.

Key words: Enalapril, myocardial infarction, cardioprotection, immunohistochemistry, antioxidant

Introduction

Human health is being seriously threatened by cardiovascular diseases (CVD), which have been regarded as the main cause of death throughout the world [1-3]. Myocardial infarction (MI) is a common presentation of ischemic heart disease (IHD) and remains the major cause of death in

33 the developed world. Though rapid advancements have been made in the treatment of coronary
34 artery diseases (CAD), MI is still a major pathological issue worldwide [4]. Increased
35 myocardial metabolic demand and decreased supply of oxygen as well as nutrients via the
36 coronary circulation to the myocardium brings about myocardial infarction hence leading to cell
37 injury. This pathological heart condition is one of the most lethal manifestations of
38 cardiovascular diseases. Acute myocardial infarction or heart attack occurs when blood stops
39 flowing to part of the heart leading to injury to the heart muscle due to the fact the heart is not
40 receiving enough oxygen [5-9].

41 Isoproterenol [1-(3, 4-dihydroxyphenyl)-2-isopropylaminoethanol hydrochloride] (ISO) a
42 synthetic catecholamine is a β -adrenergic agonist that is very important in the regulation of
43 myocardial contractility and metabolism. It serves as a standard model for the study of
44 potentially beneficial effects of numerous drugs on cardiac function [10, 11]. ISO induces
45 myocardial injury in rat because of the alteration in the physiological balance between
46 production of free radicals and antioxidative defence system [12]. It thus causes the acute
47 condition of myocardial necrosis, which can lead to cardiac dysfunctions, increased lipid
48 peroxidation, altered activities of cardiac enzymes and antioxidants [13]. It has been observed
49 that the pathophysiological and morphological changes observed in ISO-treated rats are similar
50 to those observed in human MI [14].

51 Enalapril an Angiotensin-converting-enzyme inhibitor (ACE inhibitor) is a drug used primarily
52 for the treatment of high blood pressure and congestive heart failure where it can be used alone
53 or in combination with other antihypertensive agents. ACE inhibitors have also been found to be
54 useful for other cardiovascular and kidney diseases including acute myocardial infarction,
55 diabetic nephropathy, and cardiac failure [15]. The mechanism of action of ACE inhibitors
56 involves reduction of the activity of the renin-angiotensin-aldosterone system (RAAS) [16].

57 In recent times, a novel strategy has been employed in drug discovery. It is the use of known and
58 approved drugs and compounds for newer indications. This is termed drug repurposing. In this
59 study, Isoproterenol was used to induce acute myocardial infarction and enalapril was then used
60 to ameliorate this and then to see if it could serve as a repurposed drug for myocardial infarction.

61
62

63 **Materials and Methods**

64 *Chemicals and reagents*

65 Isopreterenol, enalapril, Tween 80, Biurett's reagent, hydrogen peroxide, hydrochloric acid,
66 sulphuric acid, xylenol orange, potassium dichromate, O-diasinidine, sodium potassium tartrate,
67 copper sulphate, ethanol, sodium azide, 2-dichloro-4-nitrobenzene (CDNB) Greiss reagent,
68 phosphoric acid, sodium hydroxide, N 1-naphthyl ethylenediamine, sulphanilamide, distilled
69 water, , phosphate buffer saline, creatinine reagent, copper sulphate, tri chloro acetate, reduced
70 glutathione (GSH), thiobarbituric Acid (TBA), trichloroacetic acid (TCA), ammonium ferrous
71 sulphate, glacial acetic acid, potassium iodide, sorbitol, Ellman's reagent (DTNB), ethanol, urea
72 reagent. All other chemicals used were of analytical grade and obtained from British Drug
73 Houses (Poole, Dorset, UK). All other chemicals, reagents and drugs used were of analytical
74 grade.

75 *Experimental animals*

76 All experiments and protocols described in present study were approved by the UI-ACUREC.
77 Twenty one (21) male Wistar rats weighing between 90 to 160g were obtained from the
78 Experimental animal unit of the Faculty of Veterinary Medicine, University of Ibadan for the
79 experiment. They were allowed free access to standard rat pellets and fresh water *ad libitum*. The
80 rats were housed in the animal house unit of the Department of Veterinary Pharmacology and
81 Toxicology, University of Ibadan with a 12 hour light duration. Pre-conditioning of the rats was
82 done for two weeks before commencement of the experiment. **The institutional approval was**
83 **given to this study and the number is UI-ACUREC/App/2016/030**

84 *Cardioprotective study*

85 The animals were randomly divided into three (3) groups with seven (7) animals in each group,
86 and the treatment was as follow: Animals in the control (group A) were administered normal
87 saline, group B; isoproterenol at 85mg/kg, while group C animals were pretreated with enalapril
88 **orally** (10mg/kg) for 7 days and thereafter administered ISO (85mg/kg) **subcutaneously** on day 8
89 and 9. Blood pressure values of all the animals were carried out on day 10. At the end of the
90 experimental period, blood samples were collected for haematology and serum chemistry before

91 the rats were sacrificed by cervical dislocation. The serum in plain bottles was rapidly
92 centrifuged at 4000 revolutions per minute (rpm) for fifteen (15) minutes and processed for
93 determination of serum myeloperoxidase, total protein, and xanthine oxidase, AST, ALT and
94 nitric oxide. The heart of each rat was carefully removed and homogenized on ice and then used
95 to assay for some oxidative stress markers and antioxidant parameters. Baseline cardiovascular
96 parameters were obtained prior to the commencement of the experiment. The equipment used
97 was a non-invasive tail cuff BP monitor, the 6-channel CODA blood pressure monitor for rats
98 and mice. The haemodynamic parameters assessed were: the systolic blood pressure (SBP),
99 diastolic blood pressure (DBP), and mean arterial pressure (MAP) and were determined
100 indirectly in nonanaesthetised rats, by tail plethysmography with the use of an
101 electrospygmomanometer (CODA, Kent Scientific, USA). The average of at least nine most
102 consistent readings, taken in the quiescent state, following acclimatization, was recorded per
103 animal.

104 Blood samples for serum chemistry were collected from the rats through retro-orbital vein after
105 which the animals were sacrificed by cervical dislocation.

106

107 *Preparation of tissue homogenate*

108 The heart tissues of the rats were harvested on ice, rinsed with normal saline and homogenized in
109 aqueous potassium buffer (0.1 M, pH 7.4) and the homogenate centrifuged at 12,000 rpm (4°C)
110 for 15 min to obtain the supernatant fraction.

111 *Determination of Biochemical assay*

112 Biuret method as described by Gornal et al [17] was used to determine the protein concentrations
113 of the various samples with a slight modification. To prevent precipitation of Cu^{2+} ions as
114 cuprous oxide potassium iodide was added to the reagent. To determine the concentration of
115 reduced glutathione the method of Beutler et al [18] was used while glutathione peroxidase
116 (GPX) activity was measured by the method of Rotruck et al. [19]. In this case, hydrogen
117 peroxide was used as substrate to oxidize reduced glutathione to oxidized glutathione (GSSG).
118 Estimation of Glutathione S-transferase (GST) was by the method of Habig et al [20] using 1-

119 chloro-2, 4-dinitrobenzene as substrate. Superoxide dismutase (SOD) assay on the other hand
120 was carried out by the method of Misra and Fridovich [21]. MDA content was measured in the
121 heart as an index of lipid peroxidation [22]. Hydrogen peroxide generation was measured using
122 Wolff's [23] method while the determination of Sulfhydryl (Thiol) content was by-the method of
123 Ellman [24]. Nitric oxide was quantified as previously described [25].

124 *Histopathology*

125 Small slices of the heart were collected in 10% buffered formalin for proper fixation and after the
126 tissues have been processed and embedded in paraffin wax, sections that were about 5-6 μ m
127 thick were made and stained with haematoxylin and eosin for histopathological examination
128 [26].

129 *Immunohistochemistry of Cardiac troponins-1, CRP and IL-10*

130 The heart tissues obtained from the rats were paraffin embedded and then used for
131 immunohistochemistry. Paraffin sections were melted at 60 °C in the oven but the dewaxing of
132 the samples in xylene was followed by passage through ethanol of decreasing concentration
133 (100-80%). Peroxidase quenching in 3% H₂O₂/methanol was carried out with subsequent antigen
134 retrieval performed by microwave heating in 0.01 M citrate buffer (pH 6.0) to boil. All the
135 sections were blocked in normal goat serum (10%, HistoMark[®], KPL, Gaithersburg MD, USA)
136 and probed with cardiac troponins 1, CRP antibody and I IL-10 β (Abclonal[®]), 1:375 for 16 h in a
137 refrigerator. Detection of bound antibody was carried out using biotinylated (goat anti-rabbit,
138 2.0 μ g/ml) secondary antibody and subsequently, streptavidin peroxidase (Horse Radish
139 Peroxidase- streptavidin) according to manufacturer's protocol (HistoMark[®], KPL, Gaithersburg
140 MD, USA).

141 Diaminobenzidine (DAB, Amresco[®], USA) was used to enhance the reaction product for 6 – 10
142 min and counterstained with high definition haematoxylin (Enzo[®], NY - USA), and was
143 thereafter dehydrated in ethanol. Once the slides were covered with cover slips, they were sealed
144 with resinous solution. The immunoreactive positive expression of CRP, cardiac troponin and
145 IL-10 β intensive regions were viewed starting from low magnification on each slice then with

146 400 × magnifications using a photo microscope (Olympus) and a digital camera (Toupcam[®],
147 Touptek Photonics, Zhejiang, China).


148 *Statistical analysis*

149 All values were expressed as mean ± standard deviation (SD). The test of significance between
150 two groups was estimated by Student's t-test. One-way Analysis of Variance (ANOVA) with
151 Tukey's post-hoc test using Graph pad prism 5.0 was also performed with p-values < 0.05
152 considered statistically significant.

153 **Results**

154 In this study, ISO caused significant decreases in the levels of SBP, DBP and MAP while
155 enalapril (ENA) caused significant increase though not to the same extent as the control (Figures
156 1-3). The results of haematological analysis showed that ISO caused significant increases in the
157 levels of WBC, PCV, MCV and MCH while ENA caused significant decrease in WBC and no
158 changes relative to ISO (Table 1). ISO also caused significant increases in the levels of AST and
159 ALT while ENA caused significant decreases in the levels of these enzymes. On the other hand,
160 while ISO caused significant decrease in the level of NO, ENA caused significant increase
161 (Table 2). ISO caused significant increases in the levels of oxidative markers such as MDA,
162 H₂O₂ and MPO while ENA caused significant decreases in the levels of these markers in a
163 similar fashion to the control (Figures 4-6). Again, while ISO caused significant decrease in the
164 levels of protein thiols and non-protein thiols, ENA caused a significant increase in the levels of
165 these molecules (Figures 7 and 8). The result also showed that ISO caused significant decrease in
166 the levels of anti-oxidant markers such as SOD, GPx, GST and GSH but reverse is the case for
167 ENA (Figures 9-12). Histopathological examinations showed that while there is severe
168 infiltration of inflammatory cells into the cardiac tissue, there was no visible lesion seen in the
169 ENA and control groups (Figure 13). The immunohistochemical analysis showed that there were
170 high expressions of cardiac troponin and CRP in ISO group but lower expression of these
171 proteins in ENA and control group (Figures 14 and 15). In the case of IL-10 β , there was low
172 expression of this protein in ISO group but higher expression in ENA and control group (Figure
173 16).

174




175

176 **Figure 1:** Effect of enalapril on SBP in isoproterenol induced myocardial infarction using rats as
177 a model. The superscript 'a' showed that ISO caused significant decrease when compared to
178 control while superscript 'b' showed significant decrease when compared with ENA (n=7).

179

180



181

182 **Figure 2:** Effect of enalapril on DBP in isoproterenol induced myocardial infarction using rats as
183 a model. The superscript 'a' showed that ISO caused significant decrease in the level of this
184 parameter compared to control while 'b' showed that ENA caused significant increase relative to
185 control and ISO groups (n=7).


186

187

188

189

190



191

192 **Figure 3:** Effect of enalapril MAP in isoproterenol-induced myocardial infarction using rats as a
193 model. The superscripts showed that ISO caused significant decrease relative to ENA and control
194 groups (n=7).

195

196

197

198

199

200

201

202

203

204 **Table 1:** Effects of enalapril on RBC, WBC, HB, PCV, MCV, MCH and MCHC in
 205 **isoproterenol-induced** myocardial infarction using rats as a model (n = 7)

206

| Parameters | Control | ISO | Enalapril |
|----------------------------|------------------|---------------------------------|------------------------------|
| RBC ($\times 10^{12}/L$) | 4.75 \pm 0.90 | 4.96 \pm 0.43 | 5.03 \pm 0.69 |
| WBC (103/ μ L) | 5.47 \pm 0.38 | 6.71 \pm 1.13 ^a | 4.68 \pm 1.68 ^b |
| HB (g/dl) | 13.33 \pm 1.40 | 15.15 \pm 1.84 | 14.95 \pm 1.62 |
| PCV (%) | 45.75 \pm 4.65 | 54.25 \pm 4.25 ^a | 50.25 \pm 3.10 |
| MCV (fl) | 83.88 \pm 9.03 | 127.33 \pm 30.12 ^a | 98.87 \pm 22.76 |
| MCH (pg) | 26.41 \pm 3.48 | 38.64 \pm 8.08 ^a | 26.05 \pm 2.25 |
| MCHC (%) | 29.97 \pm 2.05 | 27.41 \pm 2.38 | 30.79 \pm 2.37 |

207 Values are mean \pm SD, n =5, ^a - **p < 0.05** compared with control, ^b - **p < 0.05** compared with ISO.

208 The superscript (a) showed that ISO caused significant decrease in the level of this parameter
 209 compared to control while (b) showed that ENA caused significant increase relative to control
 210 and ISO groups.

211

212

213

214

215

216

217

218

219

220

221 **Table 2:** Effects of enalapril on ALT, AST and NO in **isoproterenol-induced** myocardial
222 infarction using rats as a model (n=7).

| Parameters | Control | ISO | Enalapril |
|------------|------------|-------------------------|--------------------------|
| ALT | 14.51±0.02 | 14.67±0.05 ^a | 14.41±0.05 ^{ab} |
| AST | 19.91±0.01 | 19.97±0.02 ^a | 19.87±0.02 ^{ab} |
| NO | 4.11±0.68 | 1.72±0.47 ^a | 2.67±0.71 ^{ab} |

223
224 Values are mean ± SD, n =5, ^a - **p< 0.05** compared with control, ^{ab} - **p< 0.05** compared with ISO.
225 The superscript ‘a’ showed that ISO caused significant decrease in the level of this parameter
226 compared to control while ‘b’ showed that ENA caused significant increase relative to control
227 and ISO groups.

228

229


230

231

232

233

234



235

236

237 **Figure 4:** Effect of enalapril on lipid peroxidation in isoproterenol-induced myocardial
 238 infarction using rats as a model (n=5). Values are presented as mean \pm standard deviation. Grp A
 239 (Control), Grp B (ISO treated only), Grp C (ENA+ 85 mg/kg ISO). The superscript (a) showed
 240 that ISO caused significant decrease in the level of this parameter compared to control while (b)
 241 showed that ENA caused significant increase relative to control and ISO groups.

242

243

244

245


246

247

248

249

250



251

252

253 **Figure 5:** Effect of enalapril on hydrogen peroxide generation in isoproterenol-induced
 254 myocardial infarction using rats as a model (n=5). Values are presented as mean \pm standard
 255 deviation. Superscript 'a' indicates significant difference ($p < 0.05$) when compared with control
 256 (Grp A), whereas superscript 'b' indicates significant difference ($p < 0.05$) when compared with
 257 ISO treated only (Grp B). Grp A (Control), Grp B (ISO treated only), Grp C (ENA+ 85 mg/kg
 258 ISO).


259

260

261

262

263



264

265

266 **Figure 6:** Effect of enalapril on myeloperoxidase in **isoproterenol-induced** myocardial infarction
267 using rats as a model (n=5). The superscript 'a' showed that ISO caused significant increase in
268 the level of this parameter when compared to the control and ENA groups.

269

270

271

272

273

274

275


276

277

278

279

280



281

282

283

284 **Figure 7:** Effect of enalapril on protein thiol in **isoproterenol-induced** myocardial infarction
285 using rats as a model (n=5). Values are presented as mean \pm standard deviation. Superscript 'a'
286 indicates significant difference ($p<0.05$) when compared with control (Grp A), whereas
287 superscript 'b' indicates significant difference ($p<0.05$) when compared with ISO treated only
288 (Grp B). Grp A (Control), Grp B (ISO treated only), Grp C (ENA+ 85 mg/kg ISO).

289

290


291

292

293

294

295



296


297

298 **Figure 8:** Effect of enalapril on non-protein thiol in **isoproterenol-induced** myocardial infarction
 299 using rats as a model (n=5). Values are presented as mean \pm standard deviation. Superscript 'a'
 300 indicates significant difference ($p<0.05$) when compared with control (Grp A), whereas
 301 superscript 'b' indicates significant difference ($p<0.05$) when compared with ISO treated only
 302 (Grp B). Grp A (Control), Grp B (ISO treated only), Grp C (ENA+ 85 mg/kg ISO).

303

304


305



306

307 **Figure 9:** Effect of enalapril on superoxide dismutase enzyme in **isoproterenol-induced**
 308 myocardial infarction (n=5). Values are presented as mean \pm standard deviation. Superscript 'a'
 309 indicates significant difference ($p < 0.05$) when compared with control (Grp A), whereas
 310 superscript 'b' indicates significant difference ($p < 0.05$) when compared with ISO treated only
 311 (Grp B). Grp A (Control), Grp B (ISO treated only), Grp C (ENA+ 85 mg/kg ISO).

312
 313
 314
 315
 316
 317
 318
 319
 320
 321
 322



323

324


325 **Figure 10:** Effect of enalapril on glutathione peroxidase enzyme in **isoproterenol-induced**
 326 myocardial infarction using rats as a model (n=5). Values are presented as mean \pm standard
 327 deviation. Superscript 'a' indicates significant difference ($p < 0.05$) when compared with control
 328 (Grp A), whereas superscript 'b' indicates significant difference ($p < 0.05$) when compared with
 329 ISO treated only (Grp B). Grp A (Control), Grp B (ISO treated only), Grp C (ENA+ 85 mg/kg
 330 ISO).

331

332

333

334



335

336 **Figure 11:** Effect of enalapril on glutathione-s- transferase enzyme in isoproterenol-induced
 337 myocardial infarction using rats as a model (n=5). Values are presented as mean \pm standard
 338 deviation. Superscript 'a' indicates significant difference ($p<0.05$) when compared with control
 339 (Grp A), whereas superscript 'b' indicates significant difference ($p<0.05$) when compared with
 340 ISO treated only (Grp B). Grp A (Control), Grp B (ISO treated only), Grp C (ENA+ 85 mg/kg
 341 ISO).

342


343

344

345

346

347



348
349

350

351 **Figure 12:** Effect of enalapril on reduced glutathione in isoproterenol-induced myocardial
 352 infarction using rats as a model (n=5). Values are presented as mean ± standard deviation.
 353 Superscript 'a' indicates significant difference ($p<0.05$) when compared with control (Grp A),
 354 whereas superscript 'b' indicates significant difference ($p<0.05$) when compared with ISO
 355 treated only (Grp B). Grp A (Control), Grp B (ISO treated only), Grp C (ENA+ 85 mg/kg ISO).

356
357
358
359
360
361
362
363
364
365


366

367

368

369

370



371

372

373 **Figure 13:** The photomicrograph of heart from isoproterenol-induced myocardial infarction
374 using rats as a model. A (Control) shows no visible lesion. B (ISO): shows severe infiltration of
375 inflammatory cells. C (enalapril) shows no visible lesion. The slides were with H & E. Mag.
376 ×400

377


378

379

380

381

382




383

384 **Figure 14:** Immunohistochemistry of cardiac troponin in heart of isoproterenol induced
385 myocardial infarction rats. A (Control): show positive and low expression of CTnI, B (ISO):
386 shows higher expression of CTnI than control, C (enalapril) shows lower expression of CTnI
387 than B (ISO). The slides were counterstained with high definition haematoxylin. Mag. x100

388

389

390



391

392

393 **Figure 15:** Immunohistochemistry of c- reacting protein in heart of isoproterenol induced
394 myocardial infarction rats. A (Control): show positive and low expression of CRP, B (ISO):
395 shows higher expression of CRP than control, C (enalapril) shows lower expression of CRP than
396 B (ISO). The slides were counterstained with high definition haematoxylin. Mag. x100

397

398


399

400

401

402

403



404

405

406 **Figure 16:** Immunohistochemistry of interleukin-10 in heart of isoproterenol induced myocardial
407 infarction rats. A (Control): show positive and higher expression of IL-10, B (ISO): shows lower
408 expression of IL-10 than control, C (enalapril) shows higher expression of IL-10 than B (ISO).
409 The slides were counterstained with high definition haematoxylin. Mag. x100

410

411

412

413

414

415

416

417

418 **Discussion**

419 Myocardial infarction (MI), one of the main causes of death from cardiovascular disease is
420 defined as an acute condition of necrosis of the myocardium and it occurs as a result of
421 imbalance between coronary blood supply and myocardial demand [27]. MI is known to cause
422 local inflammation and apoptosis and this can result in cardiomyocyte damage [28]. ISO induces
423 cardiac necrosis by several mechanisms, including increased oxygen consumption, poor oxygen
424 utilization, increased calcium overload and accumulation, altered myocardial cell metabolism,
425 increased myocardial cAMP levels, deranged electrolyte milieu, altered membrane permeability,
426 intracellular acidosis, and increased levels of lipid peroxides [11]. The pathophysiological
427 changes that occurred in heart following isoproterenol administration in rats are comparable to
428 those taking place in human myocardial infarction [29].

429 Angiotensin converting enzyme inhibitors are known to prevent both the generation of the potent
430 vasoconstrictor angiotensin II and degradation of the powerful vasodilator bradykinin, which
431 promotes endothelial cell release of NO [30]. In this study, rats treated with ISO had significant
432 decreases in blood pressure parameters (SBD, DBP and MAP) when compared with the controls.
433 This was however prevented in the ENA-treated group. There have been earlier reports of
434 hypotension in subjects with acute myocardial infarction [31, 32]. From this study, it was
435 interesting to observe that ENA, a known antihypertensive drug, was able to preserve the blood
436 pressure measurements of ISO-treated rats comparable to the controls. This might have been a
437 consequence of its ability to prevent myocardial infarction. Studies have actually shown that
438 ACEIs have been used in the management of myocardial infarction [33, 34, 35]. Isoproterenol, a
439 β -adrenergic agonist is known to produce stress in the myocardium due to the generation of free
440 radicals by its auto-oxidation. Some of the mechanisms proposed to explain its damage to
441 cardiac myocytes include coronary hypotension, calcium overload, hypoxia, energy depletion
442 and excessive production of free radicals as a result of catecholamine autoxidation [36, 37, 38].
443 The significant decrease in the levels of systolic, diastolic and mean arterial pressure may lead to
444 coronary hypotension as seen in this study. In a study by Owens and O'Brien [39], it was
445 concluded that in patients suffering with ischaemic heart disease and hypotension, symptomatic
446 and silent ischaemia occurred in a temporally causal relation with hypotension, particularly for
447 diastolic pressures. It thus suggests that patients with coronary disease may be susceptible to

448 ischaemic events that could be incurred as a result of low blood pressure. The enalapril used in
449 this study was able to restore the haemodynamic changes caused by isoproterenol indicating **its**
450 **ability to protect against establishment of myocardial infarction.**

451 In this study, the results of haematological analysis showed that ISO caused significant increase
452 in the levels of WBC, PCV, MCV and MCH while ENA caused significant decrease in WBC
453 and no changes in the erythrocyte indices relative to control. The increase in the level of WBC
454 could be explained in terms of necrosis caused by ISO leading to white blood cell mobilization
455 [11]. The significant reduction in the level of this parameter by enalapril could also be seen as its
456 ability to counteract the toxic effect of isoproterenol.

457 **The toxicant also caused significant increase in the levels of AST and ALT while ENA caused**
458 **significant decrease in the levels of these enzymes. In heart failure, the heart has an impaired**
459 **ability to deliver blood to the body and may in the process affects the kidney and liver. The liver**
460 **can become dysfunctional, and liver enzymes can be released into the blood [40]. It thus means**
461 **that the increases noted for the liver enzymes in this study implied that isoproterenol could**
462 **impair liver functions and this was counteracted by enalapril indicating that enalapril has**
463 **beneficial effect beyond being an ACE inhibitor.**

464 It was also observed that ISO caused significant decrease in the level of NO while ENA caused
465 significant increase. Nitric oxide (NO) is known to play important functional roles in a variety of
466 physiological systems. For instance within the vasculature, NO induces vasodilation, inhibits
467 platelet aggregation, prevents neutrophil/platelet adhesion to endothelial cells, inhibits smooth
468 muscle cell proliferation and migration, regulates programmed cell death (apoptosis) and
469 maintains endothelial cell barrier function [41]. Nitric oxide (NO) is known to be deficient in
470 chronic progressive renal disease (CRD) and in end-stage renal disease (ESRD) [42, 43] and this
471 could result from arginine deficiency [44] which may be caused by a loss of functional renal
472 mass, increased endogenous NO synthase (NOS) inhibitors that accumulate in renal failure [44],
473 and/or other causes, such as increased oxidant stress [45]. Low NO production may also
474 contribute to and/or exacerbate the progression of CRD by both hemodynamic and renal growth-
475 promoting actions [46]. It should also be noted that NO blockade can lead to increased blood
476 pressure and attenuated or delayed the hypotensive effect of all ACE inhibitors [47]. ACE
477 inhibitors such as enalapril also augment the hemodynamic vasodilator action of bradykinin [48].

478 The increased level of NO in this study due to enalapril may further affirm its antihypertensive
479 property and hence cardioprotective effect.

480 ISO caused significant increase in the levels of oxidative stress markers such as MDA, H₂O₂ and
481 MPO while ENA caused significant decrease in the levels of these markers in a similar fashion to
482 the control. Oxidative stress constitutes an alteration produced by disequilibrium between
483 generation of free radicals (FR) and the antioxidant system, which can lead to a damage state, in
484 particular of the biomolecules [49, 50, 51, 52, 53]. FR generates the lipid peroxidation process in
485 an organism with malondialdehyde (MDA) level used as a marker of oxidative stress [54].
486 Myeloperoxidase (MPO) is abundant in the granules of inflammatory cells and it is an important
487 enzyme in the generation of reactive oxygen species (ROS) [55, 56, 57]. Hydrogen peroxide
488 (H₂O₂), an ROS marker has been suggested as a mediator of vascular structural and functional
489 alterations observed in hypertension [58, 59, 60, 61, 62]. The reduction of these oxidative
490 markers by enalapril is a pointer to its ability to scavenge the radicals generated by the toxicant
491 and it thus showed that enalapril has anti-oxidant activity. In fact, De Cavanagh et al [63]
492 reported that enalapril inhibits free radical formation and attenuates oxidative stress and also
493 prevents damage to the liver and kidney. This was further confirmed by the ability of this ACE
494 inhibitor to increase the levels of antioxidant enzymes such as SOD, GPx, GST and GSH
495 evaluated in this study. This view is clearly supported by a study carried out by Chandra et al
496 [64], where it was concluded that enalapril has anti-oxidative property and this may have been
497 responsible for its cardioprotective property. As a matter of fact, ENA caused a significant
498 increase in the levels of protein thiols and non-protein thiols further confirming its anti-oxidant
499 property. The thiol compounds function in the maintenance of cellular redox balance and their
500 play important role in controlling oxidative stress [65, 66, 67].

501 Cells have evolved several antioxidant strategies aimed at the detoxification of ROS with
502 glutathione redox cycle as one of the major protective systems against oxidant damage. This
503 cycle composed of the enzymes glutathione peroxidase (GPx) and glutathione reductase (GSSG-
504 Rd) and the co-substrates glutathione and NADPH [68]. Glutathione is the most abundant non-
505 protein intracellular thiol, and has a multiple role as an antioxidant agent [69]. Though the
506 mechanism(s) underlying the enhancement of glutathione and glutathione-related enzymes by

507 ACEI remains unknown, however, tissue glutathione levels and GSSG-Rd and GPx activities
508 have been shown to increase in response to experimentally induced oxidative stress [70].

509 In this study, histopathological examinations showed that while there was severe infiltration of
510 inflammatory cells into the cardiac tissue of the ISO group, there was no visible lesion seen in
511 the ENA and control groups (Figure 13). This increase in the inflammatory cells may have been
512 responsible for the increase in the levels of WBC noted in this study (Table 1). It should be noted
513 that the isoproterenol-induced myocardial alterations are similar in certain respects to those
514 occurring in human beings following a myocardial infarction [71]. **It is thought that the β -
515 adrenergic cardiostimulatory activity exerted by ISO increases cardiac oxidative metabolism to a
516 level that exceeds the amount of oxygen available to the myocytes through the unobstructed
517 coronary circulation. The area of the heart most susceptible to hypoxia caused by tachycardia
518 appears to be the left ventricular subendocardium [72, 73]. Myocyte damage observed following
519 exposure to ISO includes both apoptosis and necrosis [74]. In the study on the isoproterenol-
520 induced myocardial damage, it was discovered that the cardiac lesions varied with treatment
521 duration and doses and that numerous macrophages were observed in the necrotic areas [75]. In
522 our study, enalapril did not show any visible cardiac tissue damage possibly through its ability to
523 prevent cell infiltration thus preventing apoptosis and necrosis.**

524 The immunohistochemical analysis showed that there were high expressions of cardiac troponin
525 and CRP in ISO group but lower expression of these proteins in ENA and control groups
526 (Figures 14 and 15). In the case of IL-10B, there was low expression of this protein in ISO group
527 but higher expression in ENA and control groups (Figure 16). **Cardiac troponins are regulatory
528 proteins within the myocardium that are released into the circulation when damage to the
529 myocyte has occurred. Therefore, serum troponin is an exquisitely sensitive marker of
530 myocardial injury and is necessary for establishing the diagnosis of MI [76, 77, 78].** This study
531 has shown that ISO caused myocardial injury with upregulation of this biomarker. On the other
532 hand, the down regulation of cardiac troponin by ENA also showed that this drug has ability to
533 protect against myocardial injury in rats.

534 C-reactive protein (CRP) has the capacity to precipitate the somatic C-polysaccharide of
535 *Streptococcus pneumoniae*. It was the first acute-phase protein to be described and is an

536 exquisitely sensitive systemic marker of inflammation and tissue damage [79]. It is a known fact
537 that tissue necrosis is a potent acute-phase stimulus. In myocardial infarction, there is a major
538 CRP response with the magnitude of this response indicating the extent of myocardial necrosis
539 [80]. In all acute myocardial infarcts, CRP is co-deposited with activated complement [81, 82],
540 and research findings have shown that the CRP response did not only reflects tissue damage in
541 this context but also may actually contribute significantly to the severity of ischemic myocardial
542 injury [83]. The lowering of the level of CRP in this study by ENA is a pointer to its ability to
543 halt cardiovascular disease hence cardioprotective effect through its anti-oxidant and anti-
544 inflammatory properties.

545 **Immunohistochemistry in this study further showed that ENA caused increased level of IL-10B.**
546 **IL-10B is a Th₂-type cytokine that is produced by a wide range of immunological cell types,**
547 **including monocytes/macrophages, and it is a potent inhibitor of the proinflammatory cytokines**
548 **and chemokines [84]. Studies have shown that endogenous IL-10 limits angiotensin II (ANG II)-**
549 **mediated oxidative stress, inflammation and vascular dysfunction both *in vivo* and *in vitro*,**
550 **indicating a protective action of IL-10 in vascular diseases such as arterial hypertension [85]. As**
551 **a matter of fact, IL-10 attenuates the increases in vascular superoxide and endothelial**
552 **dysfunction during diabetes and atherosclerosis [86, 87].** In the same way, it could be suggested
553 that IL-10 might be a mediator of cardiac protection against arterial hypertension. It thus shows
554 that the cardioprotective effect of enalapril may also be linked to its anti-inflammatory property
555 as shown by the up regulation of IL-10.

556 **Conclusion**

557 In conclusion, this study has shown that enalapril, an ACE inhibitor has cardioprotective
558 properties, which it exhibited through its anti-oxidant, anti-inflammatory and anti-apoptotic
559 effects. Its antihypertensive property is also exhibited through its nitric oxide increasing ability
560 leading to vasodilation and hence decreases in peripheral resistance.

561 **Ethical Disclaimer:**

562 As per international standard or university standard ethical approval has been collected and
563 preserved by the author(s).

564

565 Conflicts of interest

566 We have no conflict of interest to declare

567

568 **Acknowledgment**

569 This study was supported with a grant (TETFUND/DESS/NRF/UIIBADAN/STI/VOL.
570 1/B2.20.11) received from the National Research Foundation of the Tertiary Education Trust
571 Fund (TETFUND), Nigeria.

572 **References**

- 573 1. Murray CJ and Lopez AD. Global Burden of Disease and Injury Series, Global Health
574 Statistics. Boston: Harvard School of Public Health 1996; *Vols. I and II*. .
- 575 2. Gatica D, Chiong M, Lavandero S and Klionsky DJ. Molecular mechanisms of
576 autophagy in the cardiovascular system. *Circulation Research* 2015; 116(3), 456-467.
- 577 3. Hina SK, Rehman ZH, Dogar N, Jahan M, Hameed ZI, Khan K, Ahmad K, Mukhtar and
578 Valeem EE. Cardioprotective effect of gemmotherapeutically treated *Withania somnifera*
579 against chemically induced myocardial injury. *Pak J Bot* 2010; 42: 1487-1499.
580
- 581 4. Boudina S, Laclau MN and Tariosse L. “Alteration of mitochondrial function in a model
582 of chronic ischemia in vivo in rat heart,” *American Journal of Physiology—Heart and*
583 *Circulatory Physiology* 2002, vol. 282, no. 3, pp. H821–H831.
584
- 585 5. Mohanty DS, Arya A, Dinda KK, Talwar S, Joshi and Gupta SK. “Mechanisms of
586 cardioprotective effect of *Withania somnifera* in experimentally induced myocardial
587 infarction,” *Basic and Clinical Pharmacology & Toxicology* 2004; vol. 94, no. 4, pp.
588 184–190.
589

- 590 6. Sabeena Farvin KH, Anandan R, Kumar SHS, Shiny KS, Sankar TV and Thankappan
591 TK. "Effect of squalene on tissue defense system in isoproterenol-induced myocardial
592 infarction in rats," *Pharmacological Research* 2004, vol. 50, no. 3, pp. 231–236.
593
- 594 7. Abel ED. Glucose transport in the heart. *Front Biosci* 2004; 9: 201–215.
595
- 596 8. Stanley WC, Recchia FA and Lopaschuk GD. Myocardial substrate metabolism in the
597 normal and failing heart. *Physiol Rev* 2005; 85: 1093–1129.
598
- 599 9. Nagoshi T, Yoshimura M, Giuseppe M, Rosano C, Gary D, Lopaschuk C and Mochizuki
600 S. Optimization of cardiac metabolism in heart failure. *Current Pharmaceutical Design*
601 2011; 17: 3846-3853.
602
- 603 10. Wexler BC. Myocardial infarction in young versus old male rats: pathophysiological
604 changes. *Am Heart J* 1987; 96: 70–80
- 605
- 606 11. Khalil MI, Ahmmed I, Ahmed R, Tanvir EM, Afroz R, Paul S, Gan SH and Alam N.
607 Amelioration of isoproterenol-induced oxidative damage in rat myocardium by *Withania*
608 *somnifera* leaf extract. *BioMed Research International Article* 2015. ID 624159, 10 pages
609
- 610 12. Rathore N, John S, Kale M and Bhatnagar D. Lipid peroxidation and antioxidant
611 enzymes in isoproterenol induced oxidative stress in rat tissues. *Pharmacol Res* 1998;38,
612 297-303
- 613 13. Banerjee SK, Sood S, Dinda AK, Das TK and Maulik SK. Chronic oral administration of
614 raw garlic protects against isoproterenol-induced myocardial necrosis in rat. *Comp*
615 *Biochem Physiol* 2003 Part C 136, 377-386.
616
- 617 14. Nirmala C and Puvanakrishnan R. Protective role of curcumin against isoproterenol
618 induced myocardial infarction in rats. *Mol Cell Biochem* 1996; 159, 85-93.
- 619 15. Jackson, Edwin K. "Chapter 30. Renin and Angiotensin". In Brunton, Laurence L.; Lazo,
620 John S.; Parker, Keith. Goodman & Gilman's The Pharmacological Basis of Therapeutics
621 (11th ed.) 2006. New York: McGraw-Hill
- 622 16. Wang W, McKinnie SM, Farhan M, Paul M, McDonald T, McLean B, Llorens-Cortes C,
623 Hazra S, Murray AG, Vederas JC and Oudit GY. "Angiotensin Converting Enzyme 2
624 Metabolizes and Partially Inactivates Pylapelin-13 and Apelin-17: Physiological Effects
625 in the Cardiovascular System". *Hypertension* 2016; 68(2): 365-377.

- 626 17. Gornal AG, Bardawill JC, David MM. Determination of serum proteins by means of
627 Biuret reaction. *J Biol Chem* 1949; 177:751e66.
628
- 629 18. Buetler E, Duron O, Kelly BM. Improved method for the determination of blood
630 glutathione. *J Lab Clin Med* 1963; 61:882e8.
631
- 632 19. Rotruck JT, Pope AL, Ganther HE, Swanson AB, Hafeman DG and Hoekstra WG.
633 Seleniu biochemical role as a component of glutathione peroxidase. *Sci* 1973;
634 179:588e90.
635
- 636 20. Habig WH, Pabst MJ and Jacoby WB. Glutathione-S-transferase activity: the enzymic
637 step in mercapturic acid formation. *J Biol Chem* 1974; 249:130e9.
638
- 639 21. Misra HP and Fridovich I. The role of superoxide anion in the autoxidation of
640 epinephrine and a simple assay for superoxide dismutase. *J Biol Chem.* 1972; 25:
641 247(10): 3170–3175.
642
- 643 22. Varshney R and Kale RK. Effect of calmodulin antagonists on radiation induced lipid
644 peroxidation in microsomes. *Intern J Biol* 1990; 158:733e41.
645
- 646 23. Wolff SF. Ferrous ion oxidation in the presence of ferric ion indicator xylenol orange for
647 measurement of hydrogen peroxides. *Methods Enzymol* 1994; 233: 182e9.
648
- 649 24. Ellman GL. Tissue sulfhydryl groups. *Arch. Biochem. Biophys* 1959; 82 (1): 70-77.
- 650 25. Olaleye SB, Adaramoye OA, Erigbali PP. Lead exposure increases oxidative stress in the
651 gastric mucosa of HCl/ethanol-exposed rats. *World J Gastroent.* 2007; 13: 5121-5126.
- 652 26. Drury R, Wallington E and Cancerson R. Carleton's histological technique". 4th ed. 1976.
653 Oxford University Press, London.
654
- 655 27. Bono DP and Boon NA. Diseases of cardiovascular system. In Davidson's Principles and
656 Practice of Medicine. Edited by Edwards CRW, Boucheir IA. Hong Kong: Churchill
657 Livingstone 1992; pp 249–340.
658
- 659 28. Nian M, Lee P, Khaper N and Liu P. Inflammatory cytokines and postmyocardial
660 infarction remodeling. *Circ Res* 2004; 94: 1543-1553.
661
- 662 29. Wang J, Bo H, Meng X, Wu Y, Bao Y and Li Y. A simple and fast experimental model
663 of myocardial infarction in the mouse. *Tex Heart Inst J* 2006; 33: 290-293.
664

- 665 30. Kerth PA and Vanhoutte PM. Effects of perindoprilat on endothelium-dependent
666 relaxations and contractions in isolated blood vessels. *Am J Hypertens* 1991; 4:226S-
667 234S.
- 668
- 669 31. Wagner JA, Hu K, Bauersachs J, Karcher J, Wiesler M, Goparaju SK, Kunos G and Ertl
670 G. Endogenous cannabinoids mediate hypotension after experimental myocardial
671 infarction. *J Am Col Cardiol* 2001; 38 (7): 2048-2054.
672
- 673 32. Ohman EM, Nanas J, Stomel RJ, Leesar MA, Nielsen DW, O’Dea D, Rogers FJ, Harber
674 D, Hudson MP, Fraulo E and Shaw LK. Thrombolysis and counterpulsation to improve
675 survival in myocardial infarction complicated by hypotension and suspected cardiogenic
676 shock or heart failure: results of the TACTICS Trial. *J Thrombos Thrombol* 2005; 19(1):
677 33-39.
678
- 679 33. Hennekens CH, Albert CM, Godfried SL, Gaziano JM, Buring JE. Adjunctive drug
680 therapy of acute myocardial infarction: evidence from clinical trials. *N Engl J Med* 1996;
681 335: 1660–1667
- 682 34. ACEIMICG. Indications for ACE inhibitors in the early treatment of acute myocardial
683 infarction: systematic overview of individual data from 100,000 patients in randomized
684 trials. ACE Inhibitor Myocardial Infarction Collaborative Group. *Jun* 1198; 97(22): 2202-
685 2212
- 686 35. Lubarsky L and Coplan NL. Angiotensin-Converting Enzyme Inhibitors in Acute
687 Myocardial Infarction: A Clinical Approach. *Preventive cardiology* 2007; 10(3): 156-
688 159.
689
- 690 36. Rona G, Chappel CI, Balazs T and Gaudry R. An infarct-like myocardial lesion and
691 other toxic manifestations produced by isoproterenol in the rat. *Arch Pathol* 1959; 76:
692 443-445.
693
- 694 37. Adameova A, Abdellatif Y and Dhalla NS. Role of excessive amounts of circulating
695 catecholamines and glucocorticoids in stress-induced heart disease. *Can J Physiol*
696 *Pharmacol* 2009; 87: 493-514.
697
- 698 38. Upaganlawar A and Balaraman R. Cardioprotective effects of *Lagenaria siceraria* fruit
699 juice on isoproterenol-induced myocardial infarction in Wistar rats: A biochemical and
700 histoarchitecture study. *J Young Pharmacists* 2011; 3: 297-303.
701

- 702 39. Owens P and O'Brien E. Hypotension in patients with coronary disease: can profound
703 hypotensive events cause myocardial ischaemic events? *Heart* 1999; 82: 477–481.
704
- 705 40. Macfarlane PS, Reid R and Callander R. Pathology Illustrated. Int Student 5th Ed.
706 Churchill Livingstone, London 2000; pp 179-188.
- 707
- 708 41. Rosselli M, Keller PJ and Dubey RK. Role of nitric oxide in the biology, physiology and
709 pathophysiology of reproduction. *Human Reproduction Update* 1998; 4 (1): 3–24.
710
- 711 42. Vallance P, Leone A, Calver A, Collier J and Moncada S. Accumulation of an
712 endogenous inhibitor of nitric oxide synthesis in chronic renal failure. *Lancet* 1992; 339:
713 572–575.
714
- 715 43. Reyes AA, Karl IE and Klahr S. Role of arginine in health and in renal disease. *Am J*
716 *Physiol* 1994; 267: F331–F346.
717
- 718 44. Morris SM Jr. Regulation of enzymes of urea and arginine synthesis. *Annu Rev Nutr*
719 1992; 12: 81–101.
720
- 721 45. Vaziri ND, Ovelisi F and Ding Y. Role of increased oxygen free radical activity in the
722 pathogenesis of uremic hypertension. *Kidney Int* 1998; 53: 1748–1754.
723
- 724 46. Zatz R and Baylis C. Chronic nitric oxide inhibition model six years on. *Rev Hypertens*
725 1998; 32: 958–964.
726
- 727 47. Cachofeiro V, Sakakibara T and Nasjletti A. Kinins, nitric oxide, and the hypotensive
728 effect of captopril and ramiprilat in hypertension. *Hypertension* 1992; 19:138-145
729
- 730 48. Bonner G, Preis S, Schunk U, Toussaint C and Kaufmann W. Hemodynamic effects of
731 bradykinin on systemic and pulmonary circulation in healthy and hypertensive humans. *J*
732 *Cardiovasc Pharmacol* 1990; 15(suppl 6):S46-S56.
733
- 734 49. Touyz RM: Oxidative stress and vascular damage in hypertension.” *Current hypertension*
735 *reports* 2000; 2 (1): 98–105.
736

- 737 50. Wilcox CS. Reactive oxygen species: roles in blood pressure and kidney function,”
738 *Current Hypertension Reports* 2002; 4 (2): 160–166.
739
- 740 51. Sabban EL and Kvetnansky R. Stress-triggered activation of gene expression in
741 catecholaminergic systems: dynamics of transcriptional events. *Trends Neurosci* 2001;
742 24(2): 91–98.
743
- 744 52. Esch T, Sefano GB and Fricchione GL, Benson H. Stress in cardiovascular diseases.
745 *Med. Sci. Monit* 2002; 8: RA93–RA101.
746
- 747 53. Agrawal A and Sharma B. Pesticides induced oxidative stress in mammalian systems. *Int*
748 *J Biol Med Res* 2010; 1(3): 90-104.
749
- 750 54. Maddock C and Pariante CM. How does stress affect you? An overview of stress,
751 immunity, depression, and disease. *Epidemiol. Psychiatr. Soc.* 2001; 10(3): 153-62.
752
- 753 55. Furtmuller, P.G., Arnhold, J., Jantschko, W., Pichler, H and Obinger, C. Redox properties
754 of the couples compound I/compound II and compound II/native enzyme of human
755 myeloperoxidase. *Biochem Biophys Res Commun* 2003; 301: 551–557.
756
- 757 56. Arnhold J, Monzani E, Furtmüller PG., Zederbauer M., Casella L and Obinger C.
758 Kinetics and thermodynamics of halide and nitrite oxidation by mammalian heme
759 peroxidases. *Eur J Inorg Chem* 2006; 19: 3801–3811.
760
- 761 57. Zederbauer M, Furtmüller PG, Brogioni, S, Jakopitsch, C, Smulevich, G, and Obinger, C.
762 Heme to protein linkages in mammalian peroxidases: impact on spectroscopic, redox, and
763 catalytic properties. *Nat Prod Rep* 2007; 24: 571–584.
764
- 765 58. Lacy F, Kailasam MT, O’Connor DT, Schmid-Scho’nbein GW and Parmer RJ. Plasma
766 hydrogen peroxide production in human essential hypertension: role of heredity, gender,
767 and ethnicity. *Hypertension* 2000; 36: 878–884.
768
- 769 59. Paravicini TM and Touyz RM. Redox signalling in hypertension. *Cardiovasc Res* 2006;
770 71:247–258.
771
- 772 60. Alvarez Y, Pe´rez-Giro´n JV, Hernanz R, Briones AM, Garcıa-Redondo A, Beltra´n A,
773 Alonso MJ, and Salaiques M. Losartan reduces the increased participation of

- 774 cyclooxygenase-2-derived products in vascular responses of hypertensive rats. *J*
775 *Pharmacol Exp Ther* 2007; 321:381–388.
776
- 777 61. Rodríguez-Martínez MA, García-Cohen EC, Baena AB, González R, Saláices M and
778 Marín J. Contractile responses elicited by hydrogen peroxide in aorta from normotensive
779 and hypertensive rats: endothelial modulation and mechanism involved. *Br J Pharmacol*
780 1998; 125:1329–1335.
781
- 782 62. Gao YJ and Lee RM. Hydrogen peroxide induces a greater contraction in mesenteric
783 arteries of spontaneously hypertensive rats through thromboxane A (2) production. *Br J*
784 *Pharmacol* 2001; 134:1639–1646.
785
- 786 63. De Cavanagh EMV, Inserra F, Toblli J, Stella I, Fraga CG and Ferder L. Enalapril
787 attenuates oxidative stress in diabetic rats. *Hypertension* 2001; 38: 1130–1136.
788
- 789 64. Chandran G, Sirajudeen KNS, Yusoff NSN, Swamy M and Samarendra MS. Effect of
790 the antihypertensive drug enalapril on oxidative stress markers and antioxidant enzymes
791 in kidney of spontaneously hypertensive rat. *Oxidative Medicine and Cellular Longevity*.
792 2014; Article ID 608512, 10 pages.
793
- 794 65. Packer L. “*Biothiols*” *Methods in Enzymology*; Academic Press Inc.: London, England,
795 1995; volumes 251, Part A & 252 Part B.
796
- 797 66. Arrigo AP. Gene expression and the thiol redox state. *Free Rad. Bio. Med.* 1999, 27,
798 936–944.
799
- 800 67. Bindoli A, Fukato JM and Forman HJ. Thiol Chemistry in peroxidase catalysis and redox
801 signalling. *Antioxid. Redox Sign.* 2008, 10, 1549–1564.
802
- 803 68. Reed DJ. Oxidative stress and mitochondrial permeability transition. in *Biothiols in*
804 *Health and Disease*, eds Packer L., Cadenas E. (Dekker, New York) 1995; pp 231–263.
805
- 806 69. Halliwell B and Gutteridge JM. Protection against oxidants in biological systems: the
807 superoxide theory of oxygen toxicity. *Free Radicals in Biology and Medicine* (Clarendon,
808 Oxford, UK), 2nd ed 1989; pp 87–187.

809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845

70. Forman HJ, Liu R.-M, Shi MM, Packer L and Cadenas E. Glutathione synthesis in oxidative stress. *Biothiols in Health and Disease* (Dekker, New York) eds 1995, pp 189–212.

71. Wexler BC and Greenberg BP. Protective effect of clofibrate on isoproterenol-induced myocardial infarction in arterio-sclerotic and nonarterio-sclerotic rats. *Atherosclerosis* 1978; 29, 373–75.

72. Balazs T, Hanig JP, and Herman EH. Toxic responses of the cardiovascular system. In Casarett and Doull's *Toxicology: the Basic Science of Poins* (C. D. Klaassen, M. O. Amdur, and J. Doull, eds), Macmillan Publishing Company, New York, USA. third edition 2006; pp 387–411.

73. Van Vleet JF, Ferrans, JV, and Herman E. Cardiovascular and skeletal muscle system. In *Handbook of Toxicologic Pathology* (W. M. Haschek, C. G. Rousseaux, and M. A. Wallig, eds), Academic Press, San Diego, CA, USA 2002; Vol. 2, pp 363–455..

74. Goldspink DF, Burniston JG, Ellison GM, Clark WA and Tan L.-B. Catecholamine-induced apoptosis and necrosis in cardiac and skeletal myocytes of the rat in vivo: the same or separate death pathways? *Exp Physiol* 2004; 89, 407–16.

75. Zhang J, Knapton A, Lipshultz SE, Weaver JL, and Herman EH. Isoproterenol-induced cardiotoxicity in Sprague-Dawley rats: correlation of reversible and irreversible myocardial injury with release of cardiac troponin T and roles of iNOS in myocardial injury. *Toxicologic Pathology* 2008; 36:277-288.

76. Gerhardt W, Nordin G and Ljungdahl L. Can troponin T replace CK MBmass as “gold standard” for acute myocardial infarction (“AMI”)? *Scand J Clin Lab Invest Suppl.* 1999; 230: 83–89.

77. Morrow DA, Cannon CP, Jesse RL, Newby K, Ravkilde J, Storrow AB, Wu AHB and Christenson RH. National Academy of Clinical Biochemistry Laboratory Medicine Practice Guidelines: clinical characteristics and utilization of biochemical markers in acute coronary syndromes. *Clin Chem.* 2007; 53(4):552–574.

- 846 78. Jaffe AS, Ravkilde J, Roberts R, Naslund U, Apple FS, Galvani M and Katus H. It's time
847 for a change to a troponin standard. *Circulation* 2002; 102: 1216-1220.
848
- 849 79. Pepys MB and Baltz ML. Acute phase proteins with special reference to C-reactive
850 protein and related proteins (pentaxins) and serum amyloid A protein. *Adv. Immunol.*
851 1983; 34: 141–212.
- 852 80. de Beer FC, Hind CR, Fox KM, Allan RM, Maseri A, and Pepys MB. Measurement of
853 serum C-reactive protein concentration in myocardial ischaemia and infarction. *Br. Heart*
854 *J.* 1982; 47: 239–243.
855
- 856 81. Kushner I, Rakita L and Kaplan MH. Studies of acute phase protein. II. Localization of
857 Cx-reactive protein in heart in induced myocardial infarction in rabbits. *J. Clin. Invest.*
858 1963; 42: 286–292.
859
- 860 82. Lagrand WK, Niessen HW, Wolbink GJ, Jaspars LH, Visser CA, Verheugt FW, Meijer
861 CJ and Hack CE. C-reactive protein colocalizes with complement in human hearts during
862 acute myocardial infarction. *Circulation.* 1997;95: 97–103.
863
- 864 83. Griselli M, Herbert J, Hutchinson WL, Taylor KM, Sohail M, Krausz T and Pepys MB.
865 C-reactive protein and complement are important mediators of tissue damage in acute
866 myocardial infarction. *J. Exp. Med.* 1999; 190: 1733–1739.
867
- 868 84. Akdis CA and Blaser K (2001): Mechanisms of interleukin-10-mediated immune
869 suppression. *Immunology* 2001; 103:131–136.
870
- 871 85. Didion SP, Kinzenbaw DA, Schrader LI, Chu Y, Faraci FM: Endogenous interleukin-10
872 inhibits angiotensin II-induced vascular dysfunction. *Hypertension* 2009; 54: 619-624.
873
- 874 86. Mallat Z, Besnard S, Duriez M, Deleuze V, Emmanuel F, Bureau MF, Soubrier F,
875 Esposito B, Duez H, Fievet C, Staels B, Duverger N, Scherman D and Tedgui A.
876 Protective role of interleukin-10 in atherosclerosis. *Circ Res* 1999; 85: e17-e24.
877
- 878 87. Gunnett CA, Heistad DD and Faraci FM. Interleukin-10 protects nitric oxide-dependent
879 relaxation during diabetes: role of superoxide. *Diabetes* 2002; 51: 1931-1937.

880

881