

## Short communication

# Total Phenols, Antioxidant Capacity and Antibacterial Activity of Manuka Honey Extract

## ABSTRACT

**Aims:** To evaluate total phenols content (TPC), antioxidant capacity (TAC) and antibacterial activity of Manuka honey extract (MHE) and to compare such properties with those for unfractionated Manuka honey.

**Study design:** *In vitro* study

**Place and Duration of Study:** School of Biomedical Sciences, Ulster University, Coleraine, UK. Between September 2016 and September 2017.

**Methodology:** MHE was prepared by solvent extraction using ethyl acetate. TPC was determined by Folin-Ciocalteu assay. The iron (III) reducing antioxidant capacity (IRAC) method was used to determine TAC. Antibacterial activity was evaluated using disc diffusion assay and 96-well microtiter plate method with absorbance measured at 600 nm.

**Results:** The TPC for MHE was 30-fold higher than the value for Manuka honey ( $33420 \pm 1685$  mg vs.  $1018 \pm 78$  mg GAE/kg) whilst TAC values were ~100-times greater ( $83,198 \pm 7064$  vs.  $793 \pm 104$  TEAC, respectively). Antibacterial activity assessed by disc diffusion for Manuka honey (18.5mm on *S. aureus* and 20mm on *E. coli*) was two times greater than for MHE (9mm for both *S. aureus* and *E. coli*). The 96-well microtiter plate assay confirmed the greater antibacterial activity for Manuka honey compared to equal concentrations MHE.

**Conclusion:** A polyphenol-rich Manuka honey extract with a high total antioxidant capacity, showed little or no antibacterial activity against *E. coli* and *S. aureus* in contrast with unfractionated Manuka honey.

**Keywords:** Manuka honey, Manuka extract, total phenols, antioxidant, antibacterial,

## 1. INTRODUCTION

Infectious diseases continue to pose a threat to human health worldwide. Wound infections are commonly caused by bacterial pathogens [1-3]. Owing to the rising occurrence of antibiotic-resistant bacterial strains, alternative ancient remedies and plant-based products such as honey are being evaluated for therapeutic use. The medicinal importance of honey has been widely documented in the world's medical literature; standardized active Manuka honey has been registered as a wound care product with appropriate medical regulatory bodies [1-3]. Sherlock et al [4] demonstrated antibacterial activity for Chilean Honey (Ulmo 90 honey) and New Zealand Manuka honey (UMF® 25+) against 5 MRSA strains. Ahmed and Othman [5] found that Tualang honey and Manuka honey could inhibit growth of Gram-positive MRSA strains including *S. aureus* and *S. pyogenes* and Gram-negative strains like *P. aeruginosa*, *E. coli* and *Enterobacter cloacae* [5].

Kwakman and Zaat [6] reported the antibacterial activity for Revamil™ honey and Manuka honey, the two leading medicinal grade honeys, arose from different mechanisms involving, hydrogen peroxide, bee defensin, methylglyoxal, and unidentified components. Several studies proposed that phenolic compounds might contribute to the non-peroxide antibacterial activity of Manuka honey [1, 6]. However, the specific contribution of phenolic components to the antibacterial action of Manuka honey has not been well investigated. The general aims of this project were to evaluate, the total phenols content (TPC), total antioxidant capacity (TAC), and antibacterial activity of an ethyl acetate extract from Manuka honey (Manuka Honey Extract; MHE) and to compare these characteristics with unfractionated Manuka honey.

## 33 2. MATERIAL AND METHODS

34

### 35 2.1 Materials

36 Manuka honey rated Unique Manuka Factor (UMF) 10+, 15+ and 18+ were purchased from Comvita Ltd  
37 (UK). Ethyl acetate, Folin & Ciocalteu reagent, sodium carbonate, gallic acid, Trizma base, ferrozine (3-  
38 (2-Pyridyl)-5,6-diphenyl-1,2,4-triazine-4',4''-disulfonic acid sodium salt), ammonium iron (III) sulfate  
39 dodecahydrate, methanol and other chemicals were purchased from Sigma-Aldrich Ltd (Gillingham, UK).  
40 Nutrient broth (Oxoid), nutrient agar (Oxoid), and penicillin-streptomycin mixture (Pen-strep) were  
41 purchased from ThermoFisher Scientific (UK). Bacteria strains (*Staphylococcus aureus*, *Escherichia coli*)  
42 were obtained from School of Biomedical Sciences, Ulster University (UK).

43

### 44 2.2 Preparation of Manuka honey extract (MHE)

45 Honey extract was prepared using ethyl acetate as solvent as described by Tan et al [7] with modification.  
46 Manuka honey (UMF 10+, 20g) was dispersed in 80ml of distilled-deionized water and 100ml ethyl  
47 acetate. The mixture was stirred using a magnetic stirrer for 24 hours. The emulsion formed was  
48 transferred to glass centrifuge tubes and centrifuged at 2,000 RPM for 15 minutes. The non-aqueous  
49 ethyl acetate phase was air-dried and the residue formed was re-dissolved in methanol solvent, filtered  
50 through 0.2  $\mu\text{m}$ , and then stored at  $-18^{\circ}\text{C}$  until used. The solids content of MHE extract was determined  
51 by drying 50  $\mu\text{l}$  of MHE and weighing the residue.

52

### 53 2.3 Determination of total phenols content

54 The total phenols content was determined using Folin-Ciocalteu method adapted for microplate analysis  
55 as described previously [8, 9].

56

### 57 2.4 Determination of antioxidant capacity

58 Antioxidant capacity was determined using the iron (III) reducing antioxidant capacity (IRAC) as described  
59 recently [8]. The IRAC reagent was prepared by dissolving 20 mg ferrozine dye in 9ml of Tris buffer and  
60 adding ferric (III) ammonium sulphate (4mg in 1ml water). For TAC determinations, samples of honey  
61 (20ul) were mixed with 280  $\mu\text{l}$  of ferrozine solution and incubating for 30 min at  $37^{\circ}\text{C}$ . TAC assay were  
62 calibrated using trolox (0-1000  $\mu\text{M}$ ) as antioxidant standard. A microplate reader (VersaMax, Molecular  
63 Devices, Sunnyvale, California, USA) was used for absorbance measurements at 562nm.

64

### 65 2.5 Antibacterial activity screening

#### 66 2.5.1. Antibacterial screening by disc diffusion assay

67 The disc diffusion assay for antibacterial activity was carried out as described previously with minor  
68 modification [10] using two bacteria strains, one Gram-positive bacteria- *S. aureus* and one Gram-  
69 negative bacteria- *E. coli*. Working in laminar flow hood, bacterial inoculated broth (200  $\mu\text{l}$ ) was  
70 transferred to blank nutrient agar plates (each bacteria x2), and allowed to dry. Thereafter, 6 blank paper  
71 discs were transferred onto agar plates using tweezer and ensuring equal spacing between each.  
72 Samples (20  $\mu\text{l}$ ) of Manuka honey (UMF10+ Manuka honey extract, 25% UMF10+, 15+, 18+ Manuka  
73 honey), MHE and controls (Pen-strep,) were slowly added to the blank disc and were left to dry briefly.  
74 Plates were incubated upside down overnight at  $37^{\circ}\text{C}$ . The diameter of zones of inhibitions was  
75 measured after 24 hours in mm.

76

#### 77 2.5.2 Micro-plate assay for antibacterial activity

78 Samples of a 24-h grown liquid culture (50 $\mu\text{L}$ ) were transferred to 96-well microtiter plate (x3), and 50 $\mu\text{L}$   
79 of sterile honey sample (25% w/v in water), MHE or antibiotic (Pen-strep) was added. The 96-well  
80 microtiter plates were then incubated at  $37^{\circ}\text{C}$  for 24h with gentle shaking and absorbances were read at  
81 600nm. The antibacterial effect (%) was determined from the expression,  $100 \cdot (1 - (A_H/A_0))$  where  $A_H$  and  
82  $A_0$  are absorbance readings with and without honey treatment.

83

### 84 2.6 Data analysis

85 Data analysis was performed using Microsoft excel and IBM SPSS Statistics Version 24. Correlation was  
 86 obtained by Pearson correlation and significance was assessed in two-tailed at level of significance of  $P=$   
 87 0.01.

88  
 89

90 **3. RESULTS AND DISCUSSION**

91

92 **3.1 Total phenols content and antioxidant capacity of Manuka honey extract**

93 The total phenols content (TPC) for MHE and unfractionated honey were expressed in mg-Gallic acid  
 94 equivalent (GAE) per kg and are shown in Table 1. The TPC for Manuka honey was  $1018.32 \pm 78.84$  mg  
 95 GAE/kg honey (Table 1) compared with values in the range of 430 - 2706 mg GAE/kg reported previously  
 96 [11]. The TPC for honey is known to vary with various factors, including monoflorality of honey, age and  
 97 geographic origins of honey samples [12]. Compared to the original honey the, MHE had a 32.8-fold  
 98 increase TPC. The TAC for honey evaluated in terms of IRAC is shown in Table 1 for ethyl acetate honey  
 99 extract, and original honey (Table 1). The TAC for MHE was 105-fold increase compared to  
 100 unfractionated Manuka honey.

101 Moniruzzaman et al [12] and Alvarez-Suarez et al [13] supported c phenols play an important role in the  
 102 antioxidant capacity of honey. The dominating phenolic components that have been identified in Manuka  
 103 honey were phenyllactic acid and a group of methoxylated benzoic acids [7, 9]. Other literature also  
 104 reported that components in Manuka honey that are responsible for antioxidant effect of honey include  
 105 flavonoids such as chrysin, quercetin, isorhamnetin and luteolin, phenolic acid like gallic acid, caffeic acid  
 106 and syringinic acid [13]

107 **Table 1. Total phenols content and total antioxidant capacity of Manuka honey (UMF10+) and the**  
 108 **Manuka honey Extract**

109

Sample	TPC (mg GAE/kg)	TAC(mg TE/kg)
Honey (UMF10+)	$1018 \pm 79$	$793 \pm 104.4$
MHE	$33420 \pm 1685$	$83198 \pm 7064$

110 \*The total phenols content (TPC) and total antioxidant capacity (TAC) for Manuka honey (UMF10+) and ethyl acetate  
 111 extract from Manuka honey (MHE) as mg Gallic acid Equivalent (GAE)/ kg or mg Trolox Equivalent (TE)/ kg sample.

112  
 113

114 **3.2 Antibacterial activity**

115 Table 2 shows the antibacterial activity using the disc diffusion assay for *S. aureus* and *E. coli*. The MHE  
 116 was adjusted to deliver similar quantities GAE per paper disc as was used for Manuka honey. For  
 117 example, paper discs were loaded with 20- $\mu$ l Manuka honey UMF10+ (250g/l) and hence the GAE  
 118 loading per disc was  $(250\text{g/l} \cdot 20 \times 10^{-6}\text{l} \cdot 1.108 \times 10^{-3}\text{g GAE/g}) = 5.1 \times 10^{-6}\text{gGAE}$ . After pre-diluting MHE by  
 119 33-fold, the GAE loading per disc was  $(7.5\text{g/l} \cdot 2 \times 10^{-6}\text{l} \cdot 33.420 \times 10^{-3}\text{g GAE/g}) = 5.0 \times 10^{-6}\text{gGAE}$ . The results  
 120 show clearly that the honey had antibacterial activity whilst MHE showed little or no antibacterial activity.  
 121 (Fig. 1).

122

123 **Table 2. Measurement of zone of inhibition (mm) by disc diffusion assay**


Bacteria	Zone of inhibition (mm)					
	Controls		Honey Samples			
	Pen-Step	Methanol	UMF10+	UMF15+	UMF18+	MHE
<i>S. aureus</i>	27.0	9.0	18.5	18.5	20.5	9.0
<i>E. coli</i>	26.0	9.5	20.0	21.0	22.5	9.0

124 \*MH= Manuka honey, Tests involve 20  $\mu$ l of 25% (w/v) added to paper discs. MHE = Manuka honey extract from  
 125 UMF10+ honey.

126

127 Antibacterial activity testing using microplate/ spectrophotometric method also showed that the honey  
 128 extract had no antibacterial activity (Figure 1). For these tests, the exposure concentration used for 10+  
 129 Manuka honey and MHE were 0.76 mM and 0.74 mM GAE, respectively. The microorganism used for

130 testing (*S. aureus* and *E. coli*) are known to be sensitive to Manuka honey [5]. Therefore, it was expected  
 131 that screening (by disc diffusion assay and spectroscopic analysis) showed antibacterial activity with  
 132 Manuka honey. Interestingly, MHE showed little or no antibacterial activity when tested at a similar  
 133 concentration as honey. Indeed, *S. aureus* measurements were consistent with increased growth after  
 134 exposure to MHE (Fig. 1).  
 135



136 **Fig. 1. Antibacterial effect for Manuka honey and Manuka hone extract (MHE)**

137 *Tested with 96-well microtiter plats with E. coli or S. aureus. Penicillin-streptomycin was used as +ve*  
 138 *control. Honey samples were 12.5% (rated UMF 10+ -UMF18+). MHE is Manuka Honey Extract See text for*  
 139 *details.*  
 140  
 141

142 Based on current results, the polyphenols from honey may not be a major factor contributing to the  
 143 antibacterial activity of Manuka honey. The findings agree with previous reports, which noted that Manuka  
 144 honey polyphenols (benzoic acids, cinnamic acids and flavonoids) could not account for antibacterial  
 145 activity [14, 15]. Alternatively, the concentration of phenolic compounds from honey might be too low to  
 146 contribute to antibacterial activity [6]. By contrast, there is considerable evidence showing a correlation  
 147 between methylglyoxal content and antibacterial activity of Manuka honey [16] whilst only slight  
 148 antibacterial activity was ascribed to the high sugar content and acidity [6]. Since, honey contains  
 149 comparable levels of polyphenols and methylglyoxal [17], further research is underway to determine if  
 150 these components interact and possible consequences on antibacterial activity.  
 151  
 152

153 **4. CONCLUSION**

154 Manuka honey organic extract, containing a high total phenols content and total antioxidant capacity  
 155 showed little or no antibacterial activity. Further in-depth research is needed to understand the  
 156 composition and characteristics of Manuka honey extracts.  
 157

158 **CONSENT**

159 No consent required

160 **ETHICAL APPROVAL (WHERE EVER APPLICABLE)**

161 No ethical approval required  
 162

163  
 164  
 165  
 166  
 167  
 168  
 169  
 170  
 171  
 172  
 173  
 174  
 175  
 176  
 177  
 178  
 179  
 180  
 181  
 182  
 183  
 184  
 185  
 186  
 187  
 188  
 189  
 190  
 191  
 192  
 193  
 194  
 195  
 196  
 197  
 198  
 199  
 200  
 201  
 202

## REFERENCES

1. McLoone P, Warnock M, Fyfe L. Honey: A realistic antimicrobial for disorders of the skin. *J Microbiol Immunol Infect.* 2016 49(2):161-7.
2. Carter DA, Blair SE, Cokcetin NN, Bouzo D, Brooks P, Therapeutic Manuka honey: no longer so alternative. *Front Microbiol.* 2016 7: 569.
3. Oryan A, Alemzadeh E, Moshiri A. Biological properties and therapeutic activities of honey in wound healing: a narrative review and meta-analysis. *J Tissue Viability.* 2016;25(2):98-118.
4. Sherlock O, Dolan A, Athman R, Power A, Gethin G, Cowman S, Humphreys H. Comparison of the antimicrobial activity of Ulmo honey from Chile and Manuka honey against methicillin-resistant *Staphylococcus aureus*, *Escherichia coli* and *Pseudomonas aeruginosa*. *BMC Compl Alt Med.* 2010;2:10(1):47.
5. Ahmed S, Othman NH. Review of the medicinal effects of Tualang honey and a comparison with Manuka honey. *MJMS.* 2013;20(3):6.-13.
6. Kwakman, PHS, Zaat, SA.J. Antibacterial components of honey. *lubmb Life.*2012 64(1): 48-55.
7. Tan ST, Holland PT, Wilkins AL, Molan PC. Extractives from New Zealand honeys. 1. White clover, Manuka and Kanuka unifloral honeys. *J Agric Food Chem.*1988.36(3): 453-460.
8. Owusu-Apenten R, Mohd Yusof H I, Nigam PS. Cryptic total antioxidant capacity of Manuka honey measured by a new microplate assay for iron (III) reducing antioxidant capacity (IRAC) and other methods. *Food Chem.* (2017): Submitted
9. Portokalakis I, Mohd Yusof HI, Ghanotakis D, Nigam P, Owusu-Apenten R , Manuka Honey-induced cytotoxicity against MCF7 breast cancer cells is correlated to total phenol content and antioxidant power.*JABB.*2016. 8(2): p. 1-10.
10. Barlow R, Barnes D, Campbell A, Nigam PS, Owusu-Apenten R. Antioxidant, anticancer and antimicrobial, effects of *Rubia cordifolia* aqueous root extract. *JABB.* 2015;5(1):6-14.
11. Stephens JM, Schlothauer RC, Morris BD, Yang D, Fearnley L, Greenwood DR, Loomes KM. Phenolic compounds and methylglyoxal in some New Zealand Manuka and Kanuka honeys. *Food Chem.*2010;120(1): 78-86.
12. Moniruzzaman M, Sulaiman SA, Khalil MI, Gan SH. Evaluation of physicochemical and antioxidant properties of sourwood and other Malaysian honeys: a comparison with Manuka honey. *Chem Cent J.* 2013;7(1):138.
13. Alvarez-Suarez JM, Gasparri M, Forbes-Hernández TY, Mazzoni L, Giampieri F. The composition and biological activity of honey: a focus on Manuka honey. *Foods.*2014;3(3):420-432.
14. Weston RJ, Mitchell KR, Allen KL. Antibacterial phenolic components of New Zealand Manuka honey. *Food Chem.* 1999;64(3):295-301.
15. Weston RJ, Brocklebank LK, Lu Y. Identification and quantitative levels of antibacterial components of some New Zealand honeys. *Food Chem.* 2000; 70(4):427-435.

- 203 16. Atrott J, Henle T. Methylglyoxal in Manuka honey—correlation with antibacterial properties. Czech J  
204 Food Sci.2009;27(Spec.):S163-S165.
- 205 17. Kwok TH, Kirkpatrick G, Mohd Yusof HI, Portokalakis I, Nigam P, Owusu-Apenten R. Rapid  
206 colorimetric determination of methylglyoxal equivalents for Manuka honey. JABB. 2016;7(1):1-6.
- 207