

Vegetation Structure and Diversity of Wadi Wasaa, Jazan, Saudi Arabia

Abstract

This study was the first report conducted on Wadi Wasaa of Jazan area in Saudi Arabia. A total of 95 species belonging to 75 genera and 31 families were recorded, both Poaceae and Euphorbiaceae were the dominant families constituted 23% of the total species of the study area. The next dominant families were Apocynaceae and Malvaceae. Chamaephytes and therophytes were the prevailed life forms, indicating a typical desert life-form spectrum (chameo-therophytic) type, followed by phanerophytes. The chorological analysis revealed a total of 26 species representing 27% were falls under monoregional, 56 species (60.0%) as bioregional area and four species were detected under pluriregional region. Cover abundance values were visually estimated and used to form ten clusters of plant community types by statistical methods with Euclidian Distance and Ward method using SPSS program (ver.20). The Shannon-Wiener diversity index was used to estimate diversity, richness and evenness of the recorded species where it revealed the highest diversity index (H) was detected in *Tamarindus indica* community, followed by the community of *Acacia asak*, whereas the lowest one calculated in *Lawsonia inermis*. At the same time, Sorensen's Index of Similarity (ISs) confirmed some different affinities among these communities.

Key words: Floristic composition, Diversity, Chorology, Community type Analysis

Introduction

The diversity of wild plants is an important aspect of our earthly environments and plays a major role in protecting the ecological consistency and balance of the region

26 (Abd El-Khalik *et al.*, 2017). The flora of Saudi Arabia is one of the richest
27 biodiversity in the Arabian Peninsula and comprises very important genetic resources
28 of crops and medicinal plants (Atiqur *et al.*, 2004). In the dry lands, wadis represent
29 one of the most prominent desert landforms, which exhibit physiographic
30 irregularities that lead to parallel variations in plant species distribution (Kassas and
31 Girgis, 1964). Life-form distribution is significantly related to the topography and
32 landform. (Zohary, 1973 and Orshan, 1986). Jazan province is situated in the
33 southwestern part of Saudi Arabia characterized by rocky slopes, cliffs and crevices
34 with granite, sandy soil whereas the hilly areas are generally formed of rocky cliffs,
35 rocky ridges, granite boulders, granite outcrops, granite sand stones and crevices (Al-
36 Farhan *et al.*, 2005). Jazan region can be broadly divided into Tihama, the Escarpments
37 and the Farasan Islands. The first two regions are part of the oldest agricultural centers
38 of the Arabian Peninsula and composed of wadis, mountains and plateaus (Al-Farhan
39 *et al.*, 2005 and Masrahi, 2012) Several studies on the floristic diversity and
40 vegetation analysis in Tihama plains of Saudi Arabia were performed by El-
41 Demerdash *et al.* (1994), Masrahi, (2012) and Marei *et al.* (2014). Wadi vegetation of
42 Saudi Arabia were studied by many authors such as Wadi Al Ammaria (El Ghenam,
43 2006, Al Yemeni 2001), Wadi Al Jufair (Al Atar *et al.*, 2012), Wadi Al Argy (Farrag,
44 2012), Wadi Al Noman (Abdel Khalik *et al.*, 2013), Wadi El Ghayl (Fahmy and
45 Hassan, 2005), Wadi Al Rummah (El Ghazali *et al.*, 2013) Wadi Talha (Al Wadie,
46 2002), Wadi Khulab (Kasem and Marei, 2017) and Wadi Tashar (El-Shabasy and
47 Kasem, 2018). Evenness Index (E) and Shannon-Wiener Diversity Index (H) methods
48 are of the most widely used approaches in measuring the diversity of species (Siraj *et*

49 *al.*, 2016). The present study aimed to investigate the floristic composition, life-form
50 and chorotype to classify, document and assess the species diversity between the
51 different community types of the Wadi Wasaa of Jazan region in Saudi Arabia.

52 **Study Area**

53 Wadi Wasaa located in the southeast of Jazan. It located between the AL-Dabha
54 Mountain in the east and AL-Hague in the west, it around 60 km²; altitude ranges
55 from 480-680m above sea level (a.s.l.). It lies between 17°48' N latitude and 42° 89'
56 E longitude (Figure 1). According to Al-Farhan *et al.* (2005) and Masrahi (2012) the
57 study area lies within the subtropical dry zone and has very hot summers and mild
58 winters; the topography is mostly mountainous with steep to moderately steep slopes
59 gradually tapering off to a relatively flat mountain plateau. December and January are
60 the coldest months (20°C) while the hottest month is, July (39°C). The maximum
61 precipitation (20.0 mm) falls during June, while the minimum precipitation of about
62 6.0 mm falls during November (Figure 2).

63 **Materials and methods**

64 A total of 46 sites located in a randomly stratified manner were selected along the
65 wadi Wasaa and conducted from January to October 2017, this period in Jazan Region
66 represents the optimum growing and flowering seasons for most plant species.
67 Locations and sample plots (25 m × 25 m) were selected randomly using the methods
68 of Muller-Dombois and Ellenberg (1974) and Barbour *et al.* (1987). The collected
69 specimens were identified and named according to Chaudhary (2001), Al-Farhan *et al.*
70 (2005), Migahid (1996) and updated according to the Plant List database (2013).
71 Plant specimens deposited at Jazan University Herbarium, KSA (JAZUH). Life-forms

72 were determined according to Raunkier (1937). A chorological analysis of the recorded
 73 species was made to assign to world geographical groups, according to Wickens
 74 (1978) and Zohary (1973). Altitude and geographical coordinates were measured
 75 using GPS for each quadrat (Geographical Position System). Cover abundance were
 76 calculated by the equation: Total number of individuals of the species/ total number of
 77 quadrats in which species has occurred. Cover abundance were converted to 1-9
 78 according to Braun-Blanquet scale (Mueller-Dombois & Ellenberg 1974) and
 79 modified later by Maarel (1979). The computer program (SPSS, ver.20) was used to
 80 analyze the vegetation structure and perform hierarchical clustering dendrogram
 81 constructed from ten plant community types (Fig. 6), these communities named by the
 82 highest mean cover abundance in each community. The species diversity of each
 83 cluster was calculated using Shannon-Weiner diversity index (1949) based on
 84 cover/abundance value of the species as input source.

85

$$H = - \sum P_i \ln P_i$$

86

87 Where, H: Shannon-Wiener Index and; P_i: proportion of individual species; ln: log
 88 base. The relative equitability (evenness) of the species in each cluster was also
 89 calculated.

90

$$\text{Equitability } J = \frac{H'}{H_{max}} = \sum_{i=1}^S \frac{P_i \ln P_i}{\ln S}$$

91 Where, S: the number of species; P_i: the proportion of individuals of the species or the
 92 abundance of the species expressed as a proportion of total cover; ln: log base.
 93 Sorenson's Similarity ratio was used to evaluate the phytogeographical similarity

94 between the plant community types. It was described using the following formula
95 (Kent and Cooker, 1992)

96

$$Ss = \frac{2a}{2a + b + c}$$

97

98

99 Where Ss: Sorensen's similarity coefficient; a: number of species common to both
100 sites; b: species number in site one; c: species number in site two.

101 **Results**

102 *Floristic analysis*

103 The floristic data on the study area, occurring between altitudinal gradients of 480-680
104 m (a.s.l), indicates a total of 95 species belonging to 75 genera and 31 families.

105 According to species richness, the majority of plants in the study area are perennials
106 (66 species, 69.5% of the total recorded species), the second most frequent growth

107 type was the annuals which revealed by 27 species (28.5% of the total species) also

108 two species of *Chenopodium fasciculosum* and *Asphodelus tenuifolius* were estimated

109 as biennial life span. Four species of *Leptadenia arborea*, *Merremia aturensis*,

110 *Dalechampia scandens* and *Cissus rotundifolia* were estimated as climber species.

111 Poaceae and Euphorbiaceae were the most dominant families represented by 14 and 8

112 species, respectively (Table 1). The next abundant families were Apocynaceae and

113 Malvaceae which represented by 6 species. Acanthaceae, Astraceae, Amaranthaceae,

114 Papilionaceae have five species constituted a total of 21%. Four species were recorded

115 in Mimosaceae while Caesalpiniaceae, Boraginaceae, Solanaceae and Zygophyllaceae

116 were finding out by three species each. Asphodalaceae, Cleomaceae, Lamiaceae,

117 Moraceae, Nyctaginaceae, Plantagonaceae and Salvadoraceae were represented by

118 two species. The remainder (11 families) contributed 12.0% of the total species
119 represented by single species each (Figure 3).

120 ***Life form spectrum***

121 According to the life form classification of Raunkiaer (1937) and as shown in Table 2
122 and Figure 4, the chamaephytes were the most dominant life form, constituted by 33
123 species representing 35% of the total recorded species followed by the therophytes
124 represented by 28 species (29%). On the other hand, 17 species of the phanerophytes
125 estimated 18-% were conducted. Also Hemicryptopyte were occurred by 11 species
126 (12%) of the total recorded taxa. Cryptophytes have the lowest contribution by six
127 species of *Asphodelus tenuifolius*, *Cyperus conglomeratus*, *Corchorus depressus*,
128 *Cenchrus ciliaris*, *Panicum turgidum* and *Sorghum bicolor* with a percentages of 6%.

129 ***Phytogeographical data***

130 Regarding the global floristic regions, monoregional, biregional and pluriregional are
131 constructed as phytochorial regions (Table 2). A total of 26 species representing 27%
132 were falls under monoregional region. In this area the highest number of 15 species
133 was recorded in Saharo-Arabian (16%) whereas the lowest one which estimated by
134 five species of *Abutilon bidentatum*, *Boerhavia elegans*, *Echinochloa colona*, *Opuntia*
135 *dillenii* and *Ricinus communis* recoded in Tropical region. Biregional area included
136 the highest number of species, i.e. 56 species with 60%, 32 species occurred in the
137 area shared by Saharo-Arabian and Sudano-Zambezian regions (34%) followed by
138 area of Saharo-Arabian shared with tropical represented by 14 species (16%). Both
139 Saharo-Arabian and Mediterranean has four species (4%). Both Mediterranean-Irano-
140 Turanian and Mediterranean-Tropical regions are represented by two species (2%).

141 The lowest one was recorded in Saharo-Zambezian and tropical regions has only one
 142 species of *Cyanthillium cinereum*. The pluriregional area (4%) has four species of
 143 *Euphorbia inarticulata*, *Dichanthium foveolatum*, *Ziziphus spina-christi* and
 144 *Corchorus tridens* falls under one main phytochoria of Mediterranean, Saharo-
 145 Arabian and Sudano-Zambezian. The remainder nine recorded taxa were distributed
 146 as follows: three species are cosmopolitan and four species pantropical and only two
 147 species of *Lawsonia inermis* and *Sorghum bicolor* are cultivated plants (Table 2 and
 148 Figure 5).

149 ***Dominant Community Types (DCT)***

150 Distribution of the plant community types among their altitudinal ranges was given in
 151 Table 4. Based on the mean cover abundance values, the description of the ten plant
 152 community types (Table 3, 4 and Figure 7) can be summarized as follow: **I-Ziziphus**
 153 ***spina-christi***, this community type consists of 5 quadrats with 35 species distributed
 154 between altitudinal ranges of 500m and 570m a.s.l., this community types found at
 155 fine calcareous soils in the wadi bed associated with *Adenium obesum*,
 156 *Anisotes trisulcus*, *Argemone ochroleuca*, *Barleria trispinosa*, *Caralluma*
 157 *retrospiciens*, *Heliotropium longiflorum*, *Lavandula coronopifolia*, *Trianthema*
 158 *crystallina* and *Tribulus terrestris*, their abundance cover is 10.4%. **II-Salvadora**
 159 ***persica***, this community type widespread in the wadi terrace, consists of three stands
 160 in which 18 species distributed between altitudinal ranges of 520m and 586m a.s.l.
 161 Associated with *Aloe officinalis*, *Anisotes trisulcus*, *Blepharis edulis*, *Calotropis*
 162 *procera*, *Corchorus depressus*, *Euphorbia triaculeata*, *Indigofera colutea*, *Ocimum*
 163 *forsskaolii*, *Ziziphus spina-christi*, *Pluchea dioscoridis* and *Senra incana*; the

164 abundance plant cover is about 5.7%. **III- *Anisotes trisulcus***, located as finely-
165 calcareous soils on wadi slopes and bed associated it contain four stands with 29
166 species distributed between altitudinal ranges of 450m and 510m a.s.l. This
167 community types associated with *Abutilon hirtum*, *Acacia asak*, *Acacia tortilis*,
168 *Adenium obesum*, *Aerva javanica*, *Cleome scaposa*, *Euphorbia triaculeata*,
169 *Forsskaolea tenacissima*, *Indigofera spinosa*, *Leptadenia arborea*, *Lavandula*
170 *coronopifolia* and *Maytenus senegalensis*; cover abundance is 9.53%. **IV-*Adenium***
171 ***obesum*** community, found at the wadi plateau and fissures it comprised of four sites
172 with 32 species distributed between altitudinal ranges of 550m and 640m a.s.l
173 associated with *Anisotes trisulcus*, *Cenchrus ciliaris*, *Eragrostis papposa*, *Eclipta*
174 *prostrate*, *Panicum turgidum*, and seedling of *Lawsonia inermis* located at sandy-
175 calcareous soils, cover abundance about 8.15%. **V-*Ricinus communis*** community
176 type, found at fine sandy soils it consists of four stands with 27 species distributed
177 between altitudinal ranges of 450m and 510m a.s.l associated with *Acalypha*,
178 *fruticosa*, *Asphodelus tenuifolius*, *Catharanthus roseus*, *Cenchrus ciliaris*,
179 *Chenopodium murale*, *Chloris barbata*, *Cyperus conglomeratus*, *Dobera glabra*,
180 *Pluchea dioscoridis* and *Acacia tortilis* seedlings, the plant cover abundance about
181 9.20%. **VI-*Acacia asak*** community type, occupies a large parts of the wadi, located
182 on slopes at sandy soils, it consists of seven plots with 42 species distributed between
183 altitudinal ranges of 550m and 580m a.s.l, associated with *Abutilon hirtum*, *Aerva*
184 *javanica*, *Anisotes trisulcus*, *Argemone ochroleuca*, *Catharanthus roseu* and *Fagonia*
185 *indica*; their cover abundance are 13.10-%. **VII-*Lawsonia inermis*** community,
186 located at sandy soils, it represents a large amount in the plateau, consists of two

187 quadrates with 15 species distributed between altitudinal ranges of 630m and 690 m
 188 a.s.l associated with *Asphodelus tenuifolius*, *Aristida adscensionis*, *Cleome viscosa*,
 189 *Cyperus conglomerates*, *Malva parviflora*, *Paspalidium desertorum*, and *Senna*
 190 *alexandrina*; cover abundance about 2.71%. **VIII-Dobera glabra** community type
 191 inhabits the wadi bed; it consists of three stands with 22 species distributed between
 192 altitudinal of 460m and 570m a.s.l, associated with *Acacia asak* branches, *A.*
 193 *ehrenbergiana*, *Acacia tortillis*, *Adenium obesum*, *Aristida adscensionis*, *Asphodelus*
 194 *tenuifolius*, *Catharanthus roseus*, *Cissus rotundifolia*, *Chenopodium murale*, *Chloris*
 195 *barbata*, *Cyperus conglomeratus*, *Delonix elata*, *Ricinus communis* seedling, *Senra*
 196 *incana* and *Tephrosia subtriflora*; cover abundance about 7.11%. **IX-Tamarindus**
 197 **indica** community type, widespread at sandy soils it consists of eight quadrates with
 198 50 species distributed between altitudinal ranges of 650m and 740m a.s.l, associated
 199 with *Abutilon bidentatum*, *Aerva javonica*, *Catharanthus roseus*, *Calotropis procera*,
 200 *Cissus rotundifolia*, *Lawsonia inermis* seedling, *Panicum turgidum*, and *Tephrosia*
 201 *subtriflora* at rough-sandy soils; their cover abundance about 16.2%. **X-Leptadenia**
 202 **arborea** community type represents a large area in the wadi it found in a dry state in
 203 most sites. It consists of six stands with 38 species distributed between altitudinal
 204 ranges of 520m-620m a.s.l, associated with *Adenium obesum*, *Calotropis procera*,
 205 *Cissus rotundifolia*, *Echinochloa colona*, *Indigofera colutea* and *Tamarindus indica*
 206 (sub-shrub); their cover abundance are 10.2%.

207 The Shannon-Wiener diversity index (H') was computed between the ten community
 208 types (Table 5). Community of *Tamarindus indica* (IX) had the highest diversity value
 209 ($H=1.51$) followed by *Leptadenia arborea* community ($H=1.37$). The next dominant

210 community types were *Acacia asak* (H=1.32) and *Ziziphus spina-christ* (H=1.27).
211 *Adenium obesum* and *Anisotes trisulcus* communities had H=1.162 and H=0.880
212 respectively. The lowest diversity index appeared in *Lawsonia inermis* community
213 type being H=0.570 (Table 5). The equitability (evenness) which measures the relative
214 abundance between the different species demonstrated the highest evenness values in
215 community type (VI) *Acacia asak* followed by community type (IX) *Tamarindus*
216 *indica*. The next evenness values were community types of *Leptadenia arborea* (X)
217 and *Dobera glabra* (VIII). The lowest value was for (VII) *Lawsonia inermis*
218 community (Table 5).

219 Similarity and dissimilarity between the different sites calculated by Sorensen's Index
220 coefficient (Table 6) detected the highest values appeared between *Ricinus communis*
221 and *Dobera glabra* (ISs=41.66%) followed by *Adenium obesum* community types and
222 *Tamarindus indica* communities (ISs=36.36%). The lowest similarity estimated
223 between *Ziziphus spina-christi* community types and *Lawsonia inermis* community
224 types (9.09%) followed by community types of *Ricinus communis* and *Tamarindus*
225 *indica* (8.33%).

226 **Discussion**


227 A total of 95 species belonging to 75 genera and 31 families were recorded from wadi
228 Wasaa, Jazan. The floristic analysis revealed four most families of Poaceae,
229 Euphorbiaceae, Malvaceae and Apocynaceae abundant in the wadi. These floristic
230 findings were in accordance with those of Marei *et al.* (2014) on Tihama Hill Slopes
231 and Kasem and Marei (2017) on wadi Khulab, Jazan of Saudi Arabia. The abundance
232 of the Poaceae might be due to water availability, including annual precipitation and

233 soil properties (Osman *et al.*, 2014, and Abdel Abd El-Khalik *et al.* 2017). Life forms
234 were diverse and the vegetation is sparse; chameophytes and therophyte are the
235 dominant, referring to the permanent vegetation that can be accompanied by
236 ephemeral (or annual) plant growth depending on the amount of precipitation in a
237 given year in accordance with finding of E1-Demerdash *et al.* (1994). Moderate of
238 cover abundance in the study area is may be due to the soil is mobile which in
239 accordance to Al-Gifri and Husse (1993) on their studies along the road from Aden to
240 Sheikh Salem (Abyan), Yemen. The intermediate diversity in the wadi was due to the
241 temperate of the rainfall sources and soil fertility which considered as a biotic factor,
242 accordingly the phytoclimate of the wadi was a chameo-therophytic type. The
243 dominance of chaemophytes-therophytes over other life forms is seen to be a response
244 to the hot dry climate, topographic variation and human and animal interference (Abd
245 El-Ghani and Abd El-Khalik, 2006). The high contributions of therophytes lead to
246 adjustment of the flora to water balance. These results are in accordance with several
247 studies in different regions of Saudi Arabia such as: Mosallam (2007) on Sudera, Taif;
248 Al-Turki and Al-Olayan (2003) In Hail Region; Al-Atar, *et al.* (2012) on wadi Al-
249 Jufair; Abd El-Ghani (1993) on Aseer regions and Kasem and Maeri (2017) on wadi
250 Khulab. Biregional area of the Saharo-Arabian, Sudano-Zambeian chorotype were
251 dominated with higher percentage than mono- and pluri-regional area, this was in
252 accordance with Kasem and Marei (2017), El-Shabasy and Kasem (2017) and Osman
253 *et al.* (2014), it represented more than one third of the total species (33-%) because
254 this area mainly deserted and located within the belt of Saharo-Sindian, at the same
255 time it is a part of that belt between Saharo and Sindian. This result was confirmed by

256 the evidence: The ratio of Saharo-Arabian, Sudano-Zambeziian chorotypes decrease
257 while moving to the north and are replaced by Mediterranean and Irano-Turanian
258 chorotypes (Danin and Plitman 1987; Abd El-Ghani and Amer 2003). The studied 46
259 plots were grouped into clusters with the aid of computer program SPSS, ver.20. Ten
260 plant communities were identified and described with varying degrees of species
261 richness, evenness and diversity. The ninth plant community (*Tamarindus indica*)
262 exhibited the highest richness (50 species). Since it is known that the increase in the
263 number samples will increase the species encountered (Mcnaughton and Wolf, 1973);
264 the community types *Salvadora persica* (II) and *Lawsonia inermis* (VII) appeared
265 with the lowest species richness compared to the other community types because they
266 represented species from only two and three sample plots respectively, and could be
267 attributed to variations in their environmental gradients that can limit the ecological
268 distributions of plant species (Lulekal, 2014), it could also be related to the effects of
269 environmental factors such as altitude aspect, soil contents and moisture, human
270 impacts and grazing intensity (Bekele, 1993). Moreover, the area covered by these
271 plants were large in size and occupies vast area of the quadrates. According to Kent
272 and Coker (1992), the Shannon is the most frequently index used for the combination
273 of species richness and relative abundance measurements; the index normally varies
274 between 1.5 and 3.5 and rarely exceeds 4.5. In the present study, the index is between
275 0.57-1.51, showing less even representation of individuals of all species in the
276 sampled quadrats. Sorensen's Index of Similarity (ISs) gives greatest weight to the
277 species that occurred in the two test areas than to those that are unique to either area
278 (Mueller-Dombois and Ellenberg, 1974). According to Sorensen's Index of Similarity

279 (ISs), the highest values calculated within the adjacent sites were in accordance with
 280 results of Tadesse & Bekele (2017). On their studied on the vegetation in Ilu Gelan
 281 district, West Shewa Zone of Oromia region, Central Ethiopia. On the other hand,
 282 Psamophytic species, such as *Senna alexandrina*, *Catharanthus roseus*, *Echinochloa*
 283 *colona*, *Datura stramonium*, *Heliotropium lasiocarpum*, *Cleome viscosa*, *Malva*
 284 *parviflora*, *Cyperus conglomeratus* and *Boerhavia elegans* were recorded from the
 285 sample plots of the wadi bed, this result matches that of Marei *et al.* 2014, as well as,
 286 the association of various species in plant communities (III) *Anisotes trisulcus* and
 287 (VI) *Acacia asak* is in agreement with Masrahi (2012).

288



289

290

Figure 1: A. Location of Jazan in Saudi Arabia, B. Study area in Jazan region

291


292

293

294


295

296




297
298
299
300
301

Figure 2: Monthly average temperature and rainfall percentages in the study area




302
303
304

Figure 3. Species percentages in the recorded families



305
306
307
308
309


Figure 4. Life-form relative spectrum of Wadi Wasaa vegetation. Ch = Chamaephyte, Th= Therophyte, Ph= Phanerophyte, He= Hemi-cryptophyte and Cr= Cryptophyte.



310
311
312
313
314
315
316
317

Figure 5. Floristic category spectrum of Wadi Wasaa. COSM= Cosmopolitan, TR= Tropical, PAN= Pantropical, SA= Saharo-Arabian, SZ = Sudano-Zambeian, ME= Mediterranean and IT= Irano-Turanian.

318




319

Figure 6. Dendrogram showing different plant community types in the study area

320

321



322

323

Figure 7. Diagram showing the abundance of ten plant community types in the wadi.

324

I-*Ziziphus spina-christi*, II- *Salvadora persica* III- *Anisotes trisulcus*, IV- *Adenium*

325

obesum, V- *Ricinus communis*. VI-*Acacia asak*, VII- *Lawsonia inermis*, VIII- *Dobera*

326

glabra, IX-*Tamarindus indica* and X- *Leptadenia arborea*

327

328

329

330

331

332

333 **Table 1:** Collected plant species from Wadi Wasaa with their families, life forms and
 334 chorotypes. Ph, phanerophytes; Ch, chamaephytes; Cr, cryptophyte; H, hemi-
 335 cryptophytes and Th, therophytes, Per=perennial, Ann=Annual,
 336 COSM=Cosmopolitan, IT=Irano-Turanian, ME=Mediterranean, PAN=Panatropical,
 337 SA= Saharo-Arabian, SZ=Sudano-Zambeian and TR=Tropical
 338

Family	Species	Life form	Habit	Life span	Chorotype
Acanthaceae	<i>Anisotes trisulcus</i> (Forssk.) Nees	Ch	Shrub	Per	SA+TR
	<i>Barleria trispinosa</i> (Forssk.) Vahl.	Ch	Sub-shrub	Per	SA
	<i>Blepharis edulis</i> (Forssk.) Pers.	Ch	Sub-shrub	Per	SA+SZ
	<i>Ecboium viride</i> (Forssk.) Alston.	Ph	Sub-shrub	Per	SA
	<i>Ruellia patula</i> Jacq.	Ch	Sub-shrub	Per	SA+TR
Aizoaceae	<i>Trianthema crystalline</i> -Vahl	Th	Herb	Ann	SA
Amaranthaceae	<i>Aerva javanica</i> (Burm.f.) Juss ex Schult.	Ch	Sub-shrub	Ann	SA+ TR
	<i>Amaranthus hybridus</i> L.	Th	Herb	Ann	PAN
	<i>A. viridis</i> L.	Ch	Herb	Ann	ME + TR
	<i>Chenopodium fasciculosum</i> Aellen	He	Herb	Biennial	SA+TR
	<i>C. carinatum</i> R. Br.	Th	Herb	Ann	SA+SZ
Apocynaceae	<i>Adenium obesum</i> (Forssk.) Roem. & Schult.	Ph	tree	Ann	SA
	<i>Calotropis procera</i> (Aiton) Dryand.	Ch	shrub	Per	SA+SZ
	<i>Caralluma retrospiciens</i> (Ehrenb.) N.E.Br.	Ch	Succulent	Per	SA+SZ
	<i>Catharanthus roseus</i> (L.) G.Don.	Ch	Herb	Ann	ME + TR
	<i>Kanahia laniflora</i> (Forssk.) R. Br.	Ch	Sub-shrub	Per	SA+SZ
	<i>Leptadenia arborea</i> (Forssk.) Schweinf	Ch	Climber	Per	SA+ SZ
	Asphodelaceae	<i>Aloe officinalis</i> Forssk.	He	Succulent	Per
	<i>Asphodelus tenuifolius</i> Cav.	Cr	Herb	Biennial	SA+ SZ
Astraceae	<i>Conyza steudelii</i> Sch.Bip. ex A.Rich.	Ph	Sub-shrub	Per	SA+TR
	<i>Cyanthillium cinereum</i> (L.) H.Rob.	Th	Herb	Ann	SZ +TR
	<i>Eclipta prostrata</i> (L) L.	Th	Herb	Per	ME+ SA
	<i>Pluchea dioscoridis</i> (L.) DC.	Ch	Sub-shrub	Per	SA+SZ
	<i>Pulicaria schimperi</i> DC.	Ch	Herb	Ann	SA+TR
Boraginaceae	<i>Heliotropium longiflorum</i> (A.DC.) Jaub. & Spach	He	Herb	Per	SA+TR
	<i>H. pterocarpum</i> (DC.&A.DC.) Hochst. & Steud. ex Bunge	He	Herb	Per	SA+SZ
	<i>H. strigosum</i> Willd.	He	Herb	Per	SA+SZ
Capparaceae	<i>Capparis cartilaginea</i> Decne.	Ch	Sub-shrub	Per	SA+SZ
Cactaceae	<i>Opuntia dillenii</i> (Ker Gawl.) Haw.	Ch	Shrub	Per	TR
Caesalpinaceae	<i>Senna alexandrina</i> Mill.	Ch	Sub-shrub	Per	SA+SZ
	<i>S. italica</i> Mill.	Ch	Sub-shrub	Per	SZ
	<i>Tamarindus indica</i> L.	Ph	Tree	Per	SA+TR
Cleomaceae	<i>Cleome scaposa</i> DC.	He	Herb	Ann	SA+TR
	<i>C. viscosa</i> L.	Th	Herb	Ann	PAN
Clesteraceae	<i>Gymnosporia senegalensis</i> (Lam.) Loes.	Ph	Tree	Per	SA
Convolvulaceae	<i>Merremia aturensis</i> (Kunth) Hallier f	He	Climber	Per	SA+SZ
Cyperaceae	<i>Cyperus conglomeratus</i> Rottb.	Cr	Herb	Per	SA
Euphorbiaceae	<i>Acalypha fruticosa</i> Forssk.	Ch	Sub-shrub	Per	SA
	<i>Chrozophora oblongifolia</i> (Del.) A. Juss. ex Spreng.	Ph	Sub-shrub	Per	SA+SZ
	<i>Dalechampia scandens</i> L.	Ph	Climber	Per	SA+TR

	<i>Euphorbia inarticulata</i> Schlecht.	Ch	Succulent	Per	ME+SA+SZ
	<i>E. hirta</i> L.	Th	Herb	Ann	COSM
	<i>E. prostrata</i> Aiton.	Th	Herb	Per	COSM
	<i>E. triaculeata</i> Forssk	Ch	Succulent	Per	SZ
	<i>Ricinus communis</i> L.	Ph	Tree	Per	TR
Lamiaceae	<i>Lavandula coronopifolia</i> Poir.	Ch	Sub-shrub	Ann	ME+SA
	<i>Ocimum forsskaolii</i> Benth.	Ch	Sub-shrub	Ann	SA+TR
Lythraceae	<i>Lawsonia inermis</i> L.	Ph	Tree	Per	Cultivated
Malvaceae	<i>Abutilon bidentatum</i> Hochst. ex A.Rich	Ch	Sub-shrub	Per	TR
	<i>A. hirtum</i> (Lamk.) Sweet	Ch	Sub-shrub	Per	PAN
	<i>Corchorus depressus</i> (L.) Stocks	Cr	Herb	Per	ME+IT
	<i>C. tridens</i> L.	Th	Herb	Ann	ME+SA+SZ
	<i>Malva parviflora</i> L.	He	Herb	Ann	ME+ IT
	<i>Senra incana</i> Cav.	Ch	Sub-shrub	Per	SA+SZ
Mimosaceae	<i>Acacia asak</i> (Forssk.) Willd	Ph	Tree	Per	SA+SZ
	<i>A. ehrenbergiana</i> Hayne	Ph	Tree	Per	SA+SZ
	<i>A. tortilis</i> (Forssk.) Hayne	Ph	Tree	Per	SA+SZ
	<i>Delonix elata</i> (L.) Gamble	Ph	Tree	Per	SA+TR
Moraceae	<i>Ficus cordata</i> ssp. <i>salicifolia</i> (Vahl) Berg.	Ph	Shrub	Per	SA
	<i>F. ingens</i> (Miq.) Miq.	Ph	Tree	Per	SA+SZ
Nyctaginaceae	<i>Boerhavia elegans</i> Choisy	He	Herb	Ann	TR
	<i>Commicarpus grandiflorus</i> (Rich.) Standley	He	Herb	Per	SA+TR
Papavaraceae	<i>Argemone ochroleuca</i> Sweet	Th	Herb	Per	PAN
Papilionaceae	<i>Crotalaria microphylla</i> M.Vahl.	Th	Herb	Per	SA+SZ
	<i>Indigofera colutea</i> (Burm.f.) Merr.	Ch	Sub-shrub	Per	SZ
	<i>I. hochstetteri</i> Bak.	Ch	Sub-shrub	Ann	SZ
	<i>I. spinosa</i> Boiss.	Ch	Sub-shrub	Per	SA+SZ
	<i>Tephrosia subtriflora</i> Baker	Ch	Sub-shrub	Per	SA
Plantaginaceae	<i>Scoparia dulcis</i> L.	Ch	Herb	Per	SA
	<i>Schweinfurthia pterosperma</i> A. Braun	Th	Herb	Ann	SA
Poaceae	<i>Aristida adscensionis</i> L.	Th	Herb	Ann	ME+SA
	<i>Cenchrus ciliaris</i> L.	Cr	Herb	Per	SA+ SZ
	<i>Chloris barbata</i> Sw.	Th	Herb	Per	SZ
	<i>C. gayana</i> Kunth	Th	Herb	Per	SA+SZ
	<i>Dichanthium foveolatum</i> (Del.) Roberty	Th	Herb	Per	ME+SA+SZ
	<i>Echinochloa colona</i> (L.) Link.	Th	Herb	Ann	TR
	<i>Eragrostis japonica</i> (Thunb.) Trin.	Th	Herb	Ann	SA+ SZ
	<i>E. papposa</i> (Roem & Schult) Steud.	Th	Herb	Per	SZ
	<i>Hyparrhenia hirta</i> (L.) Stapf	Th	Herb	Per	SA
	<i>Panicum turgidum</i> Forssk.	Cr	Herb	Per	SA+SZ
	<i>Paspalidium desertorum</i> (Rich.) Stapf.	Th	Herb	Per	SA
	<i>Sorghum bicolor</i> (L.) Moench	Cr	Herb	Ann	Cultivated
	<i>Sporobolus nervosus</i> Hochst.	Th	Herb	Per	SA+SZ
	<i>Tetrapogon cenchriformis</i> (Rich.) Clayton	Th	Herb	Ann	SA+SZ
Rhamnaceae	<i>Ziziphus spina-christi</i> (L.) Desf.	Ph	Tree	Per	ME+SA+SZ
Salvadoraceae	<i>Dobera glabra</i> (Forssk.) Juss. ex Poir	Ph	Tree	Per	SA+TR
	<i>Salvadora persica</i> L.	Ch	shrub	Per	SA+SZ
Solanaceae	<i>Datura innoxia</i> Mill.	Ch	Sub-shrub	Ann	SA
	<i>D. stramonium</i> L.	Th	Sub-shrub	Ann	COSM
	<i>Solanum surattense</i> Burm. F.	Th	Herb	Per	SA+TR
Urticaceae	<i>Forsskaolea tenacissima</i> L.	Th	Herb	Per	SA+SZ
Vitaceae	<i>Cissus rotundifolia</i> Vahl	Ch	Climber	Per	SA
Zygophyllaceae	<i>Fagonia indica</i> Burm.F.	He	Herb	Per	SA+IT

	<i>F. paulayana</i> J.Wagner & Vierh.	Th	Herb	Per	SA+SZ
	<i>Tribulus parvispinus</i> C. Presl	Th	Herb	Ann	SA+SZ

339
340
341
342

Table 2: Species number related to main floristic categories and their phytochoria percentage.

Growth type		Phytochoria				Life Form		
Type	%	Category	Type	No.	%	Form	No.	%
Annual	27	Monoregional	SA	15	16	Ch	33	35
Biennial	02		TR	5	5	Th	28	29
Perennial	66		SZ	6	6	Ph	17	18
		Biregional	SA+SZ	31	33	He	11	12
			SA +TR	16	16	Cr	6	6
			ME + IT	2	2	--	--	
			SZ+TR	1	1	--	--	
			ME+TR	2	2	--	--	
			ME+SA	4	4	--	--	
			SA +IT	1	1	--	--	
		Pleuriregional	ME+ SA +SZ	4	4	--	--	
			PAN	4	4	--	--	
			COSM	3	3	--	--	
			Cult	2	2	--	--	
			Total	14	95	100	5	95

343
344
345
346
347
348
349

Table 3: Plant species and their cover abundance in all quadrates among the dominant communities. I-*Ziziphus spina-christi*, II- *Salvadora persica* III- *Anisotes trisulcus*, IV- *Adenium obesum*, V- *Ricinus communis*. VI-*Acacia asak*, VII- *Lawsonia inermis*, VIII- *Dobera glabra*, IX-*Tamarindus indica* and X- *Leptadenia arborea*

Species	Community Types										Cover abundance
	I	II	III	IV	V	VI	VII	VIII	IX	X	
<i>Abutilon hirtum</i>	-	-	+	-	-	+	-	-	-	-	0.008
<i>Abutilon bidentatum</i>	-	-	-	-	-	-	-	-	+	-	0.060
<i>Acalypha fruticosa</i>	-	-	-	-	+	-	-	-	-	-	0.050
<i>Acacia asak</i>	-	-	+	-	-	-	-	+	-	-	13.10
<i>A. ehrenbergiana</i>	-	-	-	-	-	-	-	+	-	-	0.708
<i>A. tortilis</i>	-	-	+	-	+	-	-	+	-	-	0.300
<i>Adenium obesum</i>	+	-	-	-	-	-	-	+	-	+	8.150
<i>Aerva javanica</i>	-	-	+	-	-	+	-	-	+	-	0.045
<i>Aloe officinalis</i>	-	+	-	-	-	-	-	-	-	-	0.030
<i>Anisotes trisulcus</i>	+	+	-	+	-	+	-	-	-	-	9.530
<i>Aristida adscensionis</i>	-	-	-	-	-	-	+	+	-	-	0.007
<i>Argemone ochroleuca</i>	+	-	-	-	-	+	-	-	-	-	0.027
<i>Asphodelus tenuifolius</i>	-	-	-	-	+	-	+	+	-	-	0.005
<i>Barleria trispinosa</i>	+	-	-	-	-	-	-	-	-	-	0.077
<i>Blepharis edulis</i>	-	+	-	-	-	-	-	-	-	-	0.047
<i>Catharanthus roseus</i>	-	-	-	-	+	+	-	+	+	-	0.004
<i>Calotropis procera</i>	-	+	-	-	-	-	-	-	+	+	0.024

<i>Caralluma retrospiciens</i>	+	-	-	-	-	-	-	-	-	-	0.069
<i>Cenchrus ciliaris</i>	-	-	-	+	+	-	-	-	-	-	0.054
<i>Cissus rotundifolius</i>	-	-	-	-	-	-	-	+	+	+	0.008
<i>Chenopodium murale</i>	-	-	-	-	+	-	-	+	-	-	0.023
<i>Chloris barbata</i>	-	-	-	-	+	-	-	+	-	-	0.025
<i>Cleome viscosa</i>	-	-	-	-	-	-	+	-	-	-	0.006
<i>Cleome scaposa</i>	-	-	+	-	-	-	-	-	-	-	0.004
<i>Corchorus depressus</i>	-	+	-	-	-	-	-	-	-	-	0.054
<i>Cyperus conglomeratus</i>	-	-	-	-	+	-	+	-	-	-	0.005
<i>Delonix elata</i>	-	-	-	-	-	-	-	+	-	-	0.070
<i>Dobera glabra</i>	-	-	-	-	+	-	-	-	-	-	7.110
<i>Eragrostis papposa</i>	-	-	-	+	-	-	-	-	-	-	0.011
<i>Eclipta prostrate</i>	-	-	-	+	-	-	-	-	-	-	0.004
<i>Euphorbia triaculeata</i>	-	+	+	-	-	-	-	-	-	-	0.008
<i>Echinochloa colona</i>	-	-	-	-	-	-	-	-	-	+	0.003
<i>Fagonia indica</i>	-	-	-	-	-	+	-	-	-	-	0.007
<i>Forsskaolea tenacissima</i>	-	-	+	-	-	-	-	-	-	-	0.004
<i>Heliotropium longiflorum</i>	+	-	-	-	-	-	-	-	-	-	0.004
<i>Indigofera colutea</i>	-	+	-	-	-	-	-	-	-	+	0.005
<i>Indigofera spinosa</i>	-	-	+	-	-	-	-	-	-	-	0.006
<i>Leptadenia arborea</i>	-	-	+	-	-	-	-	-	-	-	10.20
<i>Lavandula coronopifolia</i>	+	-	+	-	-	-	-	-	-	-	0.005
<i>Lawsonia inermis</i>	-	-	-	+	-	-	-	-	+	-	2.710
<i>Maytenus senegalensis</i>	-	-	+	-	-	-	-	-	-	-	0.080
<i>Malva parviflora</i>	-	-	-	-	-	-	+	-	-	-	0.002
<i>Ocimum forsskaolii</i>	-	+	-	-	-	-	-	-	-	-	0.025
<i>Panicum turgidum</i>	-	-	-	+	-	-	-	-	+	-	0.028
<i>Pluchea dioscoridis</i>	-	-	-	-	+	-	-	-	-	-	0.010
<i>Paspalidium desertorum</i>	-	-	-	-	-	-	+	-	-	-	0.011
<i>Ricinus communis</i>	-	-	-	-	-	-	-	+	-	-	9.200
<i>Senna alexandrina</i>	-	-	-	-	-	-	+	-	-	-	0.007
<i>Senra incana</i>	-	-	-	-	-	-	-	+	-	-	0.026
<i>Tamarindus indica</i>	-	-	-	-	-	-	-	-	-	+	16.02
<i>Tephrosia subtriflora</i>	-	-	-	-	-	-	-	+	+	-	0,050
<i>Trianthema crystallina</i>	+	-	-	-	-	-	-	-	-	-	0.004
<i>Tribulus parvispinus</i>	+	-	-	-	-	-	-	-	-	-	0.005
<i>Ziziphus spina-christi</i>	-	+	-	-	-	-	-	-	-	-	10.40

350
351
352
353
354
355
356
357
358
359
360
361
362

363

364

365

Table 4: Distribution of the ten plant communities with their altitudinal ranges.

Name	Altitude		Plot Number	Plot list
	Ranges (m a.s.l.)	Means (m a.s.l.)		
I- <i>Ziziphus spina-christi</i>	500-570	616.13	5	1, 5, 6, 8 & 11
II- <i>Salvadora persica</i>	520-586	552.66	3	2, 4 & 9
III- <i>Anisotes trisulcus</i>	450-510	498.24	4	3, 7, 10 & 14
IV- <i>Adenium obesum,</i>	550-640	585.43	4	12, 13, 15 & 26
V- <i>Ricinus communis</i>	450-510	482.53	4	16, 17, 18 & 19
VI- <i>Acacia asak</i>	550-580	564.60	7	21, 24, 25, 27, 28, 29 & 23
VII- <i>Lawsonia inermis</i>	630-690	650.65	2	30 & 31
VIII- <i>Dobera glabra</i>	460-570	503.30	3	20, 22 & 23
IX- <i>Tamarindus indica</i>	650-740	685.32	8	33, 24, 35, 36, 37, 39, 41 & 46
X- <i>Leptadenia arborea</i>	520-620	576.66	6	38, 40, 43, 44, 45 & 42

366

367

368

369

370

Table 5: Shannon-Wiener diversity index for ten plant community types

Communities	Species richness (S)	Diversity index (H')	H max	Species evenness (J)
I- <i>Ziziphus spina-christi</i>	35	1.271	1.382	0.060
II- <i>Salvadora persica</i>	18	0.692	0,761	0.040
III- <i>Anisotes trisulcus</i>	29	0.880	0,940	0.040
IV- <i>Adenium obesum</i>	32	1.162	1.260	0.052
V- <i>Ricinus communis</i>	27	0.781	0.850	0.043
VI- <i>Acacia asak</i>	42	1.322	1.461	0.084
VII- <i>Lawsonia inermis</i>	15	0.570	0.631	0.032
VIII - <i>Dobera glabra</i>	22	0.742	0,811	0.061
IX- <i>Tamarindus indica</i>	50	1.513	1.652	0.080
X- <i>Leptadenia arborea</i>	38	1.370	1.493	0.071

371

372

373

374

375

376

Table 6: Similarity and dissimilarity between the 10 community types calculated by Sorensen's similarity coefficient (ISs)

Communities	<i>Ziziphus spina.</i>	<i>Salvadora persica</i>	<i>Aniso. trisulcus</i>	<i>Adenium obesum</i>	<i>Ricinus communis</i>	<i>Acacia asak</i>	<i>Lawsonia inermis</i>	<i>Dobera glabra</i>	<i>Tamarindus indica</i>
<i>Ziziphus spina-christi</i>	0.00								
<i>Salvadora persica</i>	30.7	0.00							
<i>Anisotes trisulcus</i>	24.3	30.10	0.00						
<i>Adenium obesum,</i>	27.2	15.60	28.5	0.00					
<i>Ricinus communis</i>	29.1	33.30	32.8	27.27	0.00				
<i>Acacia asak</i>	28,5	28.50	33.0	30.7	18.8	0.00			
<i>Lawsonia inermis</i>	9.09	18.50	16.6	29.0	16.6	19.0	0.00		
<i>Dobera glabra</i>	21.42	23.01	24.2	17.91	41.66	13.0	9.51	0.00	
<i>Tamarindus indica</i>	23.07	15.38	16.4	36.36	8.33	12.8.	29.1	14.2	0.00
<i>Leptadenia arborea</i>	16.66	13.04	18.18	19.4	12.6	10.2	18.18	20.0	20.0

377

378

379 **References**

- 380 Abd El-Ghani, M. (1993). Habitat features and plant communities of the Holy Places,
381 Mecca, Saudi Arabia. *Feddes Repertorium* 104:417–25.
- 382 Abd El-Ghani, M. and Abd El-Khalik, K. (2006). Floristic diversity and
383 phytogeography of the Gebel Elba national park, southeast Egypt. *Turk. J. Bot.*;
384 30:121-136.
- 385 Abd El-Ghani, M. and Amer, W.M. (2003). Soil-vegetation relationships in a coastal
386 desert plain of southern Sinai, Egypt. *J. of Arid Environments* 55:607–28.
- 387 Abdel Khalik, K., El-Sheikh, M., El-Aidarous, A., (2013). Floristic diversity and
388 vegetation analysis of Wadi Al-Noman, Holy Mecca, Saudi Arabia. *Turk. J Botany*
389 37, 894–907.
- 390 Abd El-Khalik, K. Iman, A. and Yassin A. (2017). Floristic composition and
391 vegetation: Environmental relationships of Wadi Fatimah, Mecca, Saudi Arabia,
392 *Arid Land Research and Management*, Vol.31: 316-334
- 393 Alatar A., El-Sheikh M.A., Thomas J. (2012). Vegetation analysis of Wadi Al-Jufair,
394 a hyper-arid region in Najd, Saudi Arabia. *Saudi Journal of Biological Sciences*,
395 19, 357–368
- 396 Al-Farhan, A.H., Al Turkey, T.A., Basahya, Y. (2005). *Flora of Jizan region*. Final
397 report supported by King Abdul-Aziz City for Science and Technology;1,2:545.
- 398 Al-Gifri, A.& Hussein M. A. (1993). Plant Communities along the road from Aden to
399 Sheilkh Saleim (Abyan). *Feddes. Report*. 104 : 267-270.

- 400 Al-Turki, T.A., Al-Qlayan, H.A., (2003). Contribution to the flora of Saudi Arabia:
401 Hail region. *Saudi Journal of Biological Sciences*, 10:190–222.
- 402 Al-Wadie, H., (2002). Floristic composition and vegetation of Wadi Talha, Aseer
403 Mountains, south west Saudi Arabia. *Journal of Biological Sciences* 2 (5), 285–288.
- 404 Al-Yemeni, M.N., (2001). Ecology of some plant communities in Wadi Al-Ammaria,
405 Riyadh, Saudi Arabia. *Saudi Journal of Biological Sciences* 8 (2), 145–165.
- 406 Atiqur, R.M. Mossa J.S., Al-Said, M.S., Al-Yahya M.A. (2004). Medicinal plant
407 diversity in the flora of Saudi Arabia 1: A report on seven plant families. *Fitoterapia*
408 75,149-161.
- 409 Barbourm, G., Burk, J.H. and Pitts W.D. (1987). *Terrestrial Plant Ecology*. 2nd ed.
410 Massachusetts: Benjamin/Cummings.
- 411 Bekele, T. (1993). *Vegetation ecology of remnant Afromontane forests on the Central*
412 *Plateau of Shewa, Ethiopia*. Uppsala, Sweden.
- 413 Chaudhary, S.A. (2001). In: *Flora of the Saudi Arabia Kingdom*. Ministry of
414 Agriculture and Water press, Riyadh; 1-3.
- 415 Danin, A. and Plitman, U. (1987). Revision of the plant geographical territories of
416 Israel and Sinai. *Plant Systematics and Evolution*, 156:43–53.
- 417 El Ghazali G.E.B., Al- Soqeer A.R.A. and El Tayeb G.E.A. (2013). Floristic and
418 ecological studies on the plant cover of Wadi Al Rummah, Qassim Region, Saudi
419 Arabia. *Int. Res. J. Plant Sci.* 4(10):310-318

- 420 EL Ghenem, W.M., (2006). Ecological study at wadi Al-Ammaria in El-Riyadh city-
421 Saudi Arabia. *Bulletin of Pure and Applied Sciences*, Section B 25 (1), 11–19.
- 422 El-Demerdash, M.A., Hegazy, A.K And Zilay, A.M. (1994). Vegetation-soil
423 relationships in Tihamah coastal plains of Jazan region, *Saudi Arabia. J. of Arid*
424 *Environments*; 30:161-174.
- 425 El-Shabasy, A. and Kasem,W. (2018). Systematic composition, species diversity and
426 plant chorology at Wadi Tashar, Jazan, Saudi Arabia, *J. of Medicinal Plants Studies*;
427 6(1): 83-88.
- 428 Farrag HF (2012). Floristic composition and vegetation-soil relationships in Wadi Al-
429 Argy of Taif region, Saudi Arabia. *Int. Res. J. of Pl. Sci.* 3(8): 147-157.
- 430 Kasem, W.T. and Marei, H.A. (2017). Floristic Compositions and Its Affinities to
431 Phytogeographical Regions in Wadi Khulab of Jazan, Saudi Arabia. *International*
432 *Journal of Plant & Soil Science*; 16(3):1-11.
- 433 Kassas, M. and Girgis W.A. (1964). Habitat and plant communities in the Egyptian
434 desert. V. The limestone plateau. *Journal of Ecology*, 52: 107–119.
- 435 Kent M. and P. Coker (1992). *Vegetation Description and Analysis: A practical*
436 *approach*. John Wiley and Sons, New York, 363 p.
- 437 Lulekal, E. (2014). *Plant Diversity and Ethnobotanical Study of Medicinal Plants in*
438 *Ankober District, North Shewa Zone of Amhara Region, Ethiopia*, Ph.D.
439 Dissertation. Addis Ababa University, Ethiopia.

- 440 Maarel. E. (1979). Transformation of cover-abundance values in phytosociology and
441 its effects on community similarity. *Vegetation* 39: 97–114.
- 442 Marei A., Kasem, W. and A. Gafar. (2014). Phytosociological studies of the southern
443 Sector of Tihama Hill Slopes of Jazan region, south west of Saudi Arabia. *Asian*
444 *Journal of Applied Sciences*; 2:734-744.
- 445 Masrahi, Y.A. (2012). *Brief illustrate to wild plants in Jizan region*. King Fahad
446 Library, Jeddah.; 302.
- 447 Mcnaughton S. and Wolf. L. (1973). *General Ecology-Holt, Rinehart and Winston,*
448 Inc. New York
- 449 Migahid, A.M. (1996). *Flora of Saudi Arabia*, Jeddah: King Abdul Aziz University
450 Press.;1-3.
- 451 Mosallam, H.A., (2007). Comparative study on the vegetation of protected and non-
452 protected areas, Sudera, Taif, Saudi Arabia. *Int. J. Agric. Biol.* 9, 202–214.
- 453 Mueller-Dombois D., Ellenberger, H. (1974). *Aims and Methods of Vegetation*
454 *Ecology*. New York: John Wiley and Sons Inc. 547 pp.
- 455 Orshan, G.(1986). The desert of the Middle East. In: Evenari M, Noy-Meir I &
456 Goodall DW (eds.) *Ecosystems of the World*. Vol. 12B,pp. 1–28.
- 457 Osman, A., Al-Ghamdi, F. and Bawadekji, A. (2014). Floristic diversity and
458 vegetation analysis of Wadi Arar: A typical desert Wadi of the Northern Border
459 region of Saudi Arabia. *Saudi Journal of Biological Sciences* 21, 554–565

- 460 Raunkiaer, C. (1937). *Life forms of plants and statistical plant geography*. Oxford:
461 Clarendon Press.
- 462 Shannon C.E. and Wiener. W. (1949). *The Mathematical Theory of Communication*.
463 University of Illinois, Chicago, USA.
- 464 Siraj, M., Zhang, K., Sebsebe, D. and Zerhiun W. (2016). Floristic composition and
465 plant community types in Maze National Park, southwest Ethiopia. *Applied ecology*
466 *and environmental research*, 15 (1):245-262.
- 467 Tadesse, Z, and Bekele, T. (2017). Floristic composition and plant community
468 analysis of vegetation in Ilu Gelan district, West Shewa Zone of Oromia region,
469 Central Ethiopia. *Tropical Plant Research* 4 (2): 335–350
- 470 The Plant List database (2013). Web site (<http://www.theplantlist.org/>).
- 471 Wickens, G.E. (1978). Some of the phytogeographical problems associated with
472 Egypt. *Publications Cairo University Herbarium*, 7-8:223-230.
- 473 Zohary, M. (1973). Geobotanical foundations of the Middle East. *Stuttgart: Gustav*
474 *Fischer Verlag*.

