

ANTIMICROBIAL ACTIVITY OF *PLEUROTUS SQUARROSULUS* ON CLINICAL PATHOGENIC BACTERIA AND FUNGI

ABSTRACT

Aim: To evaluate the antimicrobial activities of *Pleurotus squarrosulus* mushroom extracts on bacterial and fungal isolates.

Study design: *Pleurotus squarrosulus* was obtained from different sources in Umuahia North Local Government, Abia state, Nigeria and identified in the Department of botany, University of Nigeria, Nsukka.

Place and duration of study: Antimicrobial activities of *Pleurotus squarrosulus* was carried out in the department of microbiology between January 2016 and August 2016

Methodology: *Pleurotus squarrosulus* was extracted using ethanol, methanol and aqueous. Antimicrobial susceptibility tests were carried out by agar well diffusion technique using National Committee of Clinical Laboratory Standard. Qualitative phytochemical analysis was carried out using standard methods.

Results: Methanol, ethanol and aqueous extracts of *Pleurotus squarrosulus* were tested against *E.coli*, *B. cereus*, *S. aureus*, *P. aeruginosa*, *C. albicans* and *C. glabrata*. The different test microorganisms showed varied susceptibility to the test extracts. All the test organisms were inhibited by methanol, ethanol and aqueous extract at varied concentrations ranging between 500 mg/ml and 125 mg/ml. Statistically, inhibition of the antibacterial and antifungal control for the test organisms were significantly higher ($P < 0.05$) than that of the extracts. The phytochemical analysis revealed the presence of saponin, carbohydrates, tannins, flavonoids and proteins in all the extracts while glycoside and alkaloids, were found in some.

Conclusion: The finding of this result suggest that *Pleurotus squarrosulus* possess broad-spectrum antimicrobial activity. The potential of developing antimicrobials from plants appear rewarding.

Key words: *Pleurotus squarrosulus*, antimicrobial activities, phytochemicals, bacteria, yeast

1. INTRODUCTION

Time immemorial, mushrooms have been used as a part of regular diet due to their nutritional and medicinal values [1]. Mushrooms have been found to contain minerals, vitamins and nutritive compounds, proteins, polysaccharide and a low fat content [2]. Mushrooms are also rich sources of natural antibiotics. Their cell wall glucans have been known to poses immunomodulatory properties with many of their secondary metabolites combating bacteria, fungi and viruses [3, 4, 5, 6, 7, 8, 9]. Prior to the discovery of their high medicinal value, mushrooms have been used for hundreds of years in traditional medicine for curing various types of diseases such as antimicrobial, antioxidant, antiviral, anticancer, antitumor, anti-inflammatory, cardiovascular diseases, immunomodulating, central activities [10, 11].

Besides, mushroom has been used extensively in traditional medicine for curing of various types of diseases [12, 13, 14]. For centuries, mushrooms have been prescribed for treatment of diseases such as gastro-intestinal disorder, bleeding, high blood pressure and various bacterial infections [15]. While some of the medicinal values associated with mushroom must have arisen from surperstitious beliefs and

39 myths, they have provided information for curiosity research studies. Research has shown that some of
40 these claims are not mere myth but are authentic [16, 17]. Besides medicinal and nutritional use,
41 mushroom can be used as natural dyes for fabrics [18].

42 *Pleurotus squarrosulus* is a common edible mushroom. It was first cultivated in Germany as a
43 subsistence measure during World War I [19] and is now grown commercially around the world for food. It
44 is related to the similarly cultivated "king oyster mushroom". *Pleurotus squarrosulus* can also be used
45 industrially for mycoremediation purposes. *Pleurotus squarrosulus* is one of the more commonly sought
46 wild mushrooms, though it can also be cultivated on straw and other media. It has the bittersweet aroma
47 of benzaldehyde (which is also characteristic of anise or almonds) [20].

48 The mushroom has a broad cap spanning 5–25 cm; natural specimens range from white to gray or tan to
49 dark-brown; the margin is inrolled when young, and is smooth and often somewhat lobed or wavy. The
50 flesh is white, firm, and varies in thickness due to stipe arrangement. The gills of the mushroom are white
51 to cream, and descend on the stalk if present. If so, the stipe is off-center with a lateral attachment to
52 wood. The spore print of the mushroom is white to lilac-gray, and best viewed on dark background. The
53 mushroom's stipe is often absent. When present, it is short and thick. Due to the dearth in literature on the
54 dual value of *Pleurotus squarrosulus* as food and its antimicrobial efficacy, this study was designed.

55 2. MATERIALS AND METHODS

56 2.1 Collection and identification of materials

57 *Pleurotus squarrosulus* was collected from different sources of Umuahia North Local Government area,
58 Abia state and identified by a botanist in the Department of botany, University of Nigeria, Nsukka.

59 2.2 Test organisms used

60 Pure cultures of *Escherichia coli* JCM 20135 and *Bacillus cereus* IFO 13804 were obtained from
61 Department of Microbiology, University of Nigeria Nsukka while pure cultures of *Staphylococcus aureus*
62 ATCC 25923, *Candida albicans* ATCC 10231, *Pseudomonas aeruginosa* ATCC 25783 and *Candida*
63 *glabrata* ATCC 22018 were obtained from Spectramedics Laboratories No. 2 Adebayo close, Araromi
64 Offiri Ikenne Road, Sagamu, Ogun State.

65

66 2.3 Standard antimicrobials

67 Tetracycline (5 µg/ml), Gentamycin (5 µg/ml), Ampicillin (5 µg/ml), Oxacillin (5 µg/ml), Fluconazole (5
68 µg/ml) and Nystatin (20 µg/ml) oxoid disk were used as positive standards.

69 2.4 Sample preparation and extraction

70 Fresh *Pleurotus squarrosulus* mushrooms were thoroughly washed with ~~clean water~~, cut into
71 pieces, air-dried at room temperature and pulverized using manual grinder. Fifty grams of each
72 of the ground samples was soaked in 300 ml ethanol, cold water, and methanol for 24 hours
73 with intermittent shaking. Each sample was filtered using Whatman №1 filter paper. The filtrate
74 was poured into a crucible and allowed to dry under steady air current in order to obtain the
75 extract which was scooped and poured into well-labeled sample bottles and stored at 4°C [21].

76 77 2.5 Inoculum preparation

78 Pure cultures of *Escherichia coli* JCM 20135 and *Bacillus cereus* IFO 13804 were obtained from the
79 Department of Microbiology, University of Nigeria Nsukka while pure cultures of *Staphylococcus aureus*
80 ATCC 25923, *Candida albicans* ATCC 10231, *Pseudomonas aeruginosa* ATCC 25783 and *Candida*
81 *glabrata* ATCC 22018 were obtained from Spectramedics Laboratories, Sagamu, Ogun State, Nigeria.
82 Inoculum was prepared by emulsifying overnight colonies from an agar medium. A 0.5 McFarland
83 standard (equivalent to approximately 10⁸cfu/ml) was used. Media plates were inoculated within 30
84 minutes of standardizing the inoculum to avoid changes in inoculums density.

85 86 2.6 Determination of antimicrobial activity of mushroom extracts

87 Antimicrobial activity of mushroom extracts was determined according to the National
88 Committee of Clinical Laboratory Standards [22]. Agar well diffusion method on Sabouraud
89 dextrose agar (SDA) and Muller-Hinton agar were used for fungi and bacteria respectively. Up
90 to 100 µl of the inoculum was poured onto the agar plate and spread with glass rod under sterile
91 conditions. Wells (6mm diameter) were bored into the agar using sterile cork-borer and 0.1 ml of
92 different concentrations of the extracts (500, 250, 125, 62.5, 31.25, 15.63 and 7.81 mg/ml) was
93 applied into each well. Negative control wells were filled with dilute dimethylsulfoxide while
94 positive controls were antibiotic discs of tetracycline (10 µg/ml); ampicillin (10 µg/ml) for Gram
95 negative bacteria isolates and oxacillin (5 µg/ml); gentamicin (10 µg/ml) for Gram positive
96 bacteria isolates. Antifungal discs of fluconazole (25 µg/ml) and nystatin (20 µg/ml) (Oxoid,
97 United Kingdom) were used as positive controls for fungal isolates. This procedure was done in
98 triplicate for the entire test organisms, allowed to stand for 30 minutes on the bench and
99 incubated for 24 hours at 37±2 °C for bacteria and 72 hours at 28±2 °C for yeast. After

100 incubation, the inhibition zone diameters produced by the different concentrations of the crude
101 extracts were measured (in millimeter) and recorded. Antimicrobial activities were expressed in
102 terms of the mean value of the inhibition zone produced by the mushroom extracts.

103

104 **2.7 Determination of minimum inhibitory concentrations (MICs) of the mushroom extracts**

105 The MIC of the extracts was determined for the test organisms in triplicates at varying concentrations
106 (250, 125, 62.5, 31.25, 15.62, 7.80 and 3.90 mg/ml). To obtain these concentrations, 1.0 ml of varying
107 concentrations of the extracts with double strength (500, 250, 125, 62.5, 31.25, 15.62 and 7.80 mg/ml)
108 were constituted in different test tubes. About 1.0 ml of Mueller-Hinton broth (for bacteria) and Sabouraud
109 dextrose broth (for fungi) was added and then a loopful of the test organism, previously diluted to 0.5
110 McFarland turbidity standard, was introduced. Controls of Mueller-Hinton broth and Sabouraud dextrose
111 broth without the mushroom extract were set up. All the bacterial cultures were incubated at $37 \pm 2^\circ\text{C}$ for
112 24 hours and yeast culture incubated at $28 \pm 2^\circ\text{C}$ for 72 hours. After incubation each tube was examined
113 for microbial growth. The lowest concentration of the extract that inhibited the growth of the test
114 organisms as detected by lack of visual turbidity was designated the MIC [23].

115

116 **2.8 Determination of minimum bactericidal concentrations (MBCs) of the mushroom** 117 **extracts**

118 MBC was determined by selecting tubes that showed no bacterial growth during the MIC
119 determination. A loopful from each of the tubes was sub-cultured on the Mueller Hinton Agar
120 and incubated for 24 hours at $37^\circ\text{C} \pm 2^\circ\text{C}$. The MBC was determined as the least concentration
121 that showed no visible growth on the plate [23].

122

123 **2.9 Determination of minimum fungicidal concentrations (MFCs) of the mushroom** 124 **extracts**

125 MFC was determined by selecting tubes that showed no fungal growth during MIC determination. A
126 loopful from each of the test tubes was sub-cultured on Potato Dextrose agar. The plates were incubated
127 for 72 hours at $28 \pm 2^\circ\text{C}$. The MFC was determined as the least concentration that showed no visible
128 growth on the plate [23].

129

130 **2.10 Statistical analysis**

131 Experimental values were given as means \pm standard deviation (SD). Statistical significance of data were
132 analyzed at $P \leq 0.05$ (ANOVA) using statistical package for social sciences (SPSS, Armonk, NY, USA)
133 version 20.

134 **3. RESULTS AND DISCUSSION**

135 ~~Natural products not only provide valuable components but also an important source of bioactive~~
136 ~~compounds that provide lead information for developing useful synthetic compounds. Mushrooms contain~~
137 ~~a large number of biologically active components that impart health benefits and protection against~~
138 ~~degenerative diseases. They have been traditionally used in all over world for treatment of variety of~~
139 ~~chronic disease.~~ Antimicrobial activity of the crude extract of *Pleurotus squarrosulus* as well as
140 phytochemical characteristics were studied. Table 1 shows the result of the average MIC and MBC of the
141 ethanolic, methanolic and aqueous extracts of *P. squarrosulus* on the test organisms. The MIC of
142 ethanolic extract of *P. squarrosulus* varied between 15.63 and 31.25 mg/ml with MBC of 15.63 to 31.25
143 mg/ml. The MIC of methanolic extract of *P. squarrosulus* varied between 3.90 and 125 mg/ml with MBC of
144 7.81 to 125 mg/ml while the MIC of aqueous extract of *P. squarrosulus* varied between 31.25 and 62.50
145 mg/ml with MBC of 31.25 to 125 mg/ml.

146 Table 2 shows the result of the average MIC and MFC of the ethanolic, methanolic and aqueous extracts
147 of *P. squarrosulus* on test organisms. The MIC of ethanolic extract of *P. squarrosulus* showed 15.63
148 mg/ml for *C. albicans* and 125 mg/ml for *C. glabrata* with MFC of 31.25 and 125 mg/ml, respectively, the
149 MIC of methanolic extract of *P. squarrosulus* showed 250 mg/ml for *C. albicans* while *C. glabrata* showed
150 no activity with MFC of 250 mg/ml for *C. albicans* while the MIC of aqueous extract of *P. squarrosulus*
151 showed 7.81 mg/ml for *C. albicans* and 62.5 mg/ml for *C. glabrata* with MFC of 15.25 and 125 mg/ml,
152 respectively.

153 Table 3 shows the phytochemical analysis revealed the presence of bioactive compounds which were
154 present at varying levels. Saponins, protein and carbohydrate were detected in all the extracts while
155 glycosides, alkaloids, tannins and flavonoids were found in some.

156 Figure 1 shows the antimicrobial activity of *Pleurotus squarrosulus* methanol extract on the test
157 organisms. The mean inhibition zone diameter varied directly with increase in extract concentration. All
158 the tested strains of *E. coli*, *P. aeruginosa*, *B. cereus*, *S. aureus* and *C. albicans* were inhibited at different
159 concentrations of 500 mg/ml to 31.25 mg/ml whereas *C. glabrata* was not inhibited by the extract even at

160 the highest concentration of 500 mg/ml. However, inhibition of the antibacterial and antifungal control for
161 the test organisms were significantly higher ($p < 0.05$) than that of the extract.

162 Figure 2 presents the antimicrobial activity of *Pleurotus squarrosulus* ethanol extract on the test
163 organisms. The different test microorganisms showed varied susceptibility to the extract. All the test
164 organisms were well inhibited except *C. glabrata* that was only inhibited at concentrations of 500 mg/ml
165 and 250 mg/ml. However, inhibition of the antibacterial and antifungal control for the test organisms were
166 significantly higher ($p < 0.05$) than that of the extract.

167 Figure 3 shows the result obtained for the antimicrobial activity of *Pleurotus squarrosulus* aqueous
168 extract. *B. cereus*, *S. aureus*, *C. albicans* and *C. glabrata* were well inhibited by the extract. *E. coli* and *P.*
169 *aeruginosa* were not inhibited even at the highest concentration of 500 mg/ml. However, inhibition of the
170 antibacterial and antifungal control for the test organisms were significantly higher ($p < 0.05$) than that of
171 the extract.

172 The results indicated that extracts from mushroom ~~have antimicrobial properties~~ as reported by
173 Nwachukwu and Uzoeto [21]. Mushrooms produce various antiviral, antifungal compounds to survive in
174 the wild against competing or pathogenic agents [24, 25]. Also observed in this study is that there were
175 variations in the degree of antimicrobial activities of mushrooms. The sensitivity of isolates to the
176 mushroom extracts implies that intrinsic substance in the extracts is unknown to the microorganisms
177 which made it impossible for them to resist. The variations in the antimicrobial activities of *Pleurotus*
178 *squarrosulus* extracts may be due to the differences in their bioactive compositions or concentrations,
179 methods of extraction and mechanism of action of active ingredients [26]. The results of the present study
180 strengthened the outcomes of earlier works done by others that showed mushrooms produced a great
181 variety of antimicrobial agents. For instance, it is known that the extract from fruit bodies of several
182 *Lactarius* sp. [27, 28]; *Fomitopsis* sp. [29]; *Boletus* sp. [30]; *Cortinarius* sp. [31]; *Ganoderma lucidum*,
183 *Navesporus floccosa* and *Phellinus rimosus* [32]; *Pleurotus tuber-regium* [33]; *Amanita caesarae*,
184 *Armillaria mellea*, *Chroogomphus rutilus*, *Clavariadelphus truncates*, *Clitocybe geotropa*, *Ganoderma* sp.,
185 *Ganoderma carnosum*, *Hydnum repandum*, *Hygrophorus agathosmus*, *Lenzites betulina*, *Leucoagaricus*
186 *pudicus*, *Paxillus involutus*, *Polyporus arcularius*, *Rhizopogon roseo*, *Sarcodon imbricatus*, *Suillus*
187 *collitinus*, *Trametes versicolor*, *Tricholoma auratum*, *Tricholoma fracticum* [34]; *Lactarius deliciosus*,

188 *Sarcodon imbricatus* and *Tricholoma portentosum* [35]; *Russula delica* [36]; *Pleurotus eryngii* var. *ferulae*
 189 [37]; *Infundibulicybe geotropa*, *Lactarius controversus*, *Lactarius delicious* and *Phellinus hartigii* [38];
 190 *Lactarius indigo* [39] and *Stereum ostrea* [40] contain a wide range of antimicrobial activity.

191 4.CONCLUSION

192 This research has further illuminated the medicinal value of *Pleurotus squarrosulus* found in
 193 Umuahia North Local Government, Abia State Nigeria. From the present study, it can be
 194 concluded that *Pleurotus squarrosulus* possesses good quantities of compounds which have
 195 potent antimicrobial activity. Therefore, they have lots of potentials for use in the production of
 196 novel drugs and medicines, considering the lingering threat of multi-drug resistance.
 197 Furthermore, ~~clinical evaluation of mushrooms through in vivo based research is highly~~
 198 ~~recommended to achieve low cost, less side effect treatment and prevent recurrent infections.~~

199

200 Table 1: The MIC and MBC of crude extract of *Pleurotus squarrosulus*

Extract	Test organism	MIC (mg/ml)	MBC (mg/ml)
Ethanol	<i>B. cereus</i>	15.63	15.63
	<i>S.aureus</i>	15.63	31.25
	<i>P. aeruginosa</i>	15.63	31.25
	<i>E.coli</i>	31.25	31.25
Methanol	<i>B. cereus</i>	3.90	7.81
	<i>S.aureus</i>	31.25	62.5
	<i>P. aeruginosa</i>	62.5	62.5
	<i>E.coli</i>	125	125
	<i>B. cereus</i>	62.5	125
	<i>S.aureus</i>	31.25	31.25

Aqueous	<i>P. aeruginosa</i>	ND	ND
	<i>E.coli</i>	ND	ND

201 ND = NOT DETERMINED

202

203 Table 2: The MIC and MFC of the crude extract of *Pleurotus squarrosulus*

Extract	Test organism	MIC (mg/ml)	MFC (mg/ml)
Ethanol	<i>C.albicans</i>	15.63	31.25
	<i>C.glabata</i>	125	125
Methanol	<i>C.albicans</i>	250	250
	<i>C.glabata</i>	ND	ND
Aqueous	<i>C.albicans</i>	7.81	15.25
	<i>C.glabata</i>	62.5	125

204 ND = NOT DETERMINED

205 Table 3: PHYTOCHEMICAL ANALYSIS OF *PLEUROTUS SQUARROSULUS* IN DIFFERENT

206 SOLVENT


Solvents	Methanol	Ethanol	Aqueous
208 Saponin	++	+	+
209 Tannins	+	++	+

210	Flavonoid	+	++	+
211	Alkaloid	+	+	-
212	Proteins	++	+++	++
213	Glycosides	++	+++	-
214	Carbohydrates	++	++	++

215

Legend: - = not present, + = present in low concentration, ++ = moderate, +++ = present in high concentration,


216



217

218

219 Figure 1: The antimicrobial activity of *Pleurotus squarrosulus* methanol extract on the test
 220 organisms




221

222

223 Figure 2: The antimicrobial activity of *Pleurotus squarrosulus* ethanol extract on the test

224 organisms



225

226

227 Figure 3: The antimicrobial activity of *Pleurotus squarrosulus* aqueous extract on the test
 228 organisms

229

230

231

232 REFERENCES

- 233 1. Akpi UK, Odoh CK, Ideh EE, Adobu US. Antimicrobial Activity of *Lycoperdum perlatum* Whole fruit
234 body on Common Pathogenic Bacteria and Fungi. Afr J. Clin. Exper. Microbial. 2017; 18 (2): 79-85.
- 235 2. Subrata G, Gunjan B, Prakash P, Subhash CM, Krishnendu A. Antimicrobial Activities of Basidiocarps
236 of Wild Edible Mushrooms of West Bengal, India. Int. J. PharmTech Res. 2012; (4) 4: 1554-1560
- 237 3. Benedict RG, Brady LR. Antimicrobial activity of mushroom metabolites J. Pharmacol Sci. 1972; 61:
238 1820–1822.
- 239 4. Cohen R, Persky L, Hadar Y. Biotechnological applications and potential of Wood degrading
240 mushrooms of the genus *Pleurotus* J. App. Microbiol. Biotech. 2002; 58: 582-594.
- 241 5. Collins RA, Ng TB. Polysaccharopeptide from *Coriolus versicolor* has potential for use against human
242 immunodeficiency virus type-I infection Life Sci. 1997; 60:383-387.
- 243 6. Lindequist U, Niedermeyer THJ, Julich WD. The pharmacological potential of mushrooms. eCAM.
244 2005; 2: 285–299.
- 245 7. Smania EFA, Monache FD, Smania A, Cuneo YRA. Antifungal activity of sterols an triperpenes
246 isolated from *Ganoderma annulare*. Fitoterapia, 2003; 74: 375-377.
- 247 8. Vamanu E, Vamanu A, Pelinescu, D, Niță S, Rusu N, Popa N. Influence of the culture medium
248 composition on the exopolysaccharides synthesis by *Streptococcus* sp. IL5 strain, Acta Aliment. 2012;
249 41: 118–125.
- 250 9. Westh H, Zinn CS, Rosahi VT. An international multicenter study of antimicrobial consumption and
251 resistance in *Staphylococcus aureus* isolates from 15 hospitals in 14 countries Microb. Drug Resis.
252 2004; 10: 169 – 179.
- 253 10. Jagadish LK, Krishnan VV, Shenbhagaraman R, Kaviyaran V. Comparitive study on the
254 antioxidant, anticancer and antimicrobial property of *Agaricus bisporus* (J. E. Lange) Imbach before
255 and after boiling Afric. J. Biotech. 2009; 8 (4): 654-661.
- 256 11. Iwalokun, B.A., Usen, U.A., Otunba, A.A., Olukoya, D.K. Comparative phytochemical evaluation,
257 antimicrobial and antioxidant properties of *Pleurotus ostreatus* Afri. J. Biotechnol. 2007; 6:1732-1739.
- 258 12. Oso BA. Mushrooms in Yoruba mythology and medicinal practices. Ecol. Bot. 1997; 31 (31): 367 –
259 371.

- 260 13. Chapela IH, Lizon P. Fungi in stone age. Mycol. 1993; 7: 121 – 122
- 261 14. Tochikura TS, Nakashima H, Ohashi Y, Yamamoto H. Inhibition (in vitro) of replication and of the
262 cytopathic effect of human immunodeficiency virus by an extract of the culture medium of *Lentinus*
263 *edodes* mycelia. Med. Microbiol. Immunol. 1988; 177: 235 – 244.
- 264 15. Stamets P. Growing gourmet and medicinal mushrooms. Ten Speed Press, California. 2000
- 265 16. May JL, Wu KT, Chan PT, Wu YH, Peng JT. Flavour compounds in king oster mushroom (*Pleurotus*
266 *eryngii*). J. Agric Food Chem.1998; 46: 4587 – 4589.
- 267 17. Jonathan SG, Fasidi IO. Antimicrobial activities of two Nigerian edible macrofungi *Lycoperdon*
268 *pasilum* (Bat. Ex) and *Lycoperdon giganteum* (Pers.). Afri. J. Biomed. Res. 2003; 6: 85 – 90.
- 269 18. Udu-ibiam OE, Ogbu O, Nworie O, Ibiam UA, Agah MV, Nnachi, A. U. ~~et al.~~ Antimicrobial activities of
270 some selected edible mushrooms and spices against clinical isolates from Federal University
271 Teaching Hospital, Abakaliki (FETHA), Ebonyi State, Nigeria. Int. J. Sci. Technol. Res. 2014; 3 (5):
272 251- 255.
- 273 19. Eger G, Eden G, Wissig E. *Pleurotus ostreatus* – breeding potential of a new cultivated mushroom.
274 Theo. Appl. Gen. 1976; 47: 155 – 163
- 275 20. Beltran G, Miguel J, Estarron-Espinosa M, Ogura T. Volatile compounds secreted by the oyster
276 mushroom (*Pleurotus ostreatus*) and their antibacterial activities. J. Agric. Food Chem. 1997; 45
277 (10): 40 - 49
- 278 21. Nwachukwu,E, Uzoeto HO. Antimicrobial activity of some local mushrooms on pathogenic isolates. J.
279 Med. Plant Res. 2010; 4 (23): 2460 - 2465.
- 280 22. National Committee of Clinical Laboratory Standards. Performance Standards Antimicrobial Disc
281 Susceptibility Tests.Approved standard 5th edition. NCCLS document M2-AS, Villanova, PA, USA,
282 1998.
- 283 ~~23. Gutpe S. Medical Microbiology. New Delhi. Jaypee bros. med. Publishers and distributor Ltd, 2006~~
- 284 24. Sorimachi K, Ikehara Y, Maezato G. Inhibition of *Agaricus blazei* murill fractions of cytopathic effect
285 induced by western in vitro. Biosc. Biotechnol. biochem, 2001; 65 (7): 1645-1647.
- 286 25. Jang WJ, Hyung SW. Production of natural c9,t11 conjugated linoleic acid(c9, cLA) by submerged
287 liquid culture of mushrooms. Gyeongsang National University, South Korea, Jinju. 2004; 660-701

- 288 26. Bergendorf O, Sterner O. The sesquiterpenes of *Lactarius deliciosus* and *Lactarius deterrimus*.
289 Phytochem. 1998; 27: 97–100.
- 290 27. Anke T. Basidiomycetes: A source for new bioactive secondary metabolites Prog Ind Microbiol 1989,
291 27: 51–66.
- 292 28. Keller AC, Maillard MP, Hostettmann K. Antimicrobial steroids from the fungus *Fomitopsis pinicola*.
293 Phytochem. 1996; 41: 1041–1046.
- 294 29. Lee SJ, Yeo WH, Yun BS, Yoo ID. Isolation and sequence analysis of new peptaibol, Boletusin, from
295 *Boletus* spp. J. Pept. Sci. 1999; 5: 374–378.
- 296 30. Nicholas GM, Blunt JW, Munro MHG. Cortamidineoxide, a novel disulfide metabolite from the New
297 Zealand Basidiomycete (mushroom) *Cortinarius* species. J. Nat. Prod. 2001; 64: 341–344.
- 298 31. Sheena N, Ajith TA, Thomas M, Janardhanan KK. Antibacterial Activity of Three Macrofungi,
299 *Ganoderma lucidum*, *Navesporus floccosa* and *Phellinus rimosus* occurring in South India. Pharm.
300 Biol. 2003; 41: 564-567.
- 301 32. Ezeronye OU, Daba ASO, Onumajuru IAIC. Antibacterial Effect of Crude Polysaccharide Extracts
302 from Sclerotium and Fruit body (Sporophore) of *Pleurotus tuberregium* (Fried) Singer on Some
303 Clinical Isolates. Int. J. Mol. Med. Advance Sci. 2005; 1(3): 202-205.
- 304 33. Yamac M, Bilgili F. Antimicrobial Activities of Fruit Bodies and/or Mycelial Cultures of Some
305 Mushroom Isolates. Pharm. Biol. 2006; 44: 660-667.
- 306 34. Barros L, Calhella RC, Vaz JA, Ferreira ICFR, Baptista, P, Estevinho LM. Antimicrobial activity and
307 bioactive compounds of Portuguese wild edible mushroom methanolic extracts. Eur. Food Res.
308 Technol. 2006; 225: 151-156.
- 309 35. Turkoglu A, Kivarak I, Mercan N, Duru ME, Gezer K, Turkoglu H. Antioxodant and antimicrobial
310 activities of *Morchella conica* Pers. Afric. J. Biotechnol. 2006; 5:1146-1150.
- 311 36. Akyuz M, Kirbag S. Antimicrobial activity of *Pleurotus eryngii* var. *ferulae* grown on various agro-
312 wastes. Eur. J. Biosc. 2009; 3:58-63.
- 313 37. Altuner EM, Akata I. Antimicrobial activity of some macrofungi extracts. SAU Fen Bilimleri Dergisi Cilt.
314 2010; 14, 45-49.

- 315 38. Ochoa-Zarzosa A, Vazquez-Garciduenas SM, Robinson-Fuentes VA, Vazquez- Marrufo G.
316 Antimicrobial and cytotoxic activity from basidiocarp extracts of the edible mushroom *Lactarius indigo*
317 (Schw.) Fr. (Russulaceae), *Afric. J. Pharm. Pharmacol.* 2011; 5, 281-288.
- 318 39. Praveen K, Usha KY, Naveen M, Rajasekhar BR. Antibacterial Activity of a Mushroom-*Stereum*
319 *ostrea*. *J. Biol. Agric. Healthcare*, 2012; 2: 1-5.
- 320 40. Sivaprakasam E, Kavitha D, Balakumar R, Sridhar S, Suresh K. Antimicrobial activity of whole
321 fruiting bodies of *Trametes hirsute* (Wulf. Fr.) Pil. against some common Pathogenic Bacteria and
322 Fungus. *Int J. Pharm. Sci. Drug Res* 2011; 3(3): 219-221