

Original research papers**Alteration of sex hormone and semen parameters in adult males with subclinical hypothyroidism****ABSTRACT**

Aim: To investigate the correlation between subclinical hypothyroidism (SCH) with serum testosterone levels and semen parameters (sperm count, total motility and morphology) in men seeking medical care for sexual dysfunction and infertility.

Study design: Cross-sectional study.

Place and Duration of Study: Department of Endocrinology, Gauhati Medical College and hospital, Guwahati, India. Study done from November 2015 to December 2016.

Methodology: The patients in study were grouped into two groups: Group I – Men with subclinical hypothyroidism (n=35) and group II – Euthyroid males serving as controls (n=27). The medical history, clinical examination, semen analysis, total thyroxin(T4), total triiodothyronine(T3), thyroid-stimulating hormone(TSH), Leutinising Hormone (LH), Follicular stimulating hormone (FSH), Total testosterone (T) and prolactin (PRL) were obtained. Patients with diabetes mellitus, hypertension, chronic diseases, any testicular or pituitary diseases and prior chemo-radiotherapy were excluded from study.

Results: The ages of the patients ranged between 24 and 42 years with mean age 31.57 years. The cases had significantly lower levels of mean serum total testosterone and free T4 and higher serum prolactin levels compared to controls ($P < .01$). Linear regression analysis showed TSH levels independently associated with low testosterone levels. Also men with subclinical hypothyroidism had lower total sperm motility with preserved sperm count and morphology.

Conclusion: We found that SCH is associated with reduction in testosterone levels and total sperm motility. Our study data conclude that SCH may be a contributing factor for hypoandrogenemia and sexual dysfunction in men.

Keywords: Subclinical hypothyroidism; testosterone; sperm count; total sperm motility.

1. INTRODUCTION

Male reproductive function is orchestrated by the hypothalamo-hypophyseal testicular axis. In addition it has been seen that thyroid hormones exerts a modulatory effect on this axis. Thus thyroid hormones seem to play a major role in male sexual and spermatogenic function. Primary hypothyroidism has been associated with low levels of both total and free testosterone in men [1]. The worldwide prevalence of subclinical hypothyroidism is found to be between 4 and 10% [2, 3]. The prevalence of subclinical hypothyroidism in india is about 9.4%. In women, the prevalence was higher (11.4%), when compared with men (6.2%) [4]. But links between subclinical hypothyroidism and male sexual dysfunction and infertility are less evident. Till date there has been no concrete study that has showcased this relationship. Our study aims to investigate the association between subclinical hypothyroidism with total testosterone levels and semen parameters (sperm count, total motility and morphology) in men seeking medical care for sexual dysfunction and infertility in tertiary care centre.

26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58

2. MATERIAL AND METHODS

The study is a cross sectional study done in males of age between 20-45 years, attending Endocrinology department of Gauhati Medical College and Hospital, Assam, India from November 2015 to December 2016 for sexual dysfunction or infertility. The patients in the study were grouped into two: Group I - Men who were subclinical hypothyroidism (n=35) and group II- Men who were normal euthyroid controls (n=27). Patients with diabetes mellitus, hypertension, any chronic disease, testicular or pituitary disorder and prior chemo radiotherapy were excluded from study. Patient who fulfilled the study criteria underwent a detailed medical history and clinical examination. Serum samples were collected for free thyroxine (FT4), total triiodothyronine (T3), thyroid-stimulating hormone (TSH), leutinising hormone (LH), follicular stimulating hormone (FSH), total testosterone (T) and prolactin (PRL) estimation. Blood samples were collected in the morning after subjects had fasted for at least 8 hours. The samples were immediately centrifuged and the samples were aliquoted into 2-ml tubes .The serum was stored at -80°C until assayed. Hormonal assay was done by enzyme-enhanced chemiluminescent analyzer (IMMULITE 1000). Single semen analysis was also done in pathology department of our medical college with fresh samples collected within 30 minutes of ejaculation. World Health Organization 2010 reference values for human semen characteristics were considered in the study [5]. The study was performed according to the guidelines of the Ethics Committee of our institute with informed consent. Statistical analysis was performed by the use of SAS 9.3. Pearson correlation between TSH levels and LH, FSH, PRL, total testosterone levels and semen parameters were estimated. Linear regression model analysis was used to find independent association between TSH and total testosterone levels .Multiple regression analysis model was used to find the simultaneous effect of thyroid hormones (TSH, Free T4 and Total T3) on the serum total testosterone levels. Statistical significance was set at the 0.05 level.

3. results and discussion

3.1 Baseline Characteristics

The ages of the patients ranged between 24 and 42 years with mean age 31.57 years (SD \pm 5.2). Table 1 shows the baseline characteristics of the cases and control.


Characteristics	Case (n=35)	Control (n=27)	P value
Age(years)	31.5714286	35.5925926	0.0037
Body mass index (kg/m ²)	24.1357143	24.6518519	0.1054
Testicular volume (ml)	20.8571429	18.5185185	0.0104
TSH (mIU/L)	9.0989143	2.3629630	<.0001
Total Testosterone (ng /dl)	414.2265714	683.2000000	0.0003
Prolactin (ng /ml)	16.2294286	10.7777778	<.0001
LH (mIU/mL)	4.4051429	3.9307407	0.0403
FSH (mIU/mL)	4.7020000	4.6385185	0.215
Sperm count (million per ml)	50.8000000	52.4444444	0.7606
Semen volume (ml)	3.1657143	2.8703704	0.3125
Total sperm motility (in %)	53.1428571	82.8148148	<.0001
Duration of symptoms(years)	4.9714286	3.0740741	0.0014
Free T4 (ng /dl)	1.1731429	1.3237037	<.0001
Total T3 (nmol/L)	1.9234286	1.8400000	0.1835

59

60 As seen in table 1, there were no differences between subclinical hypothyroid cases and euthyroid control
 61 in terms of mean BMI, mean FSH, mean sperm count, mean semen volume and total T3. However, the
 62 cases had significant lower level of mean serum total testosterone levels (414.22 ± 225.48 in cases vs.
 63 683.20 ± 328.29 in controls) and free T4 (1.17 ± 0.14 in cases vs. in 1.32 ± 0.07 controls) but a higher
 64 level of serum prolactin (16.22 ± 4.70 in cases vs. in 10.77 ± 2.48 controls) in comparison to control
 65 group. LH levels were also significantly different between the two groups (4.40 ± 0.87 in cases vs. $3.93 \pm$
 66 0.89 controls). Among the semen parameters, the cases had a remarkable lower total sperm motility
 67 compared to controls groups (53.14 ± 25.83 in cases vs. 82.81 ± 16.83 in controls).

68
 69 Figure1 (A-D) shows the bar diagram comparing the mean values of various hormones (TSH, Prolactin,
 70 LH and testosterone)

71
 72



73
 74
 75
 76
 77
 78
 79
 80
 81
 82
 83

Figure 2 (E&F) shows the bar diagram comparing the mean values semen parameters among the cases and control groups. Sperm morphology was normal in both the cases and control group as per the Kruger's criteria.




Figure 2 : Each Bar represents the mean \pm SEM (standard error of mean) of sperm count and total sperm motility of the cases ($n= 35$) and the controls ($n= 27$).

84
85
86
87
88
89
90
91
92
93
94

3.2 Correlations between TSH and other hormones and semen parameters

Among the subclinical hypothyroid patients, a significant negative correlation was observed between serum TSH and total testosterone levels ($r = -0.75, P < .01$). Negative correlation was also found between total testosterone levels and serum prolactin ($r = -0.69, P < .01$). Figure 3 (G&H) demonstrates these relationship. Similarly a negative correlation was also seen among the cases between serum TSH and total sperm motility ($r = -0.73, P < .01$). No correlations were found between TSH levels and FSH, LH or sperm count.





Figure 3 : Fitted line plot showing negative linear relationship between TSH levels and testosterone levels (G) and testosterone levels and prolactin (H) among the subclinical hypothyroid males ($n=35$). The solid line shows the mean testosterone levels among the subclinical hypothyroid males and the open circles shows the distribution of TSH levels (G) and prolactin levels (H) of each cases .

95
96
97
98
99
100
101
102

3.3 Correlations between testosterone and Free T4 and total T3

A significant positive correlation was found between total testosterone levels and free T4 and Total T3 levels ($r = 0.54, P < .01$; $r = 0.65, P < .01$). Figure 4 (I&J) demonstrates these relationship.



103
104
105
106
107
108
109
110
111
112
113
114
115

Linear regression model analysis showed significant association between TSH and total testosterone levels ($P < .01$). Multiple regression model analysis was done to assess the simultaneous effect of thyroid hormones (TSH, Free T4, total T3) on serum testosterone. The model showed TSH and free T4 contributed significantly to low total testosterone level. Table 2 shows multiple regression analysis.

Table 2: Multiple regression analysis showing the relationship between serum testosterone and thyroid hormones.

Hormones	B – coefficient	P Value
TSH	-100.41507	0.0001
Free T4	-920.19167	0.0122
Total T3	147.03873	0.3160

116
117
118
119
120
121
122
123
124
125
126
127
128
129

4. DISCUSSION

In the current study we found a significant reduction in serum total testosterone levels in subclinical hypothyroid males compared to euthyroid control. It was also seen that subclinical hypothyroid patients the free T4 levels were significantly lower compared to the control group. The levels if T3 is dependent on the availability of T4 hormone. We could find a positive correlation between total testosterone levels and total T3 concentration ($r = 0.65$, $P < .01$) among the subclinical hypothyroid males. This significant correlation explains the role of total T3 in the production of testosterone. The role of T3 hormone in testosterone synthesis has been demonstrated earlier in Leydig cells of rat and goat [6, 7]. These finding supports the modulatory effects of thyroid hormones in maintaining the normal testosterone levels and male sexual function. However, it has to be understood that serum T3 concentration in subclinically

130 hypothyroid cases was along the lower side but within the normal reference range in this study. Secondly
131 linear regression model analysis showed a significant independent association between TSH and total
132 testosterone levels among subclinical hypothyroid males. The probable explanation lies in the fact that
133 thyroid-stimulating hormone receptors have also been demonstrated in human testis [8]. It is postulated
134 by some authors that TSH may have a cardinal role in regulating testosterone synthesis in leydig cells.
135 However, till date there is no concrete evidence supporting the above. Multiple regression model analysis
136 was done and showed the effects of TSH and thyroid hormones together in generation of testosterone in
137 males. The analysis showed that TSH (beta coefficient: - 100.4) along with thyroid hormones
138 simultaneously has a significant role in testosterone production.
139

140 The rise of LH levels seen in subclinical hypothyroid males has been a significant finding compared to the
141 euthyroid control. We postulate that this may be due to loss of feedback mechanism by low levels of
142 testosterone. It should be understood that LH or FSH feedbacks are under the control of free testosterone
143 levels. Our study has been based on total testosterone and hence further confirmation is needed.
144 Moreover the rise of LH and FSH levels in overt hypothyroidism itself is controversial [9,10]. We also
145 found a significant elevation of serum prolactin levels among the subclinical hypothyroid patients. This
146 findings was similar to study done by Kumar et al (2006) [11]. Thought multiple mechanism have been
147 explained as the cause of hypoandrogenemia in elevated levels of prolactin, the degree of
148 hyperprolactinemia required to cause low levels of testosterone has been a consistent finding with overt
149 hypothyroids males. The mechanisms illustrated in overt hypothyroid males include suppression of 17
150 alphahydroxylase (a key enzyme in conversions of progesterone to testosterone in Leydig cells) [12] and
151 decreasing the binding affinity of LH to its receptors (seen on a murine tumour cell line, MA-10 cells) [13]
152 by high concentration of prolactin. In our study not only a higher levels of serum prolactin were found
153 among the subclinical hypothyroid males but also a significant negative correlation found between total
154 testosterone and serum prolactin levels, ($r = -0.69$, $P < .01$) which probably explains that elevated
155 prolactin as the cause of low levels of total testosterone among the cases. Moreover Kumar et al (2006)
156 in his study has demonstrated the reduced availability of progesterone (precursor for the synthesis of
157 testosterone) in subclinical hypothyroid males. Summating all the above data we may conclude that
158 hyperprolactinemia in subclinical hypothyroid males may lead to hypoandrogenemia a picture similar to
159 overt hypothyroidism.
160

161 .
162 Finally we also learned the relationship between subclinical hypothyroidism and semen parameters.
163 Earlier studies have shown that subclinical hypothyroidism had no impact on semen parameter [14]. The
164 only semen parameter which was affected in our study was the total motility with not affecting sperm
165 count and morphology. TSH levels correlated with sperm total motility ($r = -0.73$, $P < .01$). In overt
166 hypothyroidism it has seen that low levels of thyroid hormone may affect forward progressive motility.
167 Mechanism postulated is that thyroid hormones stimulate cellular oxygen consumption [15] and increases
168 mitochondrial number over the middle and tail piece of sperm [16] and thus promoting sperm motility. In
169 our study due to limited data we were only able to compute total sperm motility.
170

171 172 **4. CONCLUSION**

173
174 In conclusion from our study it was seen that subclinical hypothyroidism may lead to low levels of total
175 testosterone levels. Spectrum of male sexual dysfunction may be a presentation of subclinical
176 hypothyroidism. Till now there is no consensus to treat subclinical hypothyroidism in male. Further studies
177 may be needed to confirm the need for supplementing thyroid hormones to this class of patients. The
178 genesis of infertility in subclinical hypothyroid males remains a domain of interest. Thought the study has
179 concluded the effects on total sperm motility without affecting sperm count and morphology, it is difficult to
180 interpret infertility in this context. The major drawback of our study was that we were unable to sub
181 classify the types of motility defects due to limited data.
182

183 **CONSENT (WHERE EVER APPLICABLE)**

184

185 We declare that 'written informed consent was obtained from the patient (or other approved parties) for
186 publication of this case report and accompanying images. A copy of the written consent is available for
187 review by the Editorial office/Chief Editor/Editorial Board members of this journal.'

188

189

190 **ETHICAL APPROVAL**

191

192 We declare that we have obtained all necessary ethical approval from our Institutional ethical Committee.

193

194 **REFERENCES**

195

196 1,Kumar A, Shekhar S, Dhole B. Thyroid and male reproduction Indian Journal Endocrinology
197 Metabolism 2014;18:23-31

198

199 2,Hollowell JG, Staehling NW, Flanders WD, Hannon WH, Gunter EW, Spencer CA, *et al.* Serum TSH,
200 T(4), and thyroid antibodies in the United States population (1988 to 1994): National Health and Nutrition
201 Examination Survey (NHANES III). J Clin Endocrinol Metab ;87:489–99

202

203 3,Canaris GJ, Manowitz NR, Mayor G, Ridgway EC. The Colorado thyroid disease prevalence study.
204 Arch Intern Med 2000;160:526–34

205

206 4, Unnikrishnan AG, Menon UV. Thyroid disorders in India: An epidemiological perspective. Indian J
207 Endocr Metab 2011;15:S78-81.

208

209 5, Cooper, TG et al. WHO reference values for human semen characteristics
210 Human. Reproduction Update 2010;16(5):559

211

212 6, Oppenheimer, J. H., Schwartz, H. L. & Surks, M. I. (1974) Tissue differences in the concentration of
213 triiodothyronine nuclear binding sites in the rat: liver, kidney, pituitary, heart, brain, spleen and testis.
214 Endocrinology 95, 897–903.

215

216 7, Jana, N. R. & Battacharya, S. (1994) Binding of thyroid hormone to the goat testicular Leydig cell
217 induces the generation of a proteinaceous factor which stimulates androgen release. Journal of
218 Endocrinology 143, 549–556.

219

220 8, Trokoudes, K. M., Sugeno, A., Hazani, E., Row, V. V. & Volpe, R. (1979) Thyroid-stimulating
221 hormone (TSH) binding to extrathyroidal human tissues: TSH and thyroid-stimulating immunoglobulin
222 effects on adenosine 3, 5 -monophosphate in testicular and adrenal tissues. Journal of Clinical
223 Endocrinology and Metabolism 48, 919–923

224

225 9, Jaya Kumar B, Khurana ML, Ammini AC, Karmarkar MG, Ahuja MM. Reproductive endocrine functions
226 in men with primary hypothyroidism: effect of thyroxine replacement. Horm Res 1990;34:215-8.

227

228 10. Donnelly P, White C. Testicular dysfunction in men with primary hypothyroidism; Reversal of
229 hypogonadotrophic hypogonadism with replacement thyroxine. Clin Endocrinol 2000;52:197-201.

230

231 11, Kumar A, Chaturvedi PK, Mohanty BP. Hypoandrogenaemia is associated with subclinical
232 hypothyroidism in men Int J Andro 2006;30:14-20.

233

234 12, Welsh, T. H., Kasson, B. G. & Hsueh, A. J. W. (1986) Direct biphasic modulation of gonadotropin-
235 stimulated testicular androgen biosynthesis by prolactin. Biology of Reproduction 34, 796–804.

236

- 237 13, Weiss-Messer, E., Ber, R. & Barkey, R. J. (1996) Prolactin and MA-10 Leydig cell steroidogenesis:
238 biphasic effects of prolactin and signal transduction. *Endocrinology* 137, 5509–5518.
239
- 240 14, Trummer H, Ramschak-Schwarzer S, Haas J, Habermann H, Pummer K, Leb, G. Thyroid hormones
241 and thyroid antibodies in infertile males. *Fertility and Sterility* 2001;76:254–57
242
- 243 15, Lei J, Nowbar S, Mariash CN, Ingbar DH. Thyroid hormone stimulates Na-K-ATPase activity and its
244 plasma membrane insertion in rat alveolar epithelial cells. *Am J Physiol Lung Cell Mol Physiology*
245 2003;285:762-72.
246
- 247 16, Wrutniak-Cabello C, Casas F, Cabello G. Thyroid hormone action in mitochondria. *J Mol Endocrinol*
248 2001;26:67-77.
249
- 250 **APPENDIX**
251