

# Alteration of sex hormone and semen parameters in adult males with subclinical hypothyroidism

**Dipti Sarma<sup>1</sup>, Uma Kaimal Saikia<sup>1</sup>, Darvin Vamadevan Das<sup>1</sup>**

<sup>1</sup> Department of Endocrinology, Gauhati Medical College and Hospital, Guwahati, India.

**Aim:** To investigate the correlation between subclinical hypothyroidism (SCH) with serum testosterone levels and semen parameters (sperm count, total motility and morphology) in men seeking medical care for sexual dysfunction and infertility.

**Study design:** Cross-sectional study.

**Place and Duration of Study:** Department of Endocrinology, Gauhati Medical College and hospital, Guwahati, India. Study done from November 2015 to December 2016.

**Methodology:** The patients in study were grouped into two groups: Group I – Men with subclinical hypothyroidism (n=35) and group II – Euthyroid males serving as controls (n=27). The medical history, clinical examination, semen analysis, total thyroxin(T4), total triiodothyronine(T3), thyroid-stimulating hormone(TSH), Leutinising Hormone (LH), Follicular stimulating hormone (FSH), Total testosterone (T) and prolactin (PRL) were obtained. Patients with diabetes mellitus, hypertension, chronic diseases, any testicular or pituitary diseases and prior chemo-radiotherapy were excluded from study.

**Results:** The ages of the patients ranged between 24 and 42 years with mean age 31.57 years. The cases had significantly lower levels of mean serum total testosterone and free T4 and higher serum prolactin levels compared to controls ( $P < .01$ ). Linear regression analysis showed TSH levels independently associated with low testosterone levels. Also men with subclinical hypothyroidism had lower total sperm motility with preserved sperm count and morphology.

**Conclusion:** We found that SCH is associated with reduction in testosterone levels and total sperm motility. Our study data conclude that SCH may be a contributing factor for hypoandrogenemia and sexual dysfunction in men.

**Keywords:** Subclinical hypothyroidism; testosterone; sperm count; total sperm motility.

## 1. INTRODUCTION

Male reproductive function is orchestrated by the hypothalamo-hypophyseal testicular axis. In addition it has been seen that thyroid hormones exerts a modulatory effect on this axis. Thus thyroid hormones seem to play a major role in male sexual and spermatogenic function. Primary hypothyroidism has been associated with low levels of both total and free testosterone in men [1]. The worldwide prevalence of subclinical hypothyroidism is found to be between 4 and 10% [2, 3]. The prevalence of subclinical hypothyroidism in india is about 9.4%. In women, the prevalence was higher (11.4%), when compared with men (6.2%) [4]. But links between subclinical hypothyroidism and male sexual dysfunction and infertility are less evident. Till date there has been no concrete study that has showcased this relationship. Our study aims to investigate the association between subclinical hypothyroidism with total testosterone levels and semen parameters (sperm count, total motility and morphology) in men seeking medical care for sexual dysfunction and infertility in tertiary care centre.

## 27 2. MATERIAL AND METHODS

28

29 The study is a cross sectional study done in males of age between 20-45 years, attending Endocrinology  
30 department of Gauhati Medical College and Hospital, Assam, India from November 2015 to December  
31 2016 for sexual dysfunction or infertility. The patients in the study were grouped into two: Group I - Men  
32 who were subclinical hypothyroidism (n=35) and group II- Men who were normal euthyroid controls  
33 (n=27). The most common sexual dysfunction among the subclinical hypothyroid males was erectile  
34 dysfunction (26 out of 35) and premature ejaculation (9 out of 35). On the other hand all men among the  
35 euthyroid control had presented with only premature ejaculation as the sexual dysfunction. Patients with  
36 diabetes mellitus, hypertension, any chronic disease, testicular or pituitary disorder and prior chemo  
37 radiotherapy were excluded from study. Patient who fulfilled the study criteria underwent a detailed  
38 medical history and clinical examination. Serum samples were collected for free thyroxine (FT4), total  
39 triiodothyronine (T3), thyroid-stimulating hormone (TSH), leutinising hormone (LH), follicular stimulating  
40 hormone (FSH), total testosterone (T) and prolactin (PRL) estimation. Blood samples were collected in  
41 the morning after subjects had fasted for at least 8 hours. The samples were immediately centrifuged and  
42 the samples were aliquoted into 2-ml tubes .The serum was stored at -80 °C until assayed. Hormonal  
43 assay was done by enzyme-enhanced chemiluminescent analyzer (IMMULITE 1000). Single semen  
44 analysis was also done in pathology department of our medical college with fresh samples collected  
45 within 30 minutes of ejaculation. World Health Organization 2010 reference values for human semen  
46 characteristics were considered in the study [5]. The study was performed according to the guidelines of  
47 the Ethics Committee of our institute with informed consent. Statistical analysis was performed by the use  
48 of SAS 9.3. Pearson correlation between TSH levels and LH, FSH, PRL, total testosterone levels and  
49 semen parameters were estimated. Linear regression model analysis was used to find independent  
50 association between TSH and total testosterone levels .Multiple regression analysis model was used to  
51 find the simultaneous effect of thyroid hormones (TSH, Free T4 and Total T3) on the serum total  
52 testosterone levels. Statistical significance was set at the 0.05 level.

53

## 54 3. Results

55

### 56 3.1 Baseline Characteristics

57 Table 1 shows the baseline characteristics of the cases and control.

58


59

Characteristics	Case (n=35)	Control (n=27)	P value
Age( years)	31.5714286	35.5925926	0.0037
Body mass index (kg/m <sup>2</sup> )	24.1357143	24.6518519	0.1054
Testicular volume (ml)	20.8571429	18.5185185	0.0104
TSH ( mIU/L )	9.0989143	2.3629630	<.0001
Total Testosterone (ng /dl )	414.2265714	683.2000000	0.0003
Prolactin (ng /ml )	16.2294286	10.7777778	<.0001
LH (mIU/mL)	4.4051429	3.9307407	0.0403
FSH (mIU/mL)	4.7020000	4.6385185	0.215
Sperm count (million per ml)	50.8000000	52.4444444	0.7606
Semen volume (ml)	3.1657143	2.8703704	0.3125
Total sperm motility (in %)	53.1428571	82.8148148	<.0001
Duration of symptoms(years)	4.9714286	3.0740741	0.0014
Free T4 (ng /dl)	1.1731429	1.3237037	<.0001
Total T3 (nmol/L)	1.9234286	1.8400000	0.1835

60

61 As seen in table 1, the ages of the patients ranged between 24 and 42 years with mean age 31.57 years  
 62 (SD  $\pm$  5.2). There were no differences between subclinical hypothyroid cases and euthyroid control in  
 63 terms of mean BMI, mean FSH, mean sperm count, mean semen volume and total T3. However, the  
 64 cases had significant lower level of mean serum total testosterone levels (414.22  $\pm$  225.48 in cases vs.  
 65 683.20  $\pm$  328.29 in controls) and free T4 (1.17  $\pm$  0.14 in cases vs. in 1.32  $\pm$  0.07 controls ) but a higher  
 66 level of serum prolactin (16.22  $\pm$  4.70 in cases vs. in 10.77  $\pm$  2.48 controls ) in comparison to control  
 67 group. LH levels were also significantly different between the two groups (4.40  $\pm$  0.87 in cases vs. 3.93  $\pm$   
 68 0.89 controls). Among the semen parameters, the cases had a remarkable lower total sperm motility  
 69 compared to controls groups (53.14  $\pm$  25.83 in cases vs. 82.81  $\pm$  16.83 in controls).

70  
 71 Figure1 (A-D) shows the bar diagram comparing the mean values of various hormones (TSH, Prolactin,  
 72 LH and testosterone)  
 73  
 74



75  
 76  
 77  
 78  
 79 Figure 2 (E&F) shows the bar diagram comparing the mean values semen parameters among the cases  
 80 and control groups. Sperm morphology was normal in both the cases and control group as per the  
 81 Kruger's criteria.  
 82  
 83  
 84  
 85




Figure 2 : Each Bar represents the mean  $\pm$  SEM (standard error of mean) of sperm count and total sperm motility of the cases (n= 35) and the controls (n= 27).

86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96

### 3.2 Correlations between TSH and other hormones and semen parameters

Among the subclinical hypothyroid patients, a significant negative correlation was observed between serum TSH and total testosterone levels ( $r = -0.75, P < .01$ ). Negative correlation was also found between total testosterone levels and serum prolactin ( $r = -0.69, P < .01$ ). Figure 3 (G&H) demonstrates these relationship. Similarly a negative correlation was also seen among the cases between serum TSH and total sperm motility ( $r = -0.73, P < .01$ ). No correlations were found between TSH levels and FSH, LH or sperm count.





Figure 3 : Fitted line plot showing negative linear relationship between TSH levels and testosterone levels (G) and testosterone levels and prolactin (H) among the subclinical hypothyroid males (n=35). The solid line shows the mean testosterone levels among the subclinical hypothyroid males and the open circles shows the distribution of TSH levels (G) and prolactin levels (H) of each cases .

97  
98  
99  
100  
101  
102  
103  
104

### 3.3 Correlations between testosterone and Free T4 and total T3

A significant positive correlation was found between total testosterone levels and free T4 and Total T3 levels ( $r = 0.54, P < .01$ ;  $r = 0.65, P < .01$ ). Figure 4 (I&J) demonstrate this relationship.



105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117

Linear regression model analysis showed significant association between TSH and total testosterone levels ( $P < .01$ ). Multiple regression model analysis was done to assess the simultaneous effect of thyroid hormones (TSH, Free T4, total T3) on serum testosterone. The model showed TSH and free T4 contributed significantly to low total testosterone level. Table 2 shows multiple regression analysis.

Table 2: Multiple regression analysis showing the relationship between serum testosterone and thyroid hormones.

Hormones	B – coefficient	P Value
TSH	-100.41507	0.0001
Free T4	-920.19167	0.0122
Total T3	147.03873	0.3160

118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131

#### 4. DISCUSSION

In the current study we found a significant reduction in serum total testosterone levels in subclinical hypothyroid males compared to euthyroid control. It was also seen that subclinical hypothyroid patients the free T4 levels were significantly lower compared to the control group. The levels if T3 is dependent on the availability of T4 hormone. We could find a positive correlation between total testosterone levels and total T3 concentration ( $r = 0.65$ ,  $P < .01$ ) among the subclinical hypothyroid males. This significant correlation explains the role of total T3 in the production of testosterone. The role of T3 hormone in testosterone synthesis has been demonstrated earlier in Leydig cells of rat and goat [6, 7]. These finding supports the modulatory effects of thyroid hormones in maintaining the normal testosterone levels and male sexual function. However, it has to be understood that serum T3 concentration in subclinically

132 hypothyroid cases was along the lower side but within the normal reference range in this study. Secondly  
133 linear regression model analysis showed a significant independent association between TSH and total  
134 testosterone levels among subclinical hypothyroid males. The probable explanation lies in the fact that  
135 thyroid-stimulating hormone receptors have also been demonstrated in human testis [8]. It is postulated  
136 by some authors that TSH may have a cardinal role in regulating testosterone synthesis in leydig cells.  
137 However, till date there is no concrete evidence supporting the above. It has to be understood that a  
138 negative relationship was found between TSH levels and total testosterone levels. Multiple regression  
139 model analysis was done and showed the effects of TSH and thyroid hormones together in generation of  
140 testosterone in males. The analysis showed that TSH (beta coefficient: - 100.4) along with thyroid  
141 hormones simultaneously has a significant role in testosterone production.

142  
143 The rise of LH levels seen in subclinical hypothyroid males has been a significant finding compared to the  
144 euthyroid control. We postulate that this may be due to loss of feedback mechanism by low levels of  
145 testosterone. It should be understood that LH or FSH feedbacks are under the control of free testosterone  
146 levels. Our study has been based on total testosterone and hence further confirmation is needed.  
147 Moreover the rise of LH and FSH levels in overt hypothyroidism itself is controversial [9,10]. We also  
148 found a significant elevation of serum prolactin levels among the subclinical hypothyroid patients. This  
149 findings was similar to study done by Kumar et al (2006) [11]. Thought multiple mechanism have been  
150 explained as the cause of hypoandrogenemia in elevated levels of prolactin, the degree of  
151 hyperprolactinemia required to cause low levels of testosterone has been a consistent finding with overt  
152 hypothyroids males. The mechanisms illustrated in overt hypothyroid males include suppression of 17  
153 alphahydroxylase (a key enzyme in conversions of progesterone to testosterone in Leydig cells) [12] and  
154 decreasing the binding affinity of LH to its receptors (seen on a murine tumour cell line, MA-10 cells) [13]  
155 by high concentration of prolactin. In our study not only a higher levels of serum prolactin were found  
156 among the subclinical hypothyroid males but also a significant negative correlation found between total  
157 testosterone and serum prolactin levels, ( $r = -0.69$ ,  $P < .01$ ) which probably explains that elevated  
158 prolactin as the cause of low levels of total testosterone among the cases. Moreover Kumar et al (2006)  
159 in his study has demonstrated the reduced availability of progesterone (precursor for the synthesis of  
160 testosterone) in subclinical hypothyroid males. Summating all the above data we may conclude that  
161 hyperprolactinemia in subclinical hypothyroid males may lead to hypoandrogenemia a picture similar to  
162 overt hypothyroidism.

163  
164  
165 Finally we also learned the relationship between subclinical hypothyroidism and semen parameters.  
166 Earlier studies have shown that subclinical hypothyroidism had no impact on semen parameter [14]. The  
167 only semen parameter which was affected in our study was the total motility without affecting sperm count  
168 and morphology. TSH levels correlated with sperm total motility ( $r = -0.73$ ,  $P < .01$ ). In overt  
169 hypothyroidism it has seen that low levels of thyroid hormone may affect forward progressive motility.  
170 Mechanism postulated is that thyroid hormones stimulate cellular oxygen consumption [15] and increases  
171 mitochondrial number over the middle and tail piece of sperm [16] and thus promoting sperm motility. **This**  
172 **probably is a major drawback in our study as we were unable to sub classify the types of motility defects**  
173 **(progressive motility, non-progressive motility or immotile sperms) among the subclinical hypothyroid**  
174 **males due to limited data. Thought motility defects have been demonstrated among hypothyroid men in**  
175 **earlier studies, motility defects among subclinical hypothyroid males have to be reconsidered. Moreover**  
176 **the mean total motility among the subclinical hypothyroid males fell into a mean value of 53%, which was**  
177 **within the lower reference as per the World Health Organization reference values for human semen**  
178 **characteristics [5] .The small sample size among the cases and control may be another drawback in the**  
179 **study. We also couldn't estimate free testosterone levels among the patients which may be needed to**  
180 **explain the rising levels of gonadotropin levels among the subclinical hypothyroid males.**

## 183 5. CONCLUSION

184  
185 In conclusion from our study it was seen that subclinical hypothyroidism may lead to low levels of total  
186 testosterone levels. Spectrum of male sexual dysfunction may be a presentation of subclinical

187 hypothyroidism. Till now there is no consensus to treat subclinical hypothyroidism in male. Further studies  
188 may be needed to confirm the need for supplementing thyroid hormones to this class of patients. The  
189 genesis of infertility in subclinical hypothyroid males remains a domain of interest. Thought the study has  
190 concluded the effects of subclinical hypothyroidism on total sperm motility more studies may be required  
191 to confirm this findings.

192

## 193 **ACKNOWLEDGEMENTS**

194

195 We would like to thank both the clinical laboratory team and the outpatient clinic team in Gauhati medical  
196 college for their great help to finalize this work

197

198

## 199 **COMPETING INTERESTS**

200

201 We declare that no competing interests exist.

202

## 203 **AUTHORS' CONTRIBUTIONS**

204

205 Dipti Sarma, Uma kaimal saikia and Darvin Das has designed the study, performed the statistical  
206 analysis, wrote the protocol, and wrote the first draft of the manuscript. All authors read and approved the  
207 final manuscript."

208

## 209 **CONSENT (WHERE EVER APPLICABLE)**

210

211 We declare that 'written informed consent was obtained from the patient (or other approved parties) for  
212 publication of this case report and accompanying images. A copy of the written consent is available for  
213 review by the Editorial office/Chief Editor/Editorial Board members of this journal."

214

215

## 216 **ETHICAL APPROVAL**

217

218 We declare that we have obtained all necessary ethical approval from our Institutional ethical Committee.

219

## 220 **REFERENCES**

221

222 1, Kumar A, Shekhar S, Dhole B. Thyroid and male reproduction. Indian J Endocr Metab 2014;18:23-31.

223

224 2,Hollowell JG, Staehling NW, Flanders WD, Hannon WH, Gunter EW, Spencer CA, *et al.* Serum TSH,  
225 T(4), and thyroid antibodies in the United States population (1988 to 1994): National Health and Nutrition  
226 Examination Survey (NHANES III). J Clin Endocrinol Metab 2002;87(2):489–99

227

228 3,Canaris GJ, Manowitz NR, Mayor G, Ridgway EC. The Colorado thyroid disease prevalence study.  
229 Arch Intern Med 2000;160:526–34

230

231 4, Unnikrishnan AG, Menon UV. Thyroid disorders in India: An epidemiological perspective. Indian J  
232 Endocr Metab 2011;15( Suppl S2):78-81

233

234 5, Cooper TG, Noonan E, von Eckardstein S, Auger J, Baker HW, Behre HM, et al. World Health  
235 Organization reference values for human semen characteristics. Hum Reprod Update. 2010;16(3):231-  
236 45.

237

238 6, Oppenheimer, J. H., Schwartz, H. L. & Surks, M. I. Tissue differences in the concentration of  
239 triiodothyronine nuclear binding sites in the rat: liver, kidney, pituitary, heart, brain, spleen and testis.  
240 Endocrinology 1974; 95:897–903.

241  
242 7, Jana NR, Bhattacharya S. Binding of thyroid hormone to goat testicular cell induces generation of a  
243 proteinaceous factor which stimulates androgen release. *J Endocrinol* 1994;143:549-56.  
244  
245 8, Trokoudes KM, Sugeno A, Hazani E, Row VV, Volpe R. Thyroid-stimulating hormone (TSH) binding  
246 to extrathyroidal human tissues: TSH and thyroid-stimulating immunoglobulin effects on adenosine 3',5'-  
247 monophosphate in testicular and adrenal tissues. *J Clin Endocrinol Metab* 1979;48:919-23.  
248  
249 9, Jaya Kumar B, Khurana ML, Ammini AC, Karmarkar MG, Ahuja MM. Reproductive endocrine functions  
250 in men with primary hypothyroidism: effect of thyroxine replacement. *Horm Res* 1990;34:215-8.  
251  
252 10. Donnelly P, White C. Testicular dysfunction in men with primary hypothyroidism; Reversal of  
253 hypogonadotropic hypogonadism with replacement thyroxine. *Clin Endocrinol* 2000;52:197-201.  
254  
255 11, Kumar A, Chaturvedi PK, Mohanty BP. Hypoandrogenaemia is associated with subclinical  
256 hypothyroidism in men. *Int J Andro* 2006;30:14-20.  
257  
258 12, Welsh TH, Kasson BG, Hsueh AW. Direct biphasic modulation of gonadotropin-stimulated testicular  
259 androgen biosynthesis by prolactin. *Biol Reprod* 1986;34:796-804.  
260  
261 13, Weiss-Messer, E., Ber, R. & Barkey, R. J. Prolactin and MA-10 Leydig cell steroidogenesis: biphasic  
262 effects of prolactin and signal transduction. *Endocrinology* 1996;137:5509-18.  
263  
264 14, Trummer H, Ramschak-Schwarzer S, Haas J, Habermann H, Pummer K, Leb, G. Thyroid hormones  
265 and thyroid antibodies in infertile males. *Fertility and Sterility* 2001;76:254-57  
266  
267 15, Lei J, Nowbar S, Mariash CN, Ingbar DH. Thyroid hormone stimulates Na-K-ATPase activity and its  
268 plasma membrane insertion in rat alveolar epithelial cells. *Am J Physiol Lung Cell Mol Physiology*  
269 2003;285:762-72.  
270  
271 16, Wrutniak-Cabello C, Casas F, Cabello G. Thyroid hormone action in mitochondria. *J Mol Endocrinol*  
272 2001;26:67-77.  
273