

**Alteration of sex hormone and semen parameters in adult males with subclinical hypothyroidism**

**Dipti Sarma<sup>1</sup>, Uma Kaimal Saikia<sup>1</sup>, Darvin Vamadevan Das<sup>1</sup>**

<sup>1</sup> *Department of Endocrinology, Gauhati Medical College and Hospital, Guwahati, India.*

**Aim:** To investigate the correlation between subclinical hypothyroidism (SCH) with serum testosterone levels and semen parameters (sperm count, total motility and morphology) in men seeking medical care for sexual dysfunction and infertility.

**Study design:** cross-sectional study.

**Place and Duration of Study:** Department of Endocrinology, Gauhati Medical College and hospital, Guwahati, India. Study done from November 2015 to December 2016.

**Methodology:** The patients in study were grouped into two groups: Group I – Men with subclinical hypothyroidism (n=35) and group II – Euthyroid males serving as controls (n=27). The medical history, clinical examination, semen analysis, total thyroxine(T4), total triiodothyronine(T3), thyroid-stimulating hormone(TSH), Luteinizing hormone (LH), Follicular stimulating hormone (FSH), Total testosterone (T) and prolactin (PRL) were obtained. Patients with diabetes mellitus, hypertension, chronic diseases, any testicular or pituitary diseases and prior chemo-radiotherapy were excluded from study.

**Results:** The age of the patients ranged between 24 and 42 years with mean age 31.57 years. The cases had significantly lower levels of mean serum total testosterone and free T4 and higher serum prolactin levels compared to controls ( $P < .01$ ). Linear regression analysis showed TSH levels independently associated with low testosterone levels. Also men with subclinical hypothyroidism had lower total sperm motility with preserved sperm count and morphology.

**Conclusion:** We found that SCH is associated with reduction in testosterone levels and total sperm motility. Our study data conclude that SCH may be a contributing factor for hypoandrogenemia and sexual dysfunction in men.

*Keywords: Subclinical hypothyroidism; testosterone; sperm count; total sperm motility.*

## **1. INTRODUCTION**

Male reproductive function is orchestrated by the hypothalamo-hypophyseal testicular axis. In addition it has been seen that thyroid hormones exerts a modulatory effect on this axis. Thus thyroid hormones seem to play a major role in male sexual and spermatogenic function. Primary hypothyroidism has been associated with low levels of both total and free testosterone in men [1]. The worldwide prevalence of subclinical hypothyroidism is found to be between 4 and 10% [2, 3]. The prevalence of subclinical hypothyroidism in India is about 9.4%. In women, the prevalence was higher (11.4%), when compared with men (6.2%) [4]. But links between subclinical hypothyroidism and male sexual dysfunction and

23 infertility are less evident. Till date there has been no concrete study that has showcased this relationship.  
24 Our study aims to investigate the association between subclinical hypothyroidism with total testosterone  
25 levels and semen parameters (sperm count, total motility and morphology) in men seeking medical care  
26 for sexual dysfunction and infertility in tertiary care centre.

## 27 28 **2. MATERIAL AND METHODS**

29  
30 The study is a cross sectional study done in males of age between 20-45 years, attending Endocrinology  
31 department of Gauhati Medical College and Hospital, Assam, India from November 2015 to December  
32 2016 for sexual dysfunction or infertility. The patients in the study were grouped into two: Group I - Men  
33 who were subclinical hypothyroidism (n=35) and group II- Men who were normal euthyroid controls  
34 (n=27). The most common sexual dysfunction among the subclinical hypothyroid males was erectile  
35 dysfunction (26 out of 35) and premature ejaculation (9 out of 35). On the other hand all men among the  
36 euthyroid control had presented with only premature ejaculation as the sexual dysfunction. Patients with  
37 diabetes mellitus, hypertension, any chronic disease, testicular or pituitary disorder and prior chemo  
38 radiotherapy were excluded from study. Patient who fulfilled the study criteria underwent a detailed  
39 medical history and clinical examination. Testicular volume was measured clinically by Prader  
40 orchidometer. Serum samples were collected for free thyroxine (FT4), total triiodothyronine (T3), thyroid-  
41 stimulating hormone (TSH), luteinizing hormone (LH), follicular stimulating hormone (FSH), total  
42 testosterone (T) and prolactin (PRL) estimation. Blood samples were collected in the morning after  
43 subjects had fasted for at least 8 hours. The samples were immediately centrifuged and the samples  
44 were aliquoted into 2-ml tubes .The serum was stored at -80 °C until assayed. Hormonal assay was done  
45 by enzyme-enhanced chemiluminescent analyzer (IMMULITE 1000). Single semen analysis was also  
46 done in pathology department of our medical college with fresh samples collected within 30 minutes of  
47 ejaculation. World Health Organization 2010 reference values for human semen characteristics were  
48 considered in the study [5]. The study was performed according to the guidelines of the Ethics Committee  
49 of our institute with informed consent. Statistical analysis was performed by the use of SAS 9.3. Pearson  
50 correlation between TSH levels and LH, FSH, PRL, total testosterone levels and semen parameters were  
51 estimated. Linear regression model analysis was used to find independent association between TSH and  
52 total testosterone levels .Multiple regression analysis model was used to find the simultaneous effect of  
53 thyroid hormones (TSH, Free T4 and Total T3) on the serum total testosterone levels. Statistical  
54 significance was set at the 0.05 level.

## 55 56 **.3. Results**

### 57 58 **3.1 Baseline Characteristics**

59 Table 1 shows the baseline characteristics of the cases and control.  
60  
61

62  
63


<b>Characteristics of subjects</b>	<b>Case (n=35) Mean ± SD</b>	<b>Control (n=27) Mean ± SD</b>	<b>P value</b>
Age( years)	31.57 ± 5.43	35.59 ± 4.48	0.0037
Body mass index (kg/m <sup>2</sup> )	24.13 ± 1.10	24.65 ± 1.36	0.1054
Testicular volume (in ml)	20.85 ± 3.53	18.51 ± 3.34	0.0104
TSH ( mIU/L )	9.09 ± 2.53	2.36 ± 0.96	<.0001
Total Testosterone (ng /dl )	414.22 ± 225.48	683.20 ± 328.29	0.0003
Prolactin (ng /ml )	16.22 ± 4.70	10.77 ± 2.48	<.0001
LH (mIU/mL)	4.40 ± 0.87	3.93 ± 0.89	0.041
FSH (mIU/mL)	4.70 ± 1.24	4.63 ± 0.87	0.215
Sperm count (million per ml)	50.80 ± 21.86	52.44 ± 19.75	0.7606
Semen volume (ml)	3.165 ± 1.15	2.87 ± 1.07	0.3125
Total sperm motility (in %)	53.14 ± 25.83	82.81 ± 16.83	<.0001
Duration of symptoms(years)	4.97 ± 2.52	3.07 ± 1.73	0.0014
Free T4 (ng /dl)	1.17 ± 0.14	1.32 ± 0.07	<.0001
Total T3 (nmol/L)	1.92± 0.27	1.84 ± 0.189	0.1835

64  
65  
66  
67  
68  
69  
70  
71  
72  
73

As seen in table 1, the age of the patients ranged between 24 and 42 years with mean age 31.57 years (SD ± 5.43). There were no differences between subclinical hypothyroid cases and euthyroid control in terms of mean BMI, mean FSH, mean sperm count, mean semen volume and total T3. However, the cases had significant lower level of mean serum total testosterone levels (414.22 ± 225.48 in cases vs. 683.20 ± 328.29 in controls) and free T4 (1.17 ± 0.14 in cases vs. in 1.32 ± 0.07 controls ) but a higher level of serum prolactin (16.22 ± 4.70 in cases vs. in 10.77 ± 2.48 controls ) in comparison to control group. LH levels were also significantly different between the two groups (4.40 ± 0.87 in cases vs. 3.93 ± 0.89 controls). Among the semen parameters, the cases had a remarkable lower total sperm motility compared to controls groups (53.14 ± 25.83 in cases vs. 82.81± 16.83 in controls).


74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84

Figure1 (A-D) shows the bar diagram comparing the mean values with standard error of mean of various hormones (TSH, Prolactin, LH and testosterone)



85  
86  
87  
88  
89  
90  
91  
92  
93

Figure 2 (E&F) shows the bar diagram comparing the mean values semen parameters among the cases and control groups. Sperm morphology was normal in both the cases and control group as per the Kruger's criteria.



94  
 95  
 96  
 97  
 98  
 99  
 100  
 101  
 102  
 103  
 104

### 3.2 Correlations between TSH and other hormones and semen parameters

Among the subclinical hypothyroid patients, a significant negative correlation was observed between serum TSH and total testosterone levels ( $r = -0.75, P < .01$ ). Negative correlation was also found between total testosterone levels and serum prolactin ( $r = -0.69, P < .01$ ). Figure 3 (G&H) demonstrates these relationship. Similarly a negative correlation was also seen among the cases between serum TSH and total sperm motility ( $r = -0.73, P < .01$ ). No correlations were found between TSH levels and FSH, LH or sperm count.





Figure 3 :Fitted line plot showing negative linear relationship between TSH levels and testosterone levels (G) and testosterone levels and prolactin (H) among the subclinical hypothyroid males (n=35). The solid line shows the mean testosterone levels among the subclinical hypothyroid males and the open circles shows the distribution of TSH levels (G) and prolactin levels (H) of each cases .

105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116

### 3.3 Correlations between testosterone and Free T4 and total T3

A significant positive correlation was found between total testosterone levels and free T4 and Total T3 levels ( $r = 0.54, P < .01$ ;  $r = 0.65, P < .01$ ). Figure 4 (I&J) demonstrate this relationship.



117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129

Linear regression model analysis showed significant association between TSH and total testosterone levels ( $P < .01$ ). Multiple regression model analysis was done to assess the simultaneous effect of thyroid hormones (TSH, Free T4, total T3) on serum testosterone. The model showed TSH and free T4 contributed significantly to low total testosterone level. Table 2 shows multiple regression analysis.

Table 2: Multiple regression analysis showing the relationship between serum testosterone and thyroid hormones.

Hormones	B – coefficient	P Value
TSH	-100.41507	0.0001
Free T4	-920.19167	0.0122
Total T3	147.03873	0.3160

130  
131  
132  
133  
134  
135  
136  
137  
138

#### 4. DISCUSSION

In the current study we found a significant reduction in serum total testosterone levels in subclinical hypothyroid males compared to euthyroid control. It was also seen that subclinical hypothyroid patients the free T4 levels were significantly lower compared to the control group. It is known that the levels of T3 are dependent on the availability of T4 hormone. We could find a positive correlation between total

139 testosterone levels and total T3 concentration ( $r = 0.65$ ,  $P < .01$ ) among the subclinical hypothyroid  
140 males. This significant correlation explains the role of total T3 in the production of testosterone. The role  
141 of T3 hormone in testosterone synthesis has been demonstrated earlier in Leydig cells of rat and goat [6,  
142 7]. These finding supports the modulatory effects of thyroid hormones in maintaining the normal  
143 testosterone levels and male sexual function. However, it has to be understood that serum T3  
144 concentration in subclinically hypothyroid cases was along the lower side but within the normal reference  
145 range in this study. Secondly linear regression model analysis showed a significant independent  
146 association between TSH and total testosterone levels among subclinical hypothyroid males. The  
147 probable explanation lies in the fact that thyroid-stimulating hormone receptors have also been  
148 demonstrated in human testis [8]. It is postulated by some authors that TSH may have a cardinal role in  
149 regulating testosterone synthesis in leydig cells. However, till date there is no concrete evidence  
150 supporting the above. It has to be understood that a negative relationship was found between TSH levels  
151 and total testosterone levels. Multiple regression model analysis was done and showed the effects of  
152 TSH and thyroid hormones together in generation of testosterone in males. The analysis showed that  
153 TSH (beta coefficient: - 100.4) along with thyroid hormones simultaneously has a significant role in  
154 testosterone production.

155  
156 The rise of LH levels seen in subclinical hypothyroid males has been a significant finding compared to the  
157 euthyroid control. We postulate that this may be due to loss of feedback mechanism by low levels of  
158 testosterone. It should be understood that LH or FSH feedbacks are under the control of free testosterone  
159 levels. Our study has been based on total testosterone and hence further confirmation is needed.  
160 Moreover the rise of LH and FSH levels in overt hypothyroidism itself is controversial [9,10]. We also  
161 found a significant elevation of serum prolactin levels among the subclinical hypothyroid patients. This  
162 was similar to study done by Kumar et al (2006) [11]. Multiple mechanism have been explained as the  
163 cause of hypoandrogenemia in elevated levels of prolactin, the degree of hyperprolactinemia required to  
164 cause low levels of testosterone has been a consistent finding with overt hypothyroids males. The  
165 mechanisms illustrated in overt hypothyroid males include suppression of 17 alphahydroxylase (a key  
166 enzyme in conversions of progesterone to testosterone in Leydig cells) [12] and decreasing the binding  
167 affinity of LH to its receptors (seen on a murine tumour cell line, MA-10 cells) [13] by high concentration of  
168 prolactin. In our study not only a higher levels of serum prolactin were found among the subclinical  
169 hypothyroid males but also a significant negative correlation found between total testosterone and serum  
170 prolactin levels, ( $r = -0.69$ ,  $P < .01$ ) which probably explains that elevated prolactin as the cause of low  
171 levels of total testosterone among the cases. Moreover Kumar et al (2006) in his study has demonstrated  
172 the reduced availability of progesterone (precursor for the synthesis of testosterone) in subclinical  
173 hypothyroid males. Summating all the above data we may conclude that hyperprolactinemia in subclinical  
174 hypothyroid males may lead to hypoandrogenemia a picture similar to overt hypothyroidism.

175  
176 .



177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214

Finally we also learned the relationship between subclinical hypothyroidism and semen parameters. Trummer et al had shown that subclinical hypothyroidism had no impact on semen parameter [14]. The only semen parameter which was affected in our study was the total motility without affecting sperm count and morphology. TSH levels correlated with sperm total motility ( $r = -0.73, P < .01$ ). In overt hypothyroidism it has been seen that low levels of thyroid hormone may affect forward progressive motility. Mechanism postulated is that thyroid hormones stimulate cellular oxygen consumption [15] and increase mitochondrial number over the middle and tail piece of sperm [16] and thus promote sperm motility. This probably is a major drawback in our study as we were unable to subclassify the types of motility defects (progressive motility, non-progressive motility or immotile sperms) among the subclinical hypothyroid males due to limited data. Though motility defects have been demonstrated among hypothyroid men in earlier studies, motility defects among subclinical hypothyroid males have to be reconsidered. Moreover the mean total motility among the subclinical hypothyroid males fell into a mean value of 53%, which was within the lower reference as per the World Health Organization reference values for human semen characteristics [5]. The small sample size among the cases and control may be another drawback in the study. We also couldn't estimate free testosterone levels among the patients which may be needed to explain the rising levels of gonadotropin levels among the subclinical hypothyroid males. Our strength of the study is that the relationship between subclinical hypothyroidism and male sexual dysfunction has been demonstrated. The comparison done between various parameters among case and control has been studied with accuracy. Till date there have been no studies in depth showcasing the relationship between male hypogonadism and altered semen characteristics in subclinical hypothyroid males. This study has opened a window for discussion regarding replacement of thyroid hormone in subclinical hypothyroid males who present with male sexual dysfunction or infertility.

## 5. CONCLUSION

In conclusion from our study it was seen that subclinical hypothyroidism may lead to low levels of total testosterone levels. Spectrum of male sexual dysfunction may be a presentation of subclinical hypothyroidism. Till now there is no consensus to treat subclinical hypothyroidism in male. Further studies may be needed to confirm the need for supplementing thyroid hormones to this class of patients. The genesis of infertility in subclinical hypothyroid males remains a domain of interest. Though the study has concluded the effects of subclinical hypothyroidism on total sperm motility more studies may be required to confirm this findings.

215 **CONSENT (WHERE EVER APPLICABLE)**

216  
217 We declare that 'written informed consent was obtained from the patient (or other approved parties) for  
218 publication of this case report and accompanying images. A copy of the written consent is available for  
219 review by the Editorial office/Chief Editor/Editorial Board members of this journal.'  
220

221

222 **ETHICAL APPROVAL**

223  
224 We declare that we have obtained all necessary ethical approval from our Institutional ethical Committee.  
225

226

227 **REFERENCES**

228 1, Kumar A, Shekhar S, Dhole B. Thyroid and male reproduction. Indian J Endocr Metab 2014;18:23-31.  
229

230 2, Hollowell JG, Staehling NW, Flanders WD, Hannon WH, Gunter EW, Spencer CA, *et al.* Serum TSH,  
231 T(4), and thyroid antibodies in the United States population (1988 to 1994): National Health and Nutrition  
232 Examination Survey (NHANES III). J Clin Endocrinol Metab 2002;87(2):489–99  
233

234 3, Canaris GJ, Manowitz NR, Mayor G, Ridgway EC. The Colorado thyroid disease prevalence study.  
235 Arch Intern Med 2000;160:526–34  
236

237 4, Unnikrishnan AG, Menon UV. Thyroid disorders in India: An epidemiological perspective. Indian J  
238 Endocr Metab 2011;15( Suppl S2):78-81  
239

240 5, Cooper TG, Noonan E, von Eckardstein S, Auger J, Baker HW, Behre HM, *et al.* World Health  
241 Organization reference values for human semen characteristics. Hum Reprod Update. 2010;16(3):231-  
242 45.  
243

244 6, Oppenheimer, J. H., Schwartz, H. L. & Surks, M. I. Tissue differences in the concentration of  
245 triiodothyronine nuclear binding sites in the rat: liver, kidney, pituitary, heart, brain, spleen and testis.  
246 Endocrinology 1974; 95:897–903.  
247

248 7, Jana NR, Bhattacharya S. Binding of thyroid hormone to goat testicular cell induces generation of a  
249 proteinaceous factor which stimulates androgen release. J Endocrinol 1994;143:549-56.  
250

251 8, Trokoudes KM, Sugeno A, Hazani E, Row VV, Volpe R. Thyroid-stimulating hormone (TSH) binding  
252 to extrathyroidal human tissues: TSH and thyroid-stimulating immunoglobulin effects on adenosine 3',5'-  
253 monophosphate in testicular and adrenal tissues. J Clin Endocrinol Metab 1979;48:919-23.  
254

255 9, Jaya Kumar B, Khurana ML, Ammini AC, Karmarkar MG, Ahuja MM. Reproductive endocrine functions  
256 in men with primary hypothyroidism: effect of thyroxine replacement. Horm Res 1990;34:215-8.  
257

258 10. Donnelly P, White C. Testicular dysfunction in men with primary hypothyroidism; Reversal of  
259 hypogonadotrophic hypogonadism with replacement thyroxine. Clin Endocrinol 2000;52:197-201.  
260

261 11, Kumar A, Chaturvedi PK, Mohanty BP. Hypoandrogenaemia is associated with subclinical  
262 hypothyroidism in men. Int J Andro 2006;30:14-20.  
263

264 12, Welsh TH, Kasson BG, Hsueh AW. Direct biphasic modulation of gonadotropin-stimulated testicular  
265 androgen biosynthesis by prolactin. Biol Reprod 1986;34:796-804.  
266

267 13, Weiss-Messer, E., Ber, R. & Barkey, R. J. Prolactin and MA-10 Leydig cell steroidogenesis: biphasic  
268 effects of prolactin and signal transduction. Endocrinology 1996;137:5509–18.  
269

270 14, Trummer H, Ramschak-Schwarzer S, Haas J, Habermann H, Pummer K, Leb, G. Thyroid hormones  
271 and thyroid antibodies in infertile males. *Fertility and Sterility* 2001;76:254–57  
272  
273 15, Lei J, Nowbar S, Mariash CN, Ingbar DH. Thyroid hormone stimulates Na-K-ATPase activity and its  
274 plasma membrane insertion in rat alveolar epithelial cells. *Am J Physiol Lung Cell Mol Physiology*  
275 2003;285:762-72.  
276  
277 16, Wrutniak-Cabello C, Casas F, Cabello G. Thyroid hormone action in mitochondria. *J Mol Endocrinol*  
278 2001;26:67-77.  
279  
280