

# QT interval variability in patients with obstructive sleep apnea

## ABSTRACT

**Aims:**

Obstructive sleep apnea (OSA) increases the risk of cardiac arrhythmias. We investigated QT interval variability among OSA patients.

**Study design:** It is a descriptive-cross-sectional study.

**Methodology:** Newly diagnosed OSA patients and healthy controls were studied. Inter-heartbeat and QT intervals were extracted from electrocardiography (for 1 hour at 3 AM). QT interval metrics including duration and variability indexes were compared between patients and controls.

**Results:** 35 patients and 13 controls were studied. There was no difference between patients and controls, neither between mild/moderate OSA versus severe OSA patients, in the measured QT interval variables. No significant correlation was found between apnea severity and the measured QT interval variables.

**Conclusion:** We found no difference between OSA patients and controls in QT variability. Also, no clear association between OSA severity and QT variability was observed.

*Keywords: Sleep apnea, arrhythmia, cardiovascular, cardiac electrophysiology*

## 1. INTRODUCTION

Obstructive sleep apnea (OSA) is associated with cardiac arrhythmias.<sup>[1]</sup> Obstructive episodes during sleep increase cardiac load which can induce cardiac remodeling and structural changes contributing to arrhythmias.<sup>[2-4]</sup> Given that the renin-angiotensin-aldosterone system's stimulation and sympathetic activity can be increased in OSA patients, the higher risk of cardiac arrhythmias and sudden death can be expected for this group of patients.<sup>[5,6]</sup> Characteristics of the QT interval, including QT variability (QTV), can provide information regarding cardiac electrical activity in OSA patients. Abnormalities of the QT interval can indicate abnormal repolarization during which cardiac vulnerability is heightened toward the development of arrhythmias. The QTV index (QTVi) assesses repolarization lability and is a predictor of cardiac arrhythmias and mortality.<sup>[7]</sup> QT interval variation may happen during and after sleep apnea episodes due to increased vagal activity and subsequent increased sympathetic tone and/or vagal withdrawal.<sup>[8]</sup> Camen et al. showed that simulated obstructive hypopnea/apnea are associated with prolongation of the QT interval.<sup>[2]</sup> However, others found no difference between OSA patients and controls in QT interval.<sup>[9]</sup> Baumert et al. reported an association between severity of OSA and QTV, reflecting alterations in cardiac sympathetic activity.<sup>[11]</sup> We evaluated QTV in patients with newly diagnosed OSA and compared it with healthy controls.

## 2. MATERIAL AND METHODS

This is a descriptive analytic study in which all patients presenting with symptoms of sleep apnea, loud and frequent snoring at night, severe obesity, and enlarged adenoids at Bamdad sleep clinic in

34 2017 were considered as candidates for polysomnography (PSG) of which 50 persons were selected  
35 through convenience sampling.

36 The inclusion criteria were patients aged at least 18 years, with OSA diagnosed by PSG. Those with  
37 cardiovascular, kidney, or pulmonary diseases or diabetes were excluded (2 cases with indications of  
38 diabetes). Thus, a sample size of 48 was considered. The study was approved by local committee of  
39 Isfahan University of Medical Sciences. A written informed consent was obtained from each  
40 participant undergoing PSG. Apnea–hypopnea index (AHI) of  $\geq 5$  events/hour was considered  
41 diagnostic.[6] Patients were categorized to mild ( $5 \leq \text{AHI} < 15$ ), moderate ( $15 \leq \text{AHI} \leq 30$ ), severe ( $\text{AHI} > 30$ )  
42 OSA.[7] and controls ( $\text{AHI} < 5$ ).<sup>[10]</sup> On a separate day, Holter electrocardiography was performed using  
43 a digital Holter recorder with three channels (I, II, and III) and sampling rate of 200 Hz (H200 recorder,  
44 Kavoshgaran Teb Kharazmi Co., Iran). Software was developed by the Department of Biomedical  
45 Engineering (Isfahan University) for signal processing. The QT and inter-heartbeat (RR) intervals  
46 were extracted from the channel II and were inspected by a cardiologist for errors. The QTV analyses  
47 were performed for an hour of sleeping period at night (3-4 AM). The QTVi was calculated as  
48  $\log[(\text{QTvar}/\text{meanQT}^2)/(\text{RRvar}/\text{meanRR}^2)]$ , where QTvar contains the variance of all QT intervals and  
49 RRvar contains the variance of all RR intervals during an hour.<sup>[11]</sup>

## 50 2.1 Statistical analysis:

51 Data analysis was performed using SPSS software (version 16.0, SPSS Inc., Chicago, IL).  
52 Continuous variables were compared between the groups using Independent sample t-Test or Mann-  
53 Whitney U test. Categorical data were compared using the Chi-square test. Pearson and Spearman  
54 correlation analyses were used to estimate the relationship between variables. A *P* value  $< 0.05$  was  
55 considered significant in all analyses.

## 56 3. RESULTS AND DISCUSSION

57  
58 35 patients and 13 controls were included into the study. The two groups were not different regarding  
59 age, gender, or BMI ( $P > 0.05$ ). The QT interval metrics are summarized in Table 1. There was no  
60 significant difference between patients and controls, neither between mild/moderate OSA versus  
61 severe OSA patients, in QT interval variables ( $P > 0.05$ ). No significant correlation was found between  
62 AHI and QT interval variables ( $P > 0.05$ ).

63

64

**Table 1: Comparison of demographic data between the patients and controls**

	Mild/Moderate OSA AHI 5 to 30, n = 19	Severe OSA AHI >30, n = 16	Controls AHI <5, n = 13	P*	P**
Age, year	47.5 ± 6.7	48.1 ± 7.6	47.7 ± 6.5	0.969†	0.805†
Male gender	9 (47.4)	8 (50)	7 (53.8)	0.500‡	0.937‡
BMI, kg/m <sup>2</sup>	28.0 ± 4.2	29.9 ± 6.6	27.2 ± 1.9	0.294†	0.311†
AHI, /h	19.0 ± 6.4	56.4 ± 22.5	2.4 ± 1.2	<0.001†	<0.001†
QT, ms	406.6±30.2	396.1±23.3	407.5±27.9	0.524†	0.266†
QTSD, ms	12.7±6.0	12.4±4.0	12.0±5.0	0.733†	0.875†
QTvar, ms <sup>2</sup>	197.8 [49.7]	171.0 [31.5]	168.9 [40.7]	0.539‡	0.832‡
logQTvar, ms <sup>2</sup>	2.13±0.36	2.15±0.25	2.09±0.34	0.767†	0.666†
QTVi, nu	0.30± [0.10]	0.18± [0.05]	0.33± [0.09]	0.143‡	0.193 ‡
logQTVi, nu	-0.74 ± 0.40	-0.89 ± 0.37	-0.63 ± 0.38	0.167†	0.152†

66 Data are presented as mean ± standard deviation [or standard of error] and number (%)

67 OSA: Obstructive sleep apnea; BMI: Body mass index; AHI: Apnea/hypopnea index (/h); QTSD: Standard  
68 deviation of QT interval; QTvar: Variance of QT interval; QTVi: QT variability index

69 \* Patients vs. controls

70 \*\* Mild/moderate OSA vs. severe OSA

71 † Independent sample *t*-Test

72 ‡ Chi-square test

73 † Mann-Whitney U Test

74

75

76

77

### 3.1 DISCUSSION

78 The cardiovascular disease is of importance particularly among OSA patients.<sup>[12]</sup> Electrocardiogram  
79 (ECG) parameters including QT interval (QT) and QT dispersion (QTd) are usually applied to evaluate  
80 myocardial repolarization.<sup>[13, 14]</sup>

81 Based on results of the current study, we found no difference between OSA patients and controls in  
82 the measured QT interval metrics, neither an association between OSA severity and QT dynamicity.

83 We analyzed cardiac activity of only one hour of sleeping time (3-4 AM). In contrast, Baumert et al.<sup>[15]</sup>  
84 analyzed all consecutive 5-min ECG segments throughout the night which provides quasi-stationary  
85 conditions of RR and QT time series for all sleeping time. QTV may be affected by sleep stages due  
86 to sleep-related variations in autonomic function.<sup>[15-16]</sup> Baumert et al. also found that QT variability  
87 was elevated in 5-min epochs that contained sleep apnea events.<sup>[15]</sup> Therefore, it is important for the  
88 future studies to perform cardiac monitoring at the same time as polysomnography to be able to  
89 analyze QTV in different sleep stages and to evaluate temporal association between apnea/hypopnea  
90 events and QT variations. Shamsuzzaman et al. showed daytime increase in rate-corrected QT in  
91 OSA patients.<sup>[18]</sup> Accordingly, daytime monitoring will provide more comprehensive information about  
92 diurnal variation of QT in OSA patients.

93 Due to potential restrictions of the device, some of limitations in the study were unavoidable such as  
94 measuring RR interval (60 / heart rate) which is required to rate correction. This was a time-limited  
95 study. Therefore, it could not be carried out in a longer period of time.

96 REM sleep duration and REM density could affect both AHI and arrhythmic activity. However, this  
97 issue can be considered as a limitation of this study.<sup>[19]</sup>

98 Our results might also be affected by some technical aspects. ECG acquisition sampling rate in our  
99 study was lower than what is recommended for QTV analysis ( $\geq 500$  Hz).<sup>[20]</sup> Measurement of QT  
100 interval, either manually or automatically by a software, is challenging. Most of the available systems  
101 utilize simple tangent and threshold methods and our method was also detection of each of the Q, R,  
102 S, and T points based a gradient-based algorithm. These techniques may be less efficient compared  
103 to techniques that use ECG waveforms with pre-defined templates.<sup>[20]</sup> There are still controversial  
104 issues such as necessity for excluding ectopic and subsequent beats, the preferred lead for QT  
105 measurement, necessity for rate correction, and increasing reproducibility by random sampling across  
106 long-term monitoring which requires further investigations.

## 107 **CONCLUSION:**

108 In summary, we found no significant difference between OSA patients and controls in QT variability.  
109 Studies with cardiac monitoring during various sleep stages as well as monitoring diurnal variations of  
110 QT interval are still required to better investigate QTV as a possible index of cardiac vulnerability  
111 toward arrhythmias in OSA patients.

## 112 **Ethical Disclaimer:**

113 As per international standard or university standard written ethical permission has been collected and  
114 preserved by the authors.

115

116

## 117 **REFERENCES:**

118

119 1. Somers VK, White DP, Amin R, Abraham WT, Costa F, Culebras A, et al. Sleep apnea and  
120 cardiovascular disease: an American Heart Association/American College of Cardiology Foundation  
121 Scientific Statement from the American Heart Association Council for High Blood Pressure Research  
122 Professional Education Committee, Council on Clinical Cardiology, Stroke Council, and Council on  
123 Cardiovascular Nursing. *J Am Coll Cardiol* 2008;52:686-717.

124 2. Camen G, Clarenbach CF, Stowhas AC, Rossi VA, Sievi NA, Stradling JR, et al. The effects of  
125 simulated obstructive apnea and hypopnea on arrhythmic potential in healthy subjects. *Eur J Appl*  
126 *Physiol* 2013;113:489-96.

127 3. Korcarz CE, Peppard PE, Young TB, Chapman CB, Hla KM, Barnet JH, et al. Effects of Obstructive  
128 Sleep Apnea and Obesity on Cardiac Remodeling: The Wisconsin Sleep Cohort Study. *Sleep*  
129 2016;39:1187-95.

130 4. Dimitri H, Ng M, Brooks AG, Kuklik P, Stiles MK, Lau DH, et al. Atrial remodeling in obstructive  
131 sleep apnea: implications for atrial fibrillation. *Heart Rhythm* 2012;9:321-7.

132 5. Horner RL. Pathophysiology of obstructive sleep apnea. *Journal of cardiopulmonary rehabilitation*  
133 *and prevention*. 2008;28(5):289-98.

134 6. Viigimae M, Karai D, Pirn P, Pilt K, Meigas K, Kaik J. QT interval variability index and QT interval  
135 duration in different sleep stages: analysis of polysomnographic recordings in nonapneic male  
136 patients. *BioMed Research International*. 2015;2015.

137

138 7. Berry RB, Budhiraja R, Gottlieb DJ, Gozal D, Iber C, Kapur VK, et al. Rules for scoring respiratory  
139 events in sleep: update of the 2007 AASM Manual for the Scoring of Sleep and Associated Events.  
140 Deliberations of the Sleep Apnea Definitions Task Force of the American Academy of Sleep Medicine.  
141 *J Clin Sleep Med* 2012;8:597-619.

- 142 8. Gillis AM, Stoohs R, Guilleminault C. Changes in the QT interval during obstructive sleep apnea.  
143 Sleep 1991;14:346-50.
- 144 9. Kilicaslan F, Tokatli A, Ozdag F, Uzun M, Uz O, Isilak Z, et al. Tp-e interval, Tp-e/QT ratio, and Tp-  
145 e/QTc ratio are prolonged in patients with moderate and severe obstructive sleep apnea. Pacing Clin  
146 Electrophysiol 2012;35:966-72.
- 147 10. Epstein LJ, Kristo D, Strollo PJ Jr, Friedman N, Malhotra A, Patil SP, et al. Clinical guideline for  
148 the evaluation, management and long-term care of obstructive sleep apnea in adults. J Clin Sleep  
149 Med 2009;5:263-76.
- 150 11. Berger RD, Kasper EK, Baughman KL, Marban E, Calkins H, Tomaselli GF. Beat-to-beat QT  
151 interval variability: novel evidence for repolarization lability in ischemic and nonischemic dilated  
152 cardiomyopathy. Circulation 1997;96:1557-65.
- 153 12. Park DH, Shin CJ, Hong SC, Yu J, Ryu SH, Kim EJ, Shin HB, Shin BH. Correlation between the  
154 severity of obstructive sleep apnea and heart rate variability indices. Journal of Korean medical  
155 science. 2008;23(2):226-31.
- 156 13. Dursunoglu D, Dursunoğlu N, Evrengül H, Özkurt S, Kılıç M, Fisekci F, Kuru Ö, Delen Ö. QT  
157 interval dispersion in obstructive sleep apnoea syndrome patients without hypertension. European  
158 Respiratory Journal. 2005;25(4):677-81.
- 159 14. Bilal N, Dikmen N, Bozkus F, Sungur A, Sarica S, Orhan I, Samur A. Obstructive sleep apnea is  
160 associated with increased QT corrected interval dispersion: the effects of continuous positive airway  
161 pressure. Brazilian journal of otorhinolaryngology. 2017 Mar 31.
- 162 15. Baumert M, Smith J, Catcheside P, McEvoy RD, Abbott D, Sanders P, et al. Variability of QT  
163 interval duration in obstructive sleep apnea: an indicator of disease severity. Sleep 2008;31:959-66.
- 164 16. Penzel T, Kantelhardt JW, Lo CC, Voigt K, Vogelmeier C. Dynamics of heart rate and sleep  
165 stages in normals and patients with sleep apnea. Neuropsychopharmacology. 2003;28 Suppl 1:S48-  
166 53.
- 167 17. Boudreau P, Yeh WH, Dumont GA, Boivin DB. Circadian variation of heart rate variability across  
168 sleep stages. Sleep. 2013;36:1919-1928.
- 169 18. Shamsuzzaman A, Amin RS, van der Walt C, Davison DE, Okcay A, Pressman GS, et al. Daytime  
170 cardiac repolarization in patients with obstructive sleep apnea. Sleep Breath 2015;19:1135-40.
- 171 19. Nigam G, Camacho M, Riaz M. Rapid Eye Movement (REM) rebound on initial exposure to CPAP  
172 therapy: a systematic review and meta-analysis. Sleep Science and Practice. 2017 Dec;1(1):13.  
173
- 174 20. Baumert M, Porta A, Vos MA, Malik M, Couderc JP, Laguna P, et al. QT interval variability in body  
175 surface ECG: measurement, physiological basis, and clinical value: position statement and consensus  
176 guidance endorsed by the European Heart Rhythm Association jointly with the ESC Working Group  
177 on Cardiac Cellular Electrophysiology. Europace 2016;18:925-44.