

1 **NF- κ B, TNF- α and IL-6 levels in Liver and Kidney of High-Fructose-Fed Rats**

2

3

Short title: High Fructose Diet

4

5 **Abstract**

6

7 **Introduction:** Fructose constituting an important part of human diet, was reported to
8 facilitate fat depositing in the abdominal region in case of excessive consumption, therefore
9 increasing the risk of chronic illness more rapidly than expected, and inducing development
10 of various diseases such as diabetes, metabolic syndrome, hypertension and atherosclerosis.

11 **Aim:** The aim of this study was to investigate nuclear factor-kappa B (NF- κ B), tumor
12 necrosis factor-alpha (TNF- α) and interleukin-6 (IL-6) levels in liver and kidney tissues of
13 high-fructose-fed rats and to determine the role of dietary addition of fructose on
14 inflammation.

15 **Methods:** The rats were randomly divided into two groups of 7 rats as control (C) and
16 fructose (F). The fructose group received 30% (v/w) fructose in drinking water for 8 weeks.
17 Serum samples were used for aspartate aminotransferase (AST), alanine aminotransferase
18 (ALT), alkaline phosphatase (ALP), blood urea nitrogen (BUN) and uric acid measurements.
19 The liver and kidney tissues of the rats were washed with 0.9% NaCl for TNF- α , IL-6 and
20 NF- κ B measurements.

21 **Results:** TNF- α , IL-6 and NF- κ B levels in liver tissues were found significantly higher in the
22 fructose group than the control group ($p < 0.001$, $p < 0.05$, $p < 0.001$, respectively). TNF- α , IL-6
23 and NF- κ B levels in the kidney tissue of the fructose group were statistically significantly
24 higher than the control group ($p < 0.001$, $p < 0.001$, $p < 0.001$, respectively).

25 **Conclusion:** Fructose fed diet increased liver and kidney damage through augmenting
26 NF- κ B, TNF- α and IL-6 levels.

27

28 **Key words:** fructose, interleukin-6, nuclear factor-kappa B, tumor necrosis factor-alpha

29

30

31

32 1.INTRODUCTION

33 As a matter of fact, eating habits today are rapidly changing worldwide, which cause an
34 increased consumption of sugar and sugar-rich nutrients in the daily diet. Significant changes
35 occurred in carbohydrate content of foods, and intake of complex carbohydrates rich in fiber
36 content decreased while sugary products with high glycemic index increased. Fructose is the
37 major sweetener of sugar-sweetened beverages as part of sucrose molecule or high fructose
38 corn syrup (HFCS) besides glucose [1]. HFCS contains 42-90% fructose and is the main
39 source of fructose in the diet. Fructose is widely used in fruit juices, canned fruits, jams,
40 jellies, breakfast cereals and pastries. One of the reasons of high preference of fructose in
41 food industry is its lower cost in comparison to saccharose. Another reason is that fructose-
42 containing food and beverages delay the feeling of satiety while inducing the feeling of a new
43 hunger sooner [2]. Given that fructose is rapidly metabolized in the body and converted
44 directly into fatty acids unlike glucose, it was reported to contribute to lactic acidosis, liver
45 steatosis, obesity, insulin resistance, diabetes and lipid metabolism disorders and also to play
46 an important role in hypertension and cardiovascular diseases [3,4].

47 A wide variety of substances are secreted from the adipose tissue, including
48 inflammatory cytokines to begin with. Cytokines are chemical signaling molecules existing
49 in peptide or glycoprotein forms and mediate the development and regulation of
50 inflammatory and immune responses in membranes [5]. These molecules are mainly
51 produced by T cells and macrophages and secreted from cells with different characteristics.
52 Cytokines are classified in 2 groups, namely proinflammatory and antiinflammatory
53 cytokines, and interleukin-6 (IL-6) and tumor necrosis factor-alpha (TNF- α) are involved in
54 the proinflammatory group [6]. IL-6 is an important cytokine playing a role in many
55 physiological and pathological events, such as immune response, acute phase response of the
56 liver, hemopoiesis, regulation of neuronal functions and osteoclast formation. The synthesis
57 and circulating level of IL-6 increase in parallel to the adipose tissue. Approximately one-
58 third of the IL-6 in the systemic circulation originates from the adipose tissue [7]. IL-6 is
59 distinctive with its endocrine activity and as a circulating cytokine, where most other
60 cytokines function via autocrine or paracrine mechanisms. TNF- α is produced by monocytes
61 and macrophages, in response to a variety of inflammatory and immunomodulatory stimuli.
62 TNF- α has a considerably wide range of bioactivities, and most cells are sensitive to TNF- α .
63 Under normal physiological conditions, TNF- α is involved in the formation of immune
64 response, cellular homeostasis as well as cell survival, proliferation, migration and

65 differentiation. TNF- α enhances lipolysis and apoptotic adipocyte death by inhibiting
66 lipogenesis [8].

67 The liver is crucial for fructose metabolism, and for this very reason has a role in
68 important events such as metabolic dysfunction of glucose, dyslipidemia and inflammation.
69 Lipid accumulation leads to the development of oxidative stress in the liver and increased
70 activation of nuclear factor kappa B (NF- κ B) [9]. NF- κ B is a transcription factor and a
71 significant regulator of many genes responsible for inflammation, immune response,
72 lymphocyte activation, cell growth and apoptosis [10]. NF- κ B exists in all cell types and is
73 involved in the production of cellular responses to stimuli such as stress, inflammatory
74 cytokines, free radicals, ultraviolet radiation, carcinogens, tumor-forming agents,
75 chemotherapeutics, bacterial and viral agents. NF- κ B plays a central role in the
76 pathophysiology of clinically important diseases of many organ systems and inflammatory
77 cell damage [11,12].

78 This study was conducted to investigate NF- κ B, TNF- α and IL-6 levels in liver and
79 kidney tissues of high-fructose-fed rats and to determine the role of dietary addition of
80 fructose on inflammation.

81

82 **2. MATERIALS AND METHODS**

83

84 **2.1. Experimental Animals and Experimental Groups**

85 Atatürk University Experimental Animals Ethics Committee approval was obtained
86 before starting the study. As experimental animals, we used 14 male Wistar rats weighing
87 about 200-250 g and 8 weeks old. The rats were randomly divided into two groups of 7 rats
88 as control (C) and fructose (F). Experimental animals were kept under appropriate
89 temperature and ventilation conditions as well as in hygienic conditions. There was no
90 restrictions for feed and water. The control group was fed with the standard diet. The fructose
91 group received 30% (v/w) fructose in drinking water for 8 weeks [13].

92 At the end of eight-week experiment, all the rats in both groups were decapitated under
93 anesthesia with ketamine (60 mg/kg, Ketalar, Eczacibasi, Istanbul, Turkey) and xylazine (12
94 mg/kg; Rompun, Bayer, Istanbul, Turkey). Blood samples of the rats were rapidly collected
95 in biochemistry tubes and centrifuged at 4000xg for 10 minutes to separate the sera. Serum
96 samples were used for aspartate aminotransferase (AST), alanine aminotransferase (ALT),
97 alkaline phosphatase (ALP), blood urea nitrogen (BUN) and uric acid measurements. The
98 liver and kidney tissues of the rats were washed with 0.9% NaCl for TNF- α , IL-6 and NF- κ B

99 measurements and packaged and stored at -80°C in the deep freezer until the day of
100 measurement.

101

102 **2.2. Biochemical Analysis**

103 Serum AST, ALT, ALP, BUN and uric acid levels were determined by autoanalyzer.
104 Liver and kidney tissues were homogenized with 1/10 cold 0.1 M phosphate buffer (pH: 7.4)
105 by using homogenizer. Tissue homogenates were centrifuged at 4000xg and at 4°C for 20
106 minutes and supernatants were obtained. In the kidney and liver tissue samples, TNF- α , IL-6
107 and NF- κB levels were measured by Enzyme-Linked Immunosorbent Assay (ELISA) method
108 using commercial kits (SunRed Biological Technology, Co., Ltd. Shanghai). Absorbances
109 were read at 450 nm in the ELISA reader. The TNF- α , IL-6 and NF- κB results were stated as
110 ng/ml.

111 **2.3. Statistical analysis**

112 Statistical analysis of the data was performed using the Statistical Package for the
113 Social Sciences (SPSS) 24.0 program and Student's T-Test. All results were expressed as
114 mean \pm standard deviation and $p < 0.05$ was considered statistically significant.

115

116 **3. RESULTS**

117 Serum AST, ALT and ALP enzyme activities with BUN and uric acid levels of the
118 study groups are presented in Table 1. Serum AST, ALT, ALP enzyme activities and BUN
119 and uric acid levels of the fructose group were statistically significantly higher than the
120 control group ($p < 0.001$). TNF- α , IL-6 and NF- κB levels in liver and kidney tissues are
121 presented in Table 2. TNF- α , IL-6 and NF- κB levels in liver tissues were found significantly
122 higher in the fructose group than the control group ($p < 0.001$, $p < 0.05$, $p < 0.001$, respectively).
123 TNF- α , IL-6 and NF- κB levels in the kidney tissue of the fructose group were statistically
124 significantly higher than the control group ($p < 0.001$, $p < 0.001$, $p < 0.001$, respectively).

125

126 **4. DISCUSSION**

127 Changes in the living conditions and eating habits lead to more fructose intake due to
128 the preference of ready-to-eat foods, hence increased calories through daily diet. Different
129 carbohydrate types such as fructose, fructose-corn syrup and sucrose abounding in processed
130 and packaged, ready-to-eat foods and beverages, can cause metabolic dysfunctions associated
131 with chronic diseases due to high-energy contents and differences in their metabolisms

132 [14,15]. In this study, we investigated the effect of 30% fructose added to drinking water on
133 both liver and kidney tissues as well as on the liver enzyme profile.

134 Due to its roles in protein synthesis, carbohydrate and lipid metabolisms and being the
135 main organ responsible for fructose metabolism, liver has an important place in xenobiotic
136 metabolism and it is the organ affected at most by the metabolic changes and damaged first.
137 For this reason, loss of liver function is vital for the human beings [16]. Both clinical and
138 experimental studies in the literature reported that liver function markers such as serum AST,
139 ALT and ALP had the tendency to increase in obesity and non-alcoholic liver steatosis
140 [17,18,19]. In this study too, we observed a statistically significant increase in the serum
141 levels of AST, ALT and ALP of the fructose-fed group compared to the control group.
142 Masterjohn et al. [20] reported plasma ALT and AST activities were elevated in comparison
143 to the control group in high fructose-fed mice. In another study, high fructose-fed rats
144 displayed higher levels of plasma ALT and AST activities than the control group [21]. There
145 are studies in the literature reporting similar results as ours as well as those with opposite
146 results. Ackerman et al. [22] showed that there was no change in ALT and AST levels of rats
147 fed a diet containing 60% fructose for 5 weeks. Various mechanisms and different mediators
148 are mentioned in fructose-induced renal damage. BUN and uric acid are metabolic waste
149 products that are excreted from the kidneys. In this study, we observed a statistically
150 significant increase in the serum levels of BUN and uric acid of the fructose-fed group
151 compared to the control group. When designing this study, we knew that fructose caused liver
152 and kidney diseases. But the primary factor initiating and developing disease process was the
153 real question that should be answered. Oxidative stress and inflammation are the issues
154 discussed at most. In our study, by means of both kidney and liver tissues, we tried to
155 determine how strong was the effect of fructose on inflammation. Cytokines are molecules
156 having roles in various biological processes such as inflammation, apoptosis, necrosis, and
157 fibrosis [23]. They are usually released from lymphocytes and monocytes and are effective in
158 intercellular communication and modulation of the immune response. By inhibiting apoptosis
159 during the inflammatory process, IL-6 triggers events that develop chronic disease
160 progression [24].

161 The release of IL-6 is induced by interleukin-1 (IL-1) and TNF- α , and IL-6 displays
162 synergistic effects with these cytokines. The best defined effects of IL-6 are on hepatocytes
163 and B lymphocytes and cause hepatocytes to synthesize many plasma proteins that contribute
164 to the acute phase response. IL-6 was also reported to be associated with signal transduction
165 pathways and expected to increase in hepatic cirrhosis and liver cancer due to regeneration

166 [25]. Yang et al. [26] examined renal tissues of high fructose-fed rats and reported that
167 fructose reduced kidney weight, led to renal tubular damage, and increased TNF- α and IL-6
168 levels significantly. In our study too, there was a statistically significant increase in TNF- α
169 and IL-6 levels in both liver and kidney tissues compared to the control group. Gersch et al.
170 [27] reported that a high-fructose diet might induce inflammation, because fructose
171 accelerated renal disease in association with an increase in monocyte-macrophage infiltration.
172 The increase in adipose tissue due to high fructose consumption is one of the most important
173 causes of the secretion of inflammatory cytokines in the systemic circulation. Adipose tissue
174 dysfunction can impair energy consumption, primarily inflammatory cytokines and insulin
175 signaling [28,29]. Both human and rat studies in the literature stated that high fructose diet
176 caused increase in visceral adiposity, and as a response to that, inflammatory cytokines
177 increased and organ-related morphological and functional changes were observed [30,31].

178 In the literature, TNF- α was reported to induce NF- κ B activation, which leads to an
179 increase in proinflammatory cytokines through various mechanisms, resulting in an increase
180 in TNF- α release and consequently development of hepatosteatosis and insulin resistance
181 [32,33]. In our study, levels of NF- κ B, a transcription factor involved in the expression of
182 cytokines that cause proinflammatory cytokine elevation, were measured in both liver and
183 kidney tissues. The liver and kidney NF- κ B levels of the group with fructose added to
184 drinking water were significantly higher than the control group. In the study by Veličković et
185 al., [34] high fructose diet was reported to increase NF- κ B and TNF α levels in the liver. In
186 another study by Zheng et al., [35] excessive fructose consumption caused inflammation by
187 increasing fatty acid biosynthesis and fat accumulation in the liver. Vasiljević et al. [36]
188 reported increased levels of hepatic TNF- α and NF- κ B in high fructose-fed rats for 9 weeks
189 and stated that high fructose contributed to the development of NF- κ B-mediated
190 inflammation. NF- κ B is one of the key regulators in inflammatory processes and stimulates
191 the synthesis of TNF- α , IL-6 and adipokine, which increase in inflammation [37,38]. We can
192 say that fructose-induced chronic inflammation leads to the activation of NF- κ B, which
193 stimulates the release of proinflammatory cytokines as a result of cell damage in liver and
194 kidney tissues due to excessive fructose consumption [39].

195 **5. CONCLUSION**

196 In conclusion, in this study investigating the effects of high fructose administration on the
197 liver and kidney, we determined that fructose increased inflammation by regulating the TNF-
198 α and IL-6 signaling pathway as a result of NF- κ B activation.

199

200 **REFERENCES**

- 201 1. Rebollo A, Roglans N, Alegret M. Way back for fructose and liver metabolism: Bench side
202 to molecular insights. *World J Gastroenterol* 2012;18(45):6552-6559.
- 203 2. Bellamkonda R, Karuna R, Sasi Bhusana Rao B, Haritha K, Manjunatha B, Silpa S,
204 Saralakumari D. Beneficiary effect of Commiphora mukul ethanolic extract against high
205 fructose diet induced abnormalities in carbohydrate and lipid metabolism in wistar rats. *J*
206 *Tradit Complement Med* 2017;8(1):203-211.
- 207 3. Nomura K, Yamanouchi T. The role of fructose-enriched diets in mechanisms of
208 nonalcoholic fatty liver disease. *J Nutr Biochem* 2012;23:203-208.
- 209 4. Park JH, Ku HJ, Kim JK, Park JW, Lee JH. Amelioration of High Fructose-Induced
210 Cardiac Hypertrophy by Naringin. *Sci. Rep* 2018;8(1): 9464.
- 211 5. Abbas AK, Lichtman AH. Basic immunology: functions and disorders of the immune
212 system. 2nd ed. Philadelphia, PA: W.B Saunders Co, Updated edition 2006-2007.
- 213 6. Avci B, Alacam H, Dilek A, Kozan A. Effects of asymmetric dimethylarginine on
214 inflammatory cytokines (TNF- α , IL-6 and IL-10) in rats. *Toxicol. Ind. Health*
215 2015;31(3):268-73.
- 216 7. Chan DC, Watts GF, Ng TW, Uchida Y, Sakai N, Yamashita S, Barrett PH. Adiponectin
217 and other adipocytokines as predictors of markers of triglyceride-rich lipoprotein metabolism.
218 *Clin Chem* 2005;51(3):578-85.
219
- 220 8. Tsatsanis C, Zacharioudaki V, Androulidaki A, Dermitzaki E, Charalampopoulos I, Minas
221 V, Gravanis A, Margioris AN. Adiponectin induces TNF-alpha and IL-6 in macrophages and
222 promotes tolerance to itself and other pro-inflammatory stimuli. *Biochem Biophys Res*
223 *Commun* 2005;335(4):1254-63.
224
- 225 9. Francini F, Castro MC, Schinella G, García ME, Maiztegui B, Raschia MA, Gagliardino
226 JJ, Massa ML. Changes induced by a fructose-rich diet on hepatic metabolism and the
227 antioxidant system. *Life Sci* 2010;86:965-971.
228
- 229 10. Gilmore TD. Introduction to NF-kappaB: players, pathways, perspectives. *Oncogene*
230 2006;25:6680-4.
231
- 232 11. Baker RG, Hayden MS, Ghosh S. NF- κ B, inflammation and metabolic disease. *Cell*
233 *Metab.* 2011;13(1):11-22.
234
- 235 12. Sanz AB, Sanchez-Niño MD, Ramos AM, Moreno JA, Santamaria B, Ruiz-Ortega M,
236 Egido J, Ortiz A. NF-kappaB in renal inflammation. *J Am Soc Nephrol* 2010;21:1254-1262.
237
- 238 13. Volynets V, Spruss A, Kanuri G, Wagnerberger S, Bischoff SC, Bergheim I. Protective
239 effect of bile acids on the onset of fructose-induced hepatic steatosis in mice. *J Lipid Res*
240 2010;51:3414-3424.
241
- 242 14. Havel PJ. Dietary fructose: implications for dysregulation of energy homeostasis and
243 lipid/carbohydrate metabolism. *Nutr Rev* 2005;63(5):133-57.
244

- 245 15. Bray GA, Nielsen SJ, Popkin BM. Consumption of high-fructose corn syrup in beverages
246 may play a role in the epidemic of obesity. *Am J Clin Nutr* 2004;79(4):537-43.
247
- 248 16. Friedman SL. Seminars in medicine of the Beth Israel Hospital, Boston. The cellular basis
249 of hepatic fibrosis. Mechanisms and treatment strategies. *N Engl J Med* 1993;328:1828-1835.
250
- 251 17. Nieto N, Friedman SL, Cederbaum AI. Stimulation and proliferation of primary rat
252 hepatic stellate cells by cytochrome P450 2E1-derived reactive oxygen species. *Hepatology*
253 2002;35:62-73.
254
- 255 18. Malaguarnera L, Di Rosa M, Zambito AM, dell'Ombra N, Nicoletti F, Malaguarnera M.
256 Chitotriosidase gene expression in Kupffer cells from patients with non-alcoholic fatty liver
257 disease. *Gut* 2006;55:1313-1320.
- 258 19. Malaguarnera L, Di Rosa M, Zambito AM, dell'Ombra N, Di Marco R, Malaguarnera M.
259 Potential role of chitotriosidase gene in nonalcoholic fatty liver disease evolution. *Am J*
260 *Gastroenterol* 2006;101:2060-2069.
- 261 20. Masterjohn C, Park Y, Lee, J. Dietary fructose feeding increases adipose methylglyoxal
262 accumulation in rats in association with low expression and activity of glyoxalase-2.
263 *Nutrients* 2013;5:3311-3328.
264
- 265 21. Giriş M, Abbasoğlu SD, Kumral A. Effect of carnosine alone or combined with α -
266 tocopherol on hepatic steatosis and oxidative stress in fructose-induced insulin-resistant rats. *J*
267 *Physiol Biochem* 2014;70:385-395.
268
- 269 22. Ackerman Z, Herman MO, Rosenthal MGT, Pappo O, Link G, Sela BA. Fructose-
270 induced fatty liver disease hepatic effects of blood pressure and plasma triglyceride
271 reduction. *Hypertension* 2005;45(5):1012-8.
- 272 23. Falasca K, Ucciferri C, Dalessandro M, Zingariello P, Mancino P, Petrarca C, Pizzigallo
273 E, Conti P, Vecchiet J. Cytokine patterns correlate with liver damage in patients with chronic
274 hepatitis B and C. *Ann Clin Lab Sci* 2006;36(2):144-150
- 275 24. Jones SA. Directing transition from innate to acquired immunity: defining a role for IL-6.
276 *J Immunol* 2005;175(6):3463-3468.
- 277 25. Schmidt-Arras D, Rose-John S. IL-6 pathway in the liver: From physiopathology to
278 therapy. *J Hepatol.* 2016;64(6):1403-15.
- 279 26. Yang M, Jiang R, Lin X. Ginger extracts improve renal injury of high fructose-fed SD
280 rats by inhibiting the expression of proinflammatory cytokines. *Xi Bao Yu Fen Zi Mian Yi*
281 *Xue Za Zhi.* 2014;30(9):929-32.
- 282 27. Gersch MS, Mu W, Cirillo P, Reungjui S, Zhang L, Roncal C, Sautin YY, Johnson RJ,
283 Nakagawa T. Fructose, but not dextrose, accelerates the progression of chronic kidney
284 disease. *Am J Physiol Renal Physiol* 2007;293:1256-1261.
- 285 28. Stanhope KL, Schwarz JM, Keim NL, Griffen SC, Bremer AA, Graham JL, Hatcher B,
286 Cox CL, Dyachenko A, Zhang W. Consuming fructose-sweetened, not glucose-sweetened,
287 beverages increases visceral adiposity and lipids and decreases insulin sensitivity in
288 overweight/obese humans. *J Clin Investig* 2009;119:1322-1334.

- 289 29. Pektas MB, Koca HB, Sadi G, Akar F. Dietary fructose activates insulin signaling and
290 inflammation in adipose tissue: Modulatory role of resveratrol. *BioMed Res Int*
291 2016;2016:8014252.
- 292 30. Zhang DM, Jiao RQ, Kong LD. High Dietary Fructose: Direct or Indirect Dangerous
293 Factors Disturbing Tissue and Organ Functions. *Nutrients* 2017; 9:1-25.
- 294 31. Jin YH, Park YJ, Ham M, Kim JB. Crosstalk between adipocytes and immune cells in
295 adipose tissue inflammation and metabolic dysregulation in obesity. *Mol Cells* 2014;37:365-
296 371.
- 297 32. Weltman MD, Farrell GC, Liddle C. Increased hepatocyte CYP2E1 expression in a rat
298 nutritional model of hepatic steatosis with inflammation. *Gastroenterol* 1996;111:1645-1653.
- 299 33. George DK, Goldwurm S, MacDonald GA, Cowley LL, Walker NI, Ward PJ, Jazwinska
300 EC, Powell LW. Increased hepatic iron concentration in nonalcoholic steatohepatitis is
301 associated with increased fibrosis. *Gastroenterol* 1998;114:311-318.
- 302 34. Veličković N, Djordjevic A, Vasiljević A, Bursać B, Milutinović DV, Matic G. Tissue-
303 specific regulation of inflammation by macrophage migration inhibitory factor
304 and glucocorticoids in fructose-fed Wistar rats. *Br J Nutr* 2013;110(3):456–465.
- 305 35. Zheng J, Peng C, Ai Y, Wang H, Xiao X, Li J. Docosahexaenoic Acid Ameliorates
306 Fructose-Induced Hepatic Steatosis Involving ER Stress Response in Primary Mouse
307 Hepatocytes. *Nutrients* 2016;8(1):55.
308
- 309 36. Vasiljević A, Bursać B, Djordjevic A, Milutinović DV, Nikolić M, Matic G, Veličković
310 N. Hepatic inflammation induced by high-fructose diet is associated with altered 11 β HSD1
311 expression in the liver of Wistar rats. *Eur J Nutr* 2014;53(6):1393-402.
312
- 313 37. Basaranoglu M, Basaranoglu G, Sabuncu T, Sentürk H. Fructose as a key player in the
314 development of fatty liver disease. *World J Gastroenterol* 2013;19(8):1166-72.
315
- 316 38. Gavrilova O, Haluzik M, Matsusue K, Cutson JJ, Johnson L, Dietz KR, Nicol CJ,
317 Vinson C, Gonzalez FJ, Reitman ML. Liver peroxisome proliferator-activated receptor
318 gamma contributes to hepatic steatosis, triglyceride clearance, and regulation of body fat
319 mass. *J Biol Chem* 2003;278(36):34268-76.
320
- 321 39. Sunami Y, Leithäuser F, Gul S, Fiedler K, Güldiken N, Espenlaub S. Hepatic activation
322 of IKK/NF-kappa B signaling induces liver fibrosis via macrophage-mediated chronic
323 inflammation. *Hepatology* 2012;56(3):1117-28.
324
- 325
- 326
- 327
- 328
- 329

330 **Table 1.** Serum biochemical parameters of the groups

	Control	Fructose
AST (U/L)	45,00±15,15	150,00±20,50 ^a
ALT (U/L)	34,00±12,50	172,00±23,34 ^a
ALP (U/L)	120,45±22,35	269,35±40,85 ^a
BUN (mg/dL)	16,45±3,12	58,35±5,22 ^a
Uric acid (mg/dL)	4.20±0,80	17,34±4,62 ^a

331 Data were given as mean ±SD.

332 AST : Aspartate aminotransferase, ALT : Alanine aminotransferase, ALP: Alkaline phosphatase,

333 BUN: Blood Urea Nitrogen

334

335 ^aSignificantly different when compared with control group, (p<0.001)

336

337 **Table 2.** Tissue biochemical parameters of the groups

	Groups	TNF- α	IL-6	NF- κ B
Liver	Control	1160,75±476,33	1,03±0,26	0,89±0,28
	Fructose	3510,64±400,37 ^a	1,75±0,32 ^b	2,62±0,57 ^a
Kidney	Control	203,08±26,54	0,16±0,06	0,36±0,08
	Fructose	528,59±101,43 ^a	0,37±0,10 ^b	0,81±0,20 ^a

338 Data were given as mean ±SD.

339 TNF- α : Tumor necrosis factor alpha, IL-6: Interleukin-6, NF- κ B: Nuclear factor kappa B

340 ^aSignificantly different when compared with control group, (p<0.001)

341 ^bSignificantly different when compared with control group, (p<0.05)

342 ^cSignificantly different when compared with control group, (p<0.01)

343 **Highlights**

344

345 • Fructose fed diet causes a potentially fatal hepatotoxicity and nephrotoxicity

346

347 • Levels of NF- κ B, a transcription factor involved in the expression of cytokines that
348 cause proinflammatory cytokine elevation, were measured in both liver and kidney
349 tissues.

350

351 • Fructose increased inflammation by regulating the TNF- α and IL-6 signaling pathway
352 as a result of NF- κ B activation.