

Original Research Article

Factors contributing to delayed breast cancer presentation: A prospective study at Parirenyatwa group of hospitals, Harare, Zimbabwe 2010-2013.

Abstract

Background: Breast cancer is one of the most common female cancers in Zimbabwe. A considerable proportion of patients delay presentation, leading to high morbidity and mortality. Delay in presentation can either be provider or patient delay. Survival is related to the stage at presentation. Delayed presentation is associated with lower survival. Understanding the reasons for delay may help in reducing delays and morbidity and mortality. This study addresses these concerns.

Aim: To determine factors contributing to delayed breast cancer presentation at Parirenyatwa Group of Hospitals

Methods: A prospective observational study of patients with the clinical and histological diagnosis of breast cancer attending Surgical Outpatient clinics awaiting surgery, or operated on from January 2010 to December 2013 were included. Patients were interviewed and specific questions relating to breast cancer risk and delay factors were recorded. Relevant investigations, including Human Immune Deficiency Virus (HIV) testing, were done and recorded. Final histology results were collected from Histopathology Department, analyzed and recorded. In addition to chi-square test for associated factors of delay and proportionate z test for percentage differences, the researchers validated the observed factors using discriminant analysis. Discriminant analysis was used to model the reasons and delay period with a cut-off point 3 months (< 3 months / ≥ 3 months).

Results: Seventy three patients were enrolled in the study. Forty nine (62.1%) were of rural domicile. Time to breast cancer presentation ranged from 1 to 52 months. The most common reason for delay (66%) was ignorance and the secondly (18%) poverty. Fifty three (72.6%) patients were unemployed ($p < 0.05$). Primary school was the highest level of education in 23 patients (31.5%), with 38 (52.1%) having attained

34 secondary level education. Fifty-seven (78.1%) patients presented with a mass
35 ($p < 0.05$) with pain occurring in 29 (39.7%) of patients. Fifty four patients (74%) had no
36 knowledge of self-breast-breast self-examination (BSE) and 37 (51%) of these patients were
37 of rural domicile ($p < 0.05$). Of the 37 rural patients with no knowledge of BSEself-breast-ex-
38 amination-35 (94.5%), had primary level education ($p < 0.005$). Fifty one (69.9%) patients
39 consented to HIV testing, 7 (13.7%) were HIV positive. A low-level of education, ignorance
40 of breast cancer, poor socio-economic status, rural residence and lack of knowledge of
41 BSEbreast self-examination (BSE)-were important predictors of breast cancer -delay to
42 presentation. old-Old age, HIV status, level of education and family history were major
43 reasons associated with breast cancer presentation delay.

44
45 **Conclusion:** The overwhelming majority of breast cancer patients attending Parire-
46 nyatwa Group of Hospitals presented with advanced disease. These patients were
47 mostly of low socio-economic status. Current health education campaigns seem to
48 be ineffective in improving breast cancer awareness. Strategies to reduce delays in
49 presentation, through various interventions focused on education and poverty allev-
50 iation need to be formulated.

51
52 **Key words:** breast cancer, presentation, delay, factors, developing countries

53 54 Introduction

55 -Breast cancer is the most common malignancy in females worldwide. It is the lead-
56 ing cause of cancer related mortality¹. Over 1-2 million cases are diagnosed every
57 year, affecting 10 to 12% of the female population, and accounting for more than
58 500,000 deaths per year worldwide^{2,3}. The Zimbabwe National Cancer Registry
59 2012 Report³ highlighted that 11% of cancer deaths were due to breast cancer, with
60 an incidence of 7%. Breast cancer mostly affects women and only a very small per-
61 centage of men.^{2,3} Factors contributing to delayed breast cancer presentation have
62 been studied elsewhere but not in Zimbabwe, despite the large number of deaths
63 due to breast cancer.

64
65 Patients who present late (figures 1-3) have lower survival rates⁴. An association
66 between stage at diagnosis and survival has been established⁴. Delayed patient
67 presentation refers to a prolonged interval between the discovery of initial symptoms

Formatted: Justified

68 and evaluation by a service provider. Delayed presentation is typically defined as an
69 interval greater than 12 weeks ⁵. ~~provider~~ Provider delay is when patients are re-
70 ferred late. This could either be due to wrong diagnoses being made or to failures in
71 the referral system, as commonly experienced in developing countries like Zim-
72 babwe. In Zimbabwe general medical practitioners and local clinics refer cases of
73 breast cancer directly to central hospitals. A proportion of patients are delayed at this
74 level. In provider delay, patients who present early are managed late thereby wor-
75 sening their outcome. In patient delay, for various reasons patients procrastinate so
76 by the time they seek medical help, the disease may be advanced. Patient delay
77 plays a major role in breast cancer related morbidity and mortality ⁵. Patients with
78 delays of 3 to 6 months have worse survival rates than those with delays of less than
79 3 months ⁶.

80
81 During the patient delay process ⁶⁻¹⁰, the time from the individual detecting the symp-
82 tom until they seek medical attention is termed "appraisal delay" ⁷ or "passive detec-
83 tion"⁸. The time from the individual recognizing the symptom to seeking help is called
84 "action appraisal" ⁹, or behavioral delay⁷. Negative attitudes towards healthcare pro-
85 viders are among the determinants of behavioral delay ¹⁰⁻²⁰. Knowledge of breast
86 cancer symptoms and self-breast examination (BSE) have been associated with less
87 appraisal and behavioral delays ^{8, 12, 13, 20-30}. Patient delay may be related to poor
88 socioeconomic status, cultural beliefs, and level of education, ignorance and acces-
89 sibility to healthcare facilities ^{14, 22, 30-40} among other factors.

90
91 The Zimbabwe National Cancer Registry (2012) report showed on average 1,-800
92 women are affected annually by breast cancer. Approximately 1,200 die from this
93 disease annually.^{2, 3} In Zimbabwe, breast cancer affects one in every 10 women.^{3, 41-}
94 ⁴⁶ This study was carried out to provide scientific data on factors associated with de-
95 layed breast cancer presentation in Zimbabwe. The aim was to identify possible
96 strategies to shorten these delays thus reducing breast cancer mortality in Zim-
97 babwe.

98
99 **AIM:** This study aimed to determine the factors associated with delay to breast can-
100 cer presentation

101

102 **Objectives:**

103
104 To determine the magnitude and reasons for delayed breast cancer presentation at
105 Parirenyatwa Group of Hospitals.

106 To determine any association between level of education and delay in presentation.

107 To determine the stage at presentation of breast cancer.

108 To determine the presenting symptoms.

109 To determine any association between HIV infection and advanced breast cancer.

110

111 **Study design:** A prospective observational study

112

113 **Sampling Procedure and Sample Size**

114

115 **Sample Size Estimation**

116 The minimum sample size n was obtained using the formula developed by Cochran
117 (in year 2006) which is -was used in populations that are large:

118

119
$$n = \frac{z^2 p(1 - p)}{\epsilon^2}$$

120 Where,

121 p = Proportion of breast cancer patients who delayed for more than three months, p
122 = 94%, calculated from a proportion of breast cancer patients delayed for more than
123 three months in a study done by Muguti *et al.*, (1993)⁴⁶ in Zimbabwe

124 ϵ = margin of error set at 6 %

125 Z= standard normal deviate set at 1.96 for 95% confidence level

126 n= pPopulation size = 61

127

128

129 **Materials and Methods**

130 All patients with a clinical and histological diagnosis of breast cancer attending Sur-
131 gical Outpatient Department clinics, admitted, awaiting surgery or operated on from
132 January 2010 to December 2013 were included in the study. Patients were inter-
133 viewed and specific questions relating to breast cancer risk and delay factors rec-

134 orded. Relevant investigations including HIV testing were done and recorded. Final
135 histology results were collected analyzed and recorded. Delayed patient presentation
136 was defined as a prolonged interval between the discovery of the initial symptom to
137 presentation to a provider, typically greater than 12 weeks (3 months).^{5,21,22} Discrimi-
138 nant analysis was used to model delay period with a cut-off point 3 months (< 3
139 months / ≥ 3 months).

140
141 **Inclusion Criteria:**
142 All female patients with a clinical and histological diagnosis of breast cancer over 15
143 years age attending clinics or admitted to Parirenyatwa University Teaching Hospital

144
145 **Exclusion Criteria:**
146 Male patients with breast cancer
147 Patients with breast cancer <15 years
148 Patients who did not have histological confirmation of breast cancer

149
150 **Statistical analysis**
151 All data was entered in Epidata Entry version 3.1 software and cleaned before anal-
152 ysis. Statistical analysis was carried out by SPSS version 16 statistical package.
153 Discriminant analysis was used to model the reasons for delay in months. Descrip-
154 tive statistics: means, standard deviations, canonical discriminant parameters were
155 determined as discriminant analysis procedure. The significance levels used to indi-
156 cate effect size were $p < 0.05$.

157
158 **Model validation**
159 Among other diagnostics parameters used were Wilk's lambda (preferred the smal-
160 lest value), and Box's M. We used a 50% Bernoulli (0.5) random sampling of the 73
161 patients to create a discriminant analysis model, setting the remaining (50%) patients
162 aside to validate the analysis. We then used the model to classify the 50% of the pa-
163 tients as delayed or not delayed. Checking for other assumptions see table 5

164
165 **Ethics statement**

166 Ethical approval was sought from Parirenyatwa and College of Health Sciences Joint
167 Research (JREC). Written consent to participate in the study and publish pictures
168 was obtained

169 170 **Conflict of Interest**

171 **The authors declare no conflict of interest.** The study was self-funded.

172 173 **Results**

174 175 **Descriptive analysis**

176 In this study ~~of out~~ 73 patients, 53 (72.6%) ~~patients~~ presented with advanced breast
177 cancer, 23 (31.5%) were ~~in~~ stage 3 and 30 (41.1%) were ~~in~~ stage 4. Forty-three pa-
178 tients (59%) self-delayed in seeking breast cancer treatment whilst only 30 (41%)
179 were treated within the recommended period (within 3 months from the first symptom
180 onset) ^{5, 21, 22}. ~~Out of 73 patients, most patients 37 (50.7%), p = 0.05 (insignificant)~~
181 ~~with advanced breast cancer (stage 3 to 4) were from rural area compared to 16~~
182 ~~(21.9%) from urban.~~ Of the 73 study patients, 49 (67.1%) were of rural -and 24
183 (32.9%) urban domicile (figure 6). ~~Thirty-seven of the 53 patients with stage 3 and 4~~
184 ~~disease were from rural areas and 16 were urban, p=0.05.~~ Time to breast cancer
185 presentation ranged from 1 to 52 months. The most common reason for delay (48
186 patients, 66%) was ignorance and secondly poverty (13 patients, 18%). Other rea-
187 sons such as unemployment were not associated with delay ($p > 0.05$), table 6. Pa-
188 tients whose highest level of education was primary education were 23 (31.5%) and
189 38 (52.1%) had secondary level education as their highest level (figure 8). The pre-
190 senting symptom in 57 (78.1%) patients was a mass ($p < 0.05$) and pain occurred in
191 39.7% of patients (Table 5). Knowledge of ~~BSE self-breast cancer examination~~ was
192 associated with level of education (shown in table 7). ~~Fifty-four Table 2 show that 54~~
193 ~~(74%) of patients had no knowledge of self-breast examination BSE and 37 (68.5%)~~
194 ~~of these patients were of rural domicile, $p < 0.05$, thus there was a significant relation-~~
195 ~~ship ($p < 0.05$).~~ Of the 37 rural patients with no knowledge of self-breast examina-
196 tion ~~BSE~~, 35 (94.6%) ~~patients~~ had only a primary education ($p < 0.005$), ~~significant re-~~
197 ~~lationship.~~ ~~Most~~ Generally more patients, 20 (27.4%), were within an age range of 51-

Comment [A1]: Need to include other stages- namely 8 patients with stage 1 and 12 stage 2 disease. Then can eliminate figure 4.

Comment [A2]: Figure 5 can be eliminated as now in text

Comment [A3]: This doesn't make grammatical sense. Should be added after next sentence.

Comment [A4]: Table 6 doesn't add anything to paper. Insignificant. Table 6 only looks at unemployment- what other factors were not associated

Comment [A5]: Grammar incorrect. Should read "Twenty-three (31.5%) of patients attained a primary school education and 38 (52.1%) went to secondary school" or something similar.

Comment [A6]: In text so do not need table 2

Comment [A7]: This is noted table 1 and 3 which are the same! Neither is referenced here.

198 60 years and followed by 15 (20.5%), aged between 41-50 years (figure 7). Fifty-one
199 patients (69.9%) consented to HIV testing, of which 7 (13.7%) were positive.

Formatted: Justified

202 Discriminant analysis

204 HIV positive status and a low level of education or ignorance (“a lack of knowledge,
205 understanding, or education”) are among the main reasons for breast cancer treat-
206 ment delay (table 8). ~~The Webster's Learner's Dictionary defines ignorance as “a~~
207 ~~lack of knowledge, understanding, or education”.~~ The findings in Figure 9 therefore
208 confirms that lack of education tops the indicated reasons. Thus ignorance or “a lack
209 of knowledge, understanding, or education” is another major reason of breast cancer
210 presentation delay.

Comment [A8]: Put reference to Webster's dictionary here so can eliminate next sentence

Comment [A9]: This could be better stated as “The lack of education being the main indicator”
Figure 9 could be put as an additional column in figure 8.

212 In standardized factor mean scores and standard deviations, the higher the mean
213 score the greater the factor contributes in categorizing the dependent variable. Small
214 standard deviations are preferred. Observations were categorized by presentation as
215 “delayed” (1) and “not delayed” (0).

Comment [A10]: This should be known by the reader.

Formatted: Justified

217 Coefficients with large absolute values correspond to variables with greater discrimi-
218 nating ability as factors associated with patients who had delayed presentation
219 ,namely old age (Coefficient; 1.061), HIV status (Coefficient; 0.89), level of education
220 (Coefficient; 0.679), and family history (Coefficient; 0.221) (table 10)

222 Discussion

224 Breast cancer is a common health problem in our environment and patients present
225 late. Factors causing delayed presentation are both patient and system related. In
226 our study the major reasons for patient delay were old age, HIV status, and low level
227 of education. In this study 43 (59%) of delays were patient related. This correlates
228 with other studies which looked at reasons for patient delay^{6, 17, 30}. A large proportion
229 of our patients were of low socioeconomic background and had low the least educa-
230 tional backgrounds. Knowledge of self breast examination BSE is was lacking. It is

231 recommended that campaigns must be directed at this population group with a view
232 to provide education regarding the early signs and symptoms of breast cancer so as
233 to change and improve their health seeking behavior ^{8, 12, 13, 14-22}. Burgess et al con-
234 cluded in their study that patients presenting late had competing demands and priori-
235 ties, fears about cancer treatments and anxieties about 'bothering the doctor'¹¹.
236 These psychosocial factors were noted in our study and need to be addressed in
237 health education campaign programmes. Although only small percentage of patients
238 were HIV positive, the majority of these presented with advanced breast cancer. The
239 stigma associated with HIV is a risk factor for delayed presentation ²⁰. This corre-
240 lates with Brazilian studies ^{40, 41, and 42}, one study reviewed breast cancer in a cohort
241 of HIV infected women. The median age at diagnosis was 46 years. The median
242 survival after breast cancer diagnosis was 12 months and breast cancer diagnosis
243 was made within 2 to 15 years of HIV-infection diagnosis. All patients were diag-
244 nosed late with breast cancer and thus had a worse prognosis ^{40, 41, 42}.

245
246 Most **B**reast cancer patients attending Parirenyatwa Group of Hospitals present
247 with advanced disease. Current health education campaigns seem not to be inter-
248 disciplinary and effective in improving breast cancer awareness; People living with
249 HIV are suffering stigma and eventually delay due to low self-esteem ²⁰. It is our col-
250 lective responsibility to reduce this delay through various interventions focused on
251 education and poverty alleviation. Follow-up studies regarding management of these
252 patients need to be done so as to recommend and formulate local guidelines

253
254 **Conclusion**
255 Factors causing delayed presentation are both patient and system related. In our
256 study the major reasons for delay were older age, HIV status, and low level of edu-
257 cation respectively. Most were patient delays with low socio-economic background
258 and low educational back-ground. Knowledge of ~~self breast examination~~BSE is lack-
259 ing. Education campaigns must be directed at this population group with a view to
260 provide education regarding the early signs and symptoms of breast cancer so as to
261 change and improve their health seeking behavior. The majority of HIV-positive pa-
262 tients presented with advanced breast cancer and HIV stigma was a risk factor for
263 delayed presentation.

Formatted: Justified

264 |
265 | Current health education campaigns seem not to be inter-disciplinary and effective in
266 | improving breast cancer awareness; People living with HIV suffering stigma and
267 | eventually delay due to low self-esteem. It is our collective responsibility to reduce
268 | this delay through various interventions focused on education and poverty allevia-
269 | tion.

270 |
271 | **Recommendations**
272 | Focused public health campaigns aimed at raising breast cancer awareness must
273 | target rural communities. ~~Breast self-~~Self-breast examination must be taught to
274 | women at all levels. Rural communities need to be encouraged to advance their
275 | education levels. Communities need to be empowered economically in order to im-
276 | prove their health seeking behaviour with special emphasis on breast cancer. Pa-
277 | tients presenting late have competing demands and priorities, fears about cancer
278 | treatments and anxieties about 'bothering the doctor. These psychosocial factors
279 | need to be addressed in health education campaign programs. Follow-up studies re-
280 | garding management of these patients need to be done so as to recommend and
281 | formulate local guidelines

282 |
283 |
284 |

285 | **References**
286 |
287 | 1. Hortobagyi GN, dela Garza Salazar J, et al. The global breast cancer burden:
288 | variations in epidemiology and survival. *Clin Breast Cancer* (2005); **6**:391-401.
289 | 2. Benson JR, Jatoi I, Keisch M. Early breast cancer. *Lancet* (2009); **373**:1642-3.
290 | 3. Chokunonga E, et al. Zimbabwe Cancer Registry 2012 Annual Report. African
291 | Cancer Registry Network. [afcrn.org/membership/membership-list/83-](http://afcrn.org/membership/membership-list/83-zimbabwe-harare)
292 | [zimbabwe-harare](http://afcrn.org/membership/membership-list/83-zimbabwe-harare). Accessed February 23, 2016.
293 | 4. Thomson CS, Forman D. Cancer survival in England and the influence of ear-
294 | ly diagnosis. *Br J Cancer* (2009);**101**:S102-9
295 | 5. Ramirez A.J. , Westcombe A. M., Burgess C. C., Sutton S., Littlejohns P., and
296 | Richards M. A., "Factors predicting delayed presentation of symptomatic

- 297 breast cancer: a systematic review," *The Lancet*, vol. 353, no. 9159,(1999).pp.
298 1127–1131.
- 299 6. Facione NC. Delay versus help seeking for breast cancer symptoms: a critical
300 review of the literature on patient and provider delay. *Soc Sci Med*
301 (1993);36:1521-34.
- 302 7. Richards MA, Westcombe AM, Love SB, Littlejohns P, Ramirez AJ. The influ-
303 ence of delay on survival in patients with breast cancer: a systematic review.
304 *Lancet* (1999);353:1119-1126.
- 305 8. Andersen BL, Cacioppo JT. Delay in seeking a cancer diagnosis: delay stag-
306 es and psychological comparison process. *Br J Soc Psychol*(1995);34:33-52.
- 307 9. Ruiter RA, de Nooijer J, van Breukelen G, Ockhuysen- VermeyCF, de Vries
308 H. Intended coping responses to cancer symptoms in healthy adults: the roles
309 of symptom knowledge, detection behavior, and perceived threat. *Cancer Ep-*
310 *idemiol Biomarkers Prev* (2008);17:818-26.
- 311 10. Ristvedt SL, Trinkaus KM. Psychological factors related to delay in consulta-
312 tion for cancer symptoms. *Psychooncology* (2005);14:339-50.
- 313 11. Burgess CC, Potts HW, Hamed H, et al. Why do older woman delay presenta-
314 tion with breast cancer symptoms? *Psychooncology* (2006);15:962-8.
- 315 12. Scott SE, Grunfeld EA, Auyeung V, Mcgurk M. Barriers and triggers to seek-
316 ing help for potentially malignant oral symptoms: implications of interventions.
317 *J Public Health Dent* (2009);69:34-40
- 318 13. De Nooijer J, Lechner L, de Vries H. Social psychological correlates of paying
319 attention to cancer symptoms and seeking medical help. *Soc Sci Med*
320 (2003);56:915-20
- 321 14. Van Osch L, Lechner L, Reubsaet A, de Nooijer J, de Vries H. Passive cancer
322 detection and medical help seeking for cancer symptoms:inadequate behavior
323 and psychological determinants. *Eur J Cancer Prev* (2007);16:266-74
- 324 15. Hunter CP, Redmond CK, Chen VW, et al. Breast Cancer factors associated
325 with stage at diagnosis in black and white woman. Black/ White Cancer Sur-
326 vival Study Group. *J Natl Cancer Inst* (1993);85:1129-37
- 327 16. Macleod U, Mitchell ED, Burgess C, Macdonald S and Ramirez AJ; Risk fac-
328 tors for delayed presentation and referral of symptomatic cancer, evidence for
329 common cancers. *British Journal of Cancer* (2011) 101, S92 – S101,pg1

- 330 17. Harirchi I, Ghaemmaghami F, Karbakhsh M, et al. Patient delay in women
331 presenting with advanced breast cancer: an Iranian study (2005). *Public*
332 *Health*, 119, 885-91.
- 333 18. Montazeri A, Ebrahimi M, Mehrdad N, et al. Delayed presentation in breast
334 cancer: a study in Iranian women (2003). *BMC Womens Health*, p3, 4.
- 335 19. Ramirez AJ, Westcombe AM, Burgess CC, et al. Factors predicting delayed
336 presentation of symptomatic breast cancer: a systematic review
337 (1999). *Lancet*, **353**, 1127-31.
- 338 20. Zimbabwe People Living with HIV Stigma Index Report
339 (2014); [http://www.stigmaindex.org/sites/default/files/reports/Zimbabwe%20Pe-](http://www.stigmaindex.org/sites/default/files/reports/Zimbabwe%20People%20Living%20with%20HIV%20Stigma%20Index%20Report_15-12-14pdf.pdf)
340 [ople%20Living%20with%20HIV%20Stigma%20Index%20Report_15-12-](http://www.stigmaindex.org/sites/default/files/reports/Zimbabwe%20People%20Living%20with%20HIV%20Stigma%20Index%20Report_15-12-14pdf.pdf)
341 [14pdf.pdf](http://www.stigmaindex.org/sites/default/files/reports/Zimbabwe%20People%20Living%20with%20HIV%20Stigma%20Index%20Report_15-12-14pdf.pdf); accessed 10/28/2016:12:29pm
- 342 21. Lancaster JM, Powell CB, Chen LM, et al. (2015): Society of Gynecologic On-
343 cology statement on risk assessment for inherited gynecologic cancer predis-
344 positions. *Gynecol Oncol* 136 (1): 3-7, [PUBMED Abstract]
- 345 22. Robson ME, Bradbury AR, Arun B, et al. (2015): American Society of Clinical
346 Oncology Policy Statement Update: Genetic and Genomic Testing for Cancer
347 Susceptibility. *J Clin Oncol* 33 (31): 3660-7, [PUBMED Abstract]
- 348 23. (n.d.). In Merriam-Webster's online dictionary (11th ed.). Retrieved from
349 <http://www.merriam-webster.com/dictionary/ignorance>; accessed on
350 1/11/2016, 6:05am.
- 351 24. Hortobagyi GN, dela Garza Salazar J, et al. The global breast cancer burden:
352 variations in epidemiology and survival. *Clin Breast Cancer* 2005; **6**:391-401.
- 353 25. Benson JR, Jatoi I, Keisch M. Early breast cancer. *Lancet* 2009; **373**:1642-3.
- 354 26. Zimbabwe Cancer Registry 2010 statistics
- 355 27. Thomson CS, Forman D. Cancer survival in England and the influence of ear-
356 ly diagnosis. *Br J Cancer* 2009; **101**:S102-9
- 357 28. Facione NC. Delay versus help seeking for breast cancer symptoms: a critical
358 review of the literature on patient and provider delay. *Soc Sci Med*
359 1993; **36**:1521-34.
- 360 29. Richards MA, Westcombe AM, Love SB, Littlejohns P, Ramirez AJ. The influ-
361 ence of delay on survival in patients with breast cancer: a systematic review.
362 *Lancet* 1999; **353**:1119-1126.

- 363 30. Andersen BL, Cacioppo JT. Delay in seeking a cancer diagnosis: delay stag-
364 es and psychological comparison process. *Br J Soc Psychol*1995;34:33-52.
- 365 31. Ruiter RA, de Nooijer J, van Breukelen G, Ockhuysen- VermeyCF, de Vries
366 H. Intended coping responses to cancer symptoms in healthy adults: the roles
367 of symptom knowledge, detection behavior, and perceived threat. *Cancer Ep-
368 idemiol Biomarkers Prev* 2008;17:818-26.
- 369 32. Ristvedt SL, Trinkaus KM. Psychological factors related to delay in consulta-
370 tion for cancer symptoms. *Psychooncology* 2005;14:339-50.
- 371 33. Burgess CC, Potts HW, Hamed H, et al. Why do older woman delay presen-
372 tation with breast cancer symptoms? *Psychooncology* 2006;15:962-8.
- 373 34. Scott SE, Grunfeld EA, Auyeung V, Mcgurk M. Barriers and triggers to seek-
374 ing help for potentially malignant oral symptoms: implications of interventions.
375 *J Public Health Dent* 2009;69:34-40
- 376 35. de Nooijer J, Lechner L, de Vries H. Social psychological correlates of paying
377 attention to cancer symptoms and seeking medical help. *Soc Sci Med*
378 2003;56:915-20
- 379 36. van Osch L, Lechner L, Reubsaet A, de Nooijer J, de Vries H. Passive cancer
380 detection and medical help seeking for cancer symptoms:inadequate behavior
381 and psychological determinants. *Eur J Cancer Prev* 2007;16:266-74
- 382 37. Hunter CP, Redmond CK, Chen VW, et al. Breast Cancer: factors associated
383 with stage at diagnosis in black and white woman. Black/ White Cancer Sur-
384 vival Study Group. *J Natl Cancer Inst* 1993;85:1129-37
- 385 38. Pace, L. E., Mpunga, T., Hategekimana, V., Dusengimana, J.-M. V., Habine-
386 za, H., Bigirimana, J. B., ... Keating, N. L. (2015). Delays in Breast Cancer
387 Presentation and Diagnosis at Two Rural Cancer Referral Centers in Rwanda.
388 *The Oncologist*, 20(7), 780–788. [http://doi.org/10.1634/theoncologist.2014-
389 0493](http://doi.org/10.1634/theoncologist.2014-0493)
- 390 39. Borges *et al*, (2013). Predicting risk of cancer during HIV infection: the role of
391 inflammatory and coagulation biomarkers. *AIDS* 27(9).
- 392 40. De Andrade AC, Luz PM, Veloso VG, *et al*, (2011). Breast cancer in a cohort
393 of human immunodeficiency virus (HIV)-infected women from Rio de Janeiro,
394 Brazil: a case series report and an incidence rate estimate. *Braz J Infect
395 Dis*;15:387-93

396 41. Shiels MS, Pfeiffer RM, Gail MH, Hall HI, Li J, Chaturvedi AK, *et al.* (2011).
397 Cancer burden in the HIV-infected population in the United States. *J Natl*
398 *Cancer Inst*, 103:753–62. 10.1093/jnci/djr076
399 42. Jessica L Castillo Email author, Paula M Luz, Bryan E Shepherd, Megan
400 Turner, Sayonara R Ribeiro, Sally S Bebawy, Juliana S Netto, Catherine C
401 McGowan, Valdiléa G Veloso, Eric A Engels, Timothy R Sterling and Beatriz
402 Grinsztejn.(2015). HIV and cancer: a comparative retrospective study of Bra-
403 zilian and U.S. *clinical cohorts Infectious Agents and Cancer BioMed Cen-*
404 *tral*.doi: 10.1186/1750-9378-10-4.
405 43. Muguti GI (1993) Experience with breast cancer in Zimbabwe. *J R Coll Surg*
406 *Edinb* 38(2): 75–78 | PubMed | ChemPort |

410
411
412
413
414
415
416
417
418



419
420
421

Figure 1: Patient 1 advanced breast cancer (Stage 4)

422
423




Figure 2: Patient 2 advanced ulcerated breast cancer (stage 4)

424
425
426
427
428




Figure 3: Patient 3 advanced ulcerated breast cancer (stage 4)

429
430
431




432
433
434
435
436

Figure 4: Stage of the disease and frequency




437
438
439


Figure 5: Prevalence of self-delay




440
441 **Figure 6: Clinical Stage vs Domicile**
442
443
444
445



446
447 **Figure 7: Breast cancer-age distribution**
448
449
450



451
452 **Figure 8: Highest level of education**
453



454
455 **Figure 9: Frequency distribution of reasons for delay**
456
457

458 **Table 1: Knowledge of self- breast examination and Residence**

Residence	Knowledge of Breast Self-Examination		Total
	Yes (%)	No (%)	
Rural	12 (16.4)	37 (50.7)	49 (67.1)
Urban	7 (9.6)	17 (23.3)	24 (32.9)
Total	19 (26.0)	54 (74.0)	73 (100)

Note: $p < 0.05$,Statistically significant association

459

460
461
462

Table 2: Knowledge of ~~breast self-self~~ breast examination

Knowledge of Self Breast Examination

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	yes	19	26.0	26.0	26.0
	no	54	74.0	74.0	100.0
	Total	73	100.0	100.0	

463
464
465
466
467
468
469
470
471
472
473
474
475
476

Table 3: Knowledge of self- breast examination and Domicile

		Knowledge of Self Breast Examination		Total
		yes	no	
Residence	Rural	12	37	49
	urban	7	17	24
Total		19	54	73

($p < 0.05$, Statistically significant)

477
478
479
480
481
482
483
484
485
486

Table 4: Relationship between knowledge of self-breast examination and age group

		Knowledge of Self Breast Examination		Total
		yes	no	
Age Group	11-20	1	1	2
	21-30	3	9	12
	31-40	5	8	13
	41-50	3	12	15
	51-60	4	16	20
	61-70	2	7	9
Total		18	53	71

487

488
489
490
491
492
493

($p > 0.05$, Not statistically significant)

Table 5: Symptoms

Symptom	Frequency	Percent
Mass	57	78.1
Nipple Discharge	12	16.4
Nipple Retraction	8	11
Pain	29	39.7
Ulcer	13	17.8

494
495
496
497
498
499
500

Table 6: Relationship between delay and employment status

Employed	Delay		Total
	no (< 3 months)	Yes (\geq 3 months)	
no	41	10	51
yes	15	3	18
Total	56	13	69

501
502
503
504
505
506
507
508
509

($p > 0.05$, Statistically insignificant)

Table 7: Relationship between [Knowledge-knowledge](#) of [breast self-self-breast](#) examination and level of education

Knowledge of Self Breast Cancer	Level of Education				Total
	Tertiary course / Diploma	Secondary	Primary	Never attended	
no	0 (0%)	14 (20.9%)	10 (14.9%)	2 (3.0%)	26 (38.8%)
yes	6 (9.0%)	22 (32.8%)	12 (17.9%)	1 (1.5%)	41 (61.2%)
Total	6 (9.0%)	36 (53.7%)	22 (32.8%)	3 (4.5%)	67 (100.0%)

Note: $p < 0.05$, Statistically significant association

510
511
512
513
514
515

516
517
518
519

Table 8: Contributions of specific reasons to delayed breast cancer presentation

Reasons	Delayed presentation score	
	No	Yes
HIV Status	20.240	24.526
Age	6.169	7.406
Early Menarche	-1.521	-2.525
Family History	.055	.148
Late Menopause	7.697	4.812
Level of Education	5.269	8.898
(Constant)	-91.994	-115.295

Note: Classification Function Coefficients determined by Fisher's linear discriminant functions

520
521
522
523
524

Table 9: Standardized discriminant Coefficients by reason

Reason	Function
	1
HIV Status	.890
Age	1.061
Early Menarche	-.524
Family History	.221
Late Menopause	-.424
Level of Education	.679

525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547

548
549
550
551

Table 10: Group means and standard deviations

Delay		Mean	Std.	N		
				Unweighted	Weighted	
Delayed (≥ 3 months)	HIV Status	1.33	0.58	3	3	
	Age	18.67	2.52	3	3	
	Early Menarche	13.00	1.00	3	3	
	Family History	1.67	0.58	3	3	
	Late Menopause	2.00	0.00	3	3	
	Level of Education	1.67	0.58	3	3	
	Knowledge of Self Breast Examination (BE)	1.33	0.58	3	3	
	Health Worker of first Contact	2.67	1.16	3	3	
	Duration of Symptoms in Months	2.67	2.08	3	3	
	Marital Status	2.00	1.00	3	3	
	Employed	1.00	0.00	3	3	
	Not delayed (< 3 months)	HIV Status	2.00	0.63	6	6
		Age	21.83	2.56	6	6
Early Menarche		14.17	1.72	6	6	
Family History		5.17	8.25	6	6	
Late Menopause		1.67	0.52	6	6	
Level of Education		2.50	0.55	6	6	
Knowledge of Self (BE)		1.17	0.41	6	6	
Health Worker of first Contact		2.33	0.82	6	6	
Duration of Symptoms in Months		2.17	1.60	6	6	
Marital Status		2.50	0.55	6	6	
Employed		1.67	0.52	6	6	

552