

METASTABLE NON-NUCLEONIC STATES OF NUCLEAR MATTER:  
PHENOMENOLOGY

It was earlier shown that the metastable states of the nuclear matter may exist when the nuclear forces are not strong enough to bind a part of the quarks into stable nucleons, which leads to local shake-ups in the nucleonic structure of the nucleus. For these anomalous excited states of the nuclear matter, called inner-shake-up or *isu*-states, the relaxation of the nuclei is initiated by the weak nuclear interaction. The existence of nuclei with a shaken-up nucleonic structure makes it possible to physically interpret a rather large group of experimental data on the initiation of low energy nuclear reactions (LENRs) and the acceleration of radioactive  $\alpha$ - and  $\beta$ -decays in a low-temperature plasma. The possible mechanisms of LENRs implemented in a Rossi E-CAT reactor are discussed. It is also suggested that the metastable *isu*-states of a different type occur as a result of high-energy collisions of particles, when heavy hadrons (baryons, mesons) are formed in the collisions of protons with characteristic energies higher than 1 TeV. This kind of concept makes it possible to physically interpret the recently recorded anomaly in the angular  $e^+e^-$  correlations of positron-electron pairs emitted in the radioactive decays of excited  ${}^8\text{Be}$  nuclei formed by the interaction between protons with kinetic energy  $\sim 1$  MeV and  ${}^7\text{Li}$  nuclei. It is this anomaly that can become the basis for introducing a fifth fundamental interaction into physics, in addition to the strong/weak nuclear, electromagnetic, and gravitational interactions.

*Keywords: metastable non-nucleonic states of nuclear matter; low energy nuclear reactions; heavy hadrons; heavy quarks; inner shake-up state of nuclear matter.*

## 1. INTRODUCTION

Recent studies [1, 2] suggested that there might be a fifth fundamental interaction, in addition to the strong/weak nuclear, electromagnetic, and gravitational ones. They were initiated by the study [3], in which the authors studied the radioactive decays of excited  ${}^8\text{Be}$  nuclei with energies of 17.64 and 18.15 MeV, formed in the interaction of protons with kinetic energy  $E_p = 1.10$  MeV and  ${}^7\text{Li}$  nuclei, using  $\text{LiF}_2$  and  $\text{LiO}_2$  targets. The above excited states were recorded as resonances at  $E_p = 0.441$  MeV and  $E_p = 1.03$  MeV in the process  ${}^7\text{Li}(p, \gamma){}^8\text{Be}$  under study. The authors [3] studied the formation of a positron-electron pair  $e^+ - e^-$  resulting from the internal conversion that accompanies the birth of two  $\alpha$ -particles in the radioactive decay of the  ${}^8\text{Be}$  nuclei. They expected a sharp drop in the probability of the correlated formation of an  $e^+ - e^-$  pair as the opening angle  $\Theta$  between positrons and electrons in the laboratory frame of reference increases. However, they recorded an “anomalous” increase in the angular function within an angular range of  $\Theta \sim 130\text{-}140^\circ$ , considering this anomaly as a result of the formation of the  $e^+e^-$  pair in the decay of a hypothetical neutral isoscalar boson formed in the above process, with a rest mass equal to  $16.7 \text{ MeV}/c^2$ , where  $c$  is the speed of light in vacuum. In this decay, the opening angle would be  $180^\circ$  in the system of the center of mass of the  $e^+e^-$  pair. It was suggested that the introduced isoscalar boson, with its expected lifetime of  $\sim 10^{-14}$  s, might be a good candidate for the relatively light gauge boson performing the role of the mediator in the secluded WIMP dark matter scenario. However, the later analysis [1, 2] showed that the  ${}^8\text{Be}$  anomaly,

50 which is consistent with all existing experimental constraints, can be adequately interpreted  
 51 only when one puts forward a hypothesis of existence of another type of boson, a protophobic  
 52 gauge vector boson  $X$ , which is produced in the decay of the excited state of  ${}^8\text{Be}^*$  down to the  
 53 ground state,  ${}^8\text{Be}^* \rightarrow {}^8\text{Be}^*X$ , and then decays as  $X \rightarrow e^+e^-$ . It is the boson that could be  
 54 related to the elusive dark matter in the Universe. According to [1, 2], this boson, with a mass  
 55 of about  $17 \text{ MeV}/c^2$ , is the mediator of the weak force, interacts with neutrons but is  
 56 “protophobic” and ignores protons. The latter allows one to explain why the  $X$  boson might  
 57 have avoided earlier detection. It was shown that the Standard Model can be easily extended  
 58 to accommodate a light gauge boson with protophobic quark couplings [2]. As a result, the  
 59 postulated boson was associated with a new, fifth, fundamental interaction, which should be  
 60 introduced into the physical science [1, 2].

61

62 It will be shown below that the recorded anomalies in the angular  $e^+e^-$  correlations in  
 63 the radioactive decay of excited  ${}^8\text{Be}$  nuclei can be qualitatively interpreted on the basis of a  
 64 new concept rather than the fundamental hypotheses made in [1, 2]. We assume that  
 65 metastable states can occur in the nuclear matter when the nuclear forces are not strong  
 66 enough to bind a part of the quarks into nucleons, which gives rise to local shake-ups in the  
 67 nucleonic structure of the nucleus. For these anomalous excited states, the relaxation  
 68 dynamics of the nuclei crucially depends on the weak nuclear interaction. Earlier, this  
 69 assumption made it possible to physically interpret a rather large set of experimental data on  
 70 the initiation of LENRs and acceleration of radioactive  $\alpha$ - и  $\beta$ -decays in a low-temperature  
 71 plasma. It will be shown that such states of the nuclear matter with a shaken-up nucleonic  
 72 structure can occur in the high-energy collisions of nuclei too, such as protons in the colliding  
 73 beams with characteristic energies higher than 1 TeV. The concept to be introduced will  
 74 allow one to understand why the decay of highly excited hadrons (baryons, mesons) formed  
 75 in these collisions is effectuated by the weak nuclear interaction.

76

## 77 2. ELECTRON FACTOR IN INITIATING NUCLEAR PROCESSES

78

79 Phenomenological approach [4-7] implies that the dynamic interrelation between the electron  
 80 and nuclear subsystems of an atom, which is mediated by the electromagnetic component of  
 81 the physical vacuum (EM vacuum), is the key factor in initiating LENRs [8-12] and the  
 82 radioactive decay of nuclei [5, 8, 13]. This interrelation manifests itself in experimentally  
 83 recorded facts that the occurrence of radioactive decay of nuclei is accounted for by the  
 84 positive difference between the total mass of the initial atom subsystems, electron and nuclear  
 85 (whole atom rather than the nucleus alone), and the total mass of its decay products [14, 15].  
 86 When the mechanisms of LENRs and the decay of atomic nucleus  ${}^A_ZN$  ( $Z$  and  $A$  are the  
 87 atomic and mass numbers of the nucleus  $N$ , respectively) are considered, the nuclear matter  
 88 is usually represented in the form of interacting nucleons. In the  $K$ -capture, however, when  
 89 the electron of the inner shells of an atom interacts with the surface of the nucleus, giving rise  
 90 to a new daughter nucleus, the nucleonic structure of the nuclear matter is unchanged. At the  
 91 initial irreversible stage of this process, the electron interacting with the nucleus surface emits  
 92 a neutrino  $\nu$ . The resulting virtual vector  $W^-$ -boson, integrated into the nuclear matter,  
 93 interacts with the  $u$ -quark of one of the protons and is converted to a  $d$ -quark. As a result, this  
 94 proton is converted to a neutron, and a nucleus  ${}^A_{Z-1}M$  is formed. However, the situation can  
 95 drastically change when the  $K$ -capture is energetically forbidden, which are the cases under  
 96 consideration below, and the electron can acquire a rather high (on chemical scales) kinetic  
 97 energy  $E_e \sim 3\text{-}5 \text{ eV}$ , which can occur in a low-temperature plasma. In this case, when the  
 98 electron shells are not yet ionized by these electrons, the scattering of electrons with the

99 above kinetic energy (de Broglie wavelength  $\lambda \approx 0.5$  nm) by atoms and ions initiates the  
 100 oscillation of the electron subsystems of the atoms and ions, increasing the probability of  
 101 interaction between the electrons of the inner subshells of the atoms and ions and their  
 102 respective nuclei.


103 The first, irreversible stage of this interaction is characterized by the emission of a  
 104 neutrino  $\nu$  and the integration of a vector  $W^-$ -boson into the nuclear matter of the initial  
 105 nucleus  ${}^A_Z N$  :



107  
 108 As a result, the nucleonic structure of the formed nucleus  ${}^A_{Z-1} M_{isu}$ , with a charge less by one  
 109 than that of the initial nucleus, is locally shaken up. Indeed, the interaction between the  
 110 vector  $W^-$ -boson and the  $u$ -quark of one of the protons of the nucleus  ${}^A_Z N$  can only produce a  
 111 virtual  $d$ -quark followed by a chain of virtual conversions of quarks involving vector  $W$ -  
 112 bosons. At the same time, the deficit of the total mass of this nucleus prevents from the  
 113 formation of a neutron. The resulting state of local anomaly in the nuclear matter with a  
 114 shaken-up nucleonic structure is characterized as a metastable inner-shake-up state, or  $isu$ -  
 115 state. The latter is indicated by the subscript on the right of the nucleus symbol in the right-  
 116 hand side of (1). The subscript in the electron symbol in the left-hand side of (1) indicates  
 117 that this stage of the process is activated. The initiated chain of virtual conversions of quarks  
 118 in which the vector  $W$ -bosons are involved must be interrupted by the irreversible decay of  
 119 the virtual  $W^-$ -boson producing the initial nucleus, electron, and antineutrino  $\tilde{\nu}$  :



123  
 124 Consequently, the overall process can be represented as an inelastic scattering of an electron  
 125 by the initial nucleus:



127  
 128 The nuclei in which the nuclear matter is in a metastable  $isu$ -state will be called “ $\beta$ -nuclei”.  
 129 The threshold energy for this process producing a  $\nu\tilde{\nu}$  pair, which is accounted for by the  
 130 neutrino-antineutrino rest masses, is about 0.3 eV [16].

131  
 132 It is common knowledge that the nucleus is a system of nucleons bound into a whole  
 133 by exchange interactions in which the quarks are exchanged using pions. Therefore, the  
 134 formation of three quarks not bound into a nucleon in a nucleus, which in this case can be  
 135 regarded as “markers” of new degrees of freedom, in fact, means that the intensity of nuclear  
 136 forces is not enough to provide the traditional proton-neutron arrangement of the nuclear  
 137 matter in the system under consideration. The subsequent relaxation of the locally formed  
 138  $isu$ -state, which can be transferred by the mediating pions to other nucleons of the nucleus, is  
 139 initiated only by the weak nuclear interactions, which are effectuated by the mediating quarks  
 140 in the formation and absorption of gauge vector neutral  $Z^0$  and charged  $W^\pm$ -bosons. In the  
 141 case under consideration, this relaxation terminates with the decay of the virtual vector  $W^-$ -  
 142 boson followed by the formation of the initial nucleus in the emission of an electron and  
 143 antineutrino. The lifetime of the formed  $\beta$ -nuclei found in the metastable  $isu$ -state can be  
 144 rather long, from tens of minutes to several years, and the nuclei in this state can be directly  
 145 involved in various nuclear processes [4, 5].

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190


191

192

It should be noted that the relaxation rearrangement of the nuclear matter when the products of these nuclear conversions are formed is effectuated primarily by forming a purely nucleonic structure of the nucleus, obeying the principle of least action. While the relaxation processes of de-excitation in nuclei with a proton-neutron (nucleonic) structure can go via the excited states of the nucleus and include  $\gamma$ -quanta emission steps, this type of relaxation in  $\beta$ -nuclei is virtually impossible. Therefore, if the atomic nuclei in which the nuclear matter is in a partial “non-nucleonic” state are involved in the processes, the mechanism of relaxation of the formed products is always accompanied by energy loss due to the emission of neutrino-antineutrino pairs, or due to the URCA process [17], rather than the emission of  $\gamma$ -quanta by excited nuclei, as in the relaxation of nuclear products characterized by the proton-neutron arrangement of the nuclear matter. It is for this reason that the corresponding nuclear processes are safe for the environment.

Of special interest are the cases in which the formation of *isu*-states in the nuclear matter is initiated in initially radioactive nuclei because the relaxation process with a vector  $W^-$ -boson decay can initiate a general radioactive decay of the *isu*-state nucleus that results in the formation of daughter products of the decay of the initial radioactive nucleus. According to [5, 18], the general stability of the nuclear matter in a metastable *isu*-state can be lost by changing the boundary conditions for the components of the electric field intensity vector of the EM vacuum at the surface of the nucleus in whose volume the nucleonic matter shake-up occurred. The index characterizing the  ${}_{Z-1}^A M_{isu}$  nucleus instability that occurs in the process (1) is the absolute value of the structural energy deficit  $\Delta Q$  ( $\Delta Q < 0$ ) of this metastable *isu*-state nucleus, which is defined as  $\Delta Q = (m_{\frac{A}{2}N} - m_{\frac{A}{Z-1}M})c^2$ . In this case, the mass of the  ${}_{Z-1}^A M_{isu}$  nucleus is taken as  $m_{\frac{A}{Z-1}M_{isu}} = m_{\frac{A}{2}N} + m_e$ , where  $m_{\frac{A}{2}N}$  is the mass of the  ${}_{Z}^A N$  nucleus and  $m_e$  is the rest mass of the electron.

For example, in the laser ablation of metal samples in an aqueous solution of uranyl, when a low-temperature plasma is formed in the vapor near the metal surface, the interaction between the plasma electrons and the  ${}^{238}\text{U}$  nuclei initiates the formation of “ $\beta$ -protactinium” nuclei followed by a  $\beta$ -decay of the  ${}_{91}^{238}\text{Pa}_{isu}$  nuclei that produces thorium-234 and helium-4 nuclei as the products of decay of the initial uranium-238 nucleus:



In this case, the effective rate constant for the initiated decays of  ${}^{238}\text{U}$  nucleus increases by 9 orders of magnitude, giving rise to a kind of “ $e^-$ -catalysis” [4]. The deficit  $\Delta Q$  of structural energy for the formed  $\beta$ -protactinium nucleus is  $\Delta Q \approx -3.46 \text{ MeV}$ . An unexpected result was recorded in experiments with a beryllium sample. The beryllium nanoparticles formed in the solution after one-hour laser action showed an anomalously high rate of formation of thorium-234 nuclei for more than 500 days after the laser ablation was completed. The half-life for the nuclei initiated by the laser ablation that produce thorium-234 was 2.5 years. Naturally, this phenomenon could be associated with the accumulation of  $\beta$ -protactinium nuclei in beryllium nanoparticles in the laser ablation, which lasted as short as an hour.

Additional examples are the  $\beta^-$ -decay of  ${}_{27}^{60}\text{Co}$ ,  ${}_{55}^{137}\text{Cs}$  and  ${}_{56}^{140}\text{Ba}$  nuclei initiated by the  $e^-$ -catalysis mechanism, for which the half-life  $T_{1/2}$  is 1925 days, 30.1 years, and 12.8 days, respectively [4, 7]:

193

194  ${}_{27}^{60}\text{Co} + e_{he}^- \rightarrow {}_{26}^{60}\text{Fe}_{isu} + \nu \rightarrow {}_{28}^{60}\text{Ni} + 2e^- + \nu + 2\bar{\nu} + Q(2.82 \text{ MeV}),$  (5)

195  ${}_{55}^{137}\text{Cs} + e_{he}^- \rightarrow {}_{54}^{137}\text{Xe}_{isu} + \nu \rightarrow {}_{56}^{137}\text{Ba} + 2e^- + \nu + 2\bar{\nu} + Q(1.18 \text{ MeV}).$  (6)


196  ${}_{56}^{140}\text{Ba} + e_{he}^- \rightarrow {}_{55}^{140}\text{Cs}_{isu} + \nu \rightarrow {}_{57}^{140}\text{La} + e^- + \nu + 2\bar{\nu} + Q(1.05 \text{ MeV}).$  (7)

197

198 In these examples, the deficits of structural energy  $\Delta Q$ , which prevent the *isu*-state  ${}_{26}^{60}\text{Fe}_{isu}$ ,  
 199  ${}_{54}^{137}\text{Xe}_{isu}$  and  ${}_{55}^{140}\text{Cs}_{isu}$  nuclei from coming to the stable ground states of the nuclear matter  
 200 referring to the  ${}_{26}^{60}\text{Fe}$ ,  ${}_{54}^{137}\text{Xe}$ , and  ${}_{55}^{140}\text{Cs}$  nuclei, is -0.237, -4.17, and -6.22 MeV, respectively .  
 201 It can be expected that the initiation effect of electrons on the  $\beta^-$ -decay of nuclei in a low-  
 202 temperature plasma will be best manifested when the absolute value of structural energy  
 203 deficit  $\Delta Q$  for the *isu*-state nuclei to be formed is the highest. This implies that in the above  
 204 cases the acceleration of radioactive decay would be clearly seen for the  ${}_{55}^{137}\text{Cs}$  and  ${}_{56}^{140}\text{Ba}$   
 205 nuclei and minimal for the  ${}_{27}^{60}\text{Co}$  nuclei. The available experimental data [8] on the initiated  
 206 decays of  ${}_{55}^{137}\text{Cs}$ ,  ${}_{56}^{140}\text{Ba}$ , and  ${}_{27}^{60}\text{Co}$  validate this conclusion: the half-lives of  $\beta^-$ -active cesium-  
 207 137 (30.1 years) and barium-140 nuclei (12.8 days) drop to about 380 and 2.7 days,  
 208 respectively, whereas the half-life of cobalt-60, equal to 1925 days, remains practically  
 209 unchanged. The Feynman diagrams for the  $\beta^-$ -decays, positron  $\beta^+$ -decays, and  $\alpha$ -decays of  
 210 nuclei initiated by the  $e^-$ -catalysis mechanism are plotted in Fig. 1.

211

212




213

214


215

216



217

218



219  
220

*d*

221 Fig. 1. Feynman diagrams for initiated (a)  $\beta^-$ -decays, (b)  $\beta^+$ -decays, and (c, d)  $\alpha$ -decay  
222

223 The unexpected character of the result that the decay of a radioactive nucleus can be affected  
224 by external actions consists in the fact that this effect is associated with electrons, which  
225 cannot interact with the nucleons of the nucleus as nuclear matter fragments, but can initiate,  
226 with the help of vector  $W^-$ -bosons, local shake-ups in the nucleonic structure of the nucleus.  
227 At the same time, the experiments show that the external excitation of a radioactive nucleus  
228 as a whole system (for example, by the action of  $\gamma$ -radiation) cannot affect the rate of  
229 radioactive decay and, hence, the above initiation of the nucleus instability. In these cases, the  
230 nuclear matter manifests itself as a whole system of interacting nucleons with their inherent  
231 individual characteristics.

232

233 Section 6 will show how the external actions of very high energy can give rise to *isu*-  
234 states in the nuclear matter, which account for the decay of nuclei.

235

236

### 237 3. POSSIBLE MECHANISMS OF NUCLEAR-CHEMICAL REACTIONS

238 The simplest  $\beta$ -nuclei are  $\beta$ -neutrons and  $\beta$ -dineutrons, which can be formed by the  
239 interaction of high-energy electrons with protons  $p^+$  and deuterons  $d^+$ , respectively; for  
240 example, in the laser ablation of metals in an ordinary or heavy water, as well as in a protium-  
241 or deuteron-containing glow-discharge plasma:

242 
$$p^+ + e^-_{he} \rightarrow {}^1n_{isu} + \nu, \quad (8)$$

243 
$$d^+ + e^-_{he} \rightarrow {}^2n_{isu} + \nu. \quad (9)$$

244 If the half-lives  $T_{1/2}$  of these  $\beta$ -nuclei are sufficiently long, the neutral nuclei  ${}^1n_{isu}$  and  ${}^2n_{isu}$ ,  
245 which are characterized by the baryon numbers equal to one and two, the rest masses equal to  
246 the masses of the hydrogen atom and deuterium, respectively, and by zero lepton charges, can  
247 be efficiently involved in various nuclear processes [4-7, 13].

248

249 Analysis of experimental data on the synthesis of tritium nuclei  $t^+$  in the laser ablation  
250 of metals in a heavy water shows that the half-life  $T_{1/2}$  of the  $\beta$ -dineutron decay,

251 
$${}^2n_{isu} \rightarrow d^+ + e^- + \tilde{\nu}, \quad (10)$$

252 which produces a deuteron, electron, and antineutrino, is rather long, at least, tens of minutes  
 253 [13]. It was assumed that the synthesis occurs by the interaction between a tritium nucleus  $t^+$   
 254 and nucleus  ${}^2n_{isu}$ :

$$255 \quad d^+ + {}^2n_{isu} \rightarrow t^+ + n + Q(3.25MeV), \quad (11)$$

256 where  $n$  stands for a neutron. This is accompanied by another process:

$$257 \quad d^+ + {}^2n_{isu} \rightarrow {}^3He + n + e^- + \tilde{\nu} + Q(3.27MeV), \quad (12)$$

258 which is a result of the weak nuclear interaction.

259 The authors [13] also postulated that the interaction between electrons and a tritium nuclei  $t^+$   
 260 may produce a hypothetical  $\beta$ -trineutron  ${}^3n_{isu}$ :

$$261 \quad t^+ + e^-_{he} \rightarrow {}^3n_{isu} + \nu. \quad (13)$$

262

263 The rest mass of the neutral nucleus  ${}^3n_{isu}$  was assumed to be equal to the rest mass of  
 264 the tritium atom. It is the formation of  ${}^3n_{isu}$  that the initiated decay of tritium nuclei in the  
 265 laser ablation of metals in aqueous media and the synthesis of tritium nuclei can pass through  
 266 [13]:

$$267 \quad t^+ + e^-_{he} \rightarrow {}^3n_{isu} + \nu \rightarrow {}^3He + 2e^- + \nu + 2\tilde{\nu} + Q(0.019MeV). \quad (14)$$

268 It should be noted that the half-life  $T_{1/2}$  of  ${}^3n_{isu}$  in the  $e^-$ -catalysis is of the same order of  
 269 magnitude as that of  ${}^2n_{isu}$ , which is many orders of magnitude shorter than the half-life of the  
 270 tritium nucleus ( $T_{1/2} = 12.3$  years) [13].

271 It is shown in [5] that the introduced concept of a  $\beta$ -nuclei with a rather long lifetime  
 272 formed in the glow discharge in a deuterium-containing gas makes it possible to physically  
 273 interpret a group of data [9, 10] on the initiated radioactive decay of  $W$  nuclei in the surface  
 274 layers of a tungsten cathode (foil). Note that although 5 isotopes of tungsten  
 275 ( ${}^{180}_{74}W$ ,  ${}^{182}_{74}W$ ,  ${}^{183}_{74}W$ ,  ${}^{184}_{74}W$ ,  ${}^{186}_{74}W$ ) are potentially  $\alpha$ -radioactive nuclei,

$$276 \quad {}^A_{74}W \rightarrow {}^{A-4}_{72}Hf + {}^4_2He + Q_A, \quad (15)$$

277 they are usually considered as stable isotopes because of an anomalously large period of their  
 278 half-life,  $T_{1/2} \approx 10^{17} - 10^{19}$  years, which is many orders of magnitude greater than the lifetime  
 279 of the Universe. The values of heat release  $Q_A$  in the radioactive  $\alpha$ -decay of tungsten nuclei  
 280 with mass numbers  $A$  equal to 180, 182, 183, 184, and 186, is 2.52, 1.77, 1.68, 1.66, and 1.12  
 281 MeV, respectively. Based on the energy consideration alone, one can admit that there are  $\alpha$ -  
 282 decays producing several  $\alpha$  particles for the above stable isotopes of tungsten, including the  
 283 decay producing nine  $\alpha$  particles for the tungsten-180 isotope.





284

285 The concept dealing with the formation of metastable  $isu$ -state nuclei that we are  
 286 developing distinguishes three mechanisms for initiated nuclear conversions, including  
 287 radioactive decays of nuclei:

288 1. *Mechanism of nuclear fusion.* The neutral particles  ${}^A n_{isu}$  ( $A = 1, 2, 3$ ) with long enough  
 289 lifetimes formed in a low-temperature plasma can diffuse along grain boundaries deep into  
 290 the cathode and interact with the metal (tungsten) nuclei in the cathode surface layers. In this  
 291 case, the interaction and fusion of  ${}^2n_{isu}$  nuclei with  ${}^A_{74}W$  isotopes can give rise to excited  
 292  ${}^{A+2}_{74}W^*$  nuclei at the first process step. In addition to the overall excitation energy, indicated  
 293 by the asterisk, equal to about 10 MeV relative to the stable ground state of these nuclei, their  
 294 nuclear matter due to their fusion with  ${}^2n_{isu}$  can be partially in an unbalanced  $isu$ -state with a

295 lost stability in the nucleus bulk. All of this causes the resulting conversions accompanied by  
 296 the emission of  $\alpha$  particles and daughter isotopes. Note that in contrast to the nuclear  
 297 reactions that occur in the collision of reactants in the gaseous phase, the energy factor alone  
 298 due to the possible effect of the environment is enough to effectuate the above nuclear  
 299 conversions in the region of grain boundaries of the solid metal phase, with the unmatched  
 300 spins and parities of the colliding and resulting nuclei.

301  
 302 Experimental works studying the conversions in the glow discharge in a deuterium-  
 303 containing gas recorded the formation of new elements in the surface layer of the tungsten  
 304 cathode after it was treated by the plasma for 4 to 7 hours, which include not only stable  
 305 isotopes of erbium, ytterbium, lutetium and Hafnium, but also radioactive isotopes of  
 306 ytterbium and hafnium [9, 10]. While the formation of the stable isotopes could be assumed  
 307 to be related to the diffusion of impurity elements from the cathode bulk to its surface treated  
 308 by the plasma, the formation of the radioactive isotopes definitely points to the radioactive  
 309 decay of tungsten isotopes. As all possible reactions for the initiated decay of various  
 310 tungsten isotopes are already reported [5], only a few examples are given below for  
 311 illustration:





316  
 317 In (16) - (19) it is taken into account that in addition to the major masses 169 to 180, the mass  
 318 spectra of the products recorded the birth and growth of the peak of mass 9. It should be  
 319 noted that the absence of the basic mass 4, corresponding to helium nuclei, in the mass  
 320 spectra recorded in [9, 10] can be attributed to the extremely low solubility of helium in  
 321 tungsten [19] and the high diffusivity of helium in the zone between the boundaries of foil  
 322 grains. It is obvious that these transport processes can be accomplished only when the  
 323 lifetime of  ${}^2n_{isu}$  is long enough for the diffusive transport of these neutral nuclei along the foil  
 324 grain boundaries to surface layers. This agrees with the conclusion that this time must be no  
 325 less than tens of minutes for the synthesis of tritium in the laser ablation of metals in a heavy  
 326 water [13].

327  
 328

## 329 2. Mechanism of $e^-$ -catalysis.

330 The above consideration implies that there may be another way of initiating the  $\alpha$ -decay of  
 331 tungsten isotopes in a glow discharge in experimental studies, when electrons with kinetic  
 332 energy  $E_e \sim 3\text{-}5$  eV interact directly with stable isotopes of tungsten in the  $e^-$ -catalysis.  
 333 Possible examples of these processes are given below:



336

337 It should be noted that the concept of  $e^-$ -catalysis can be helpful in understanding the  
 338 formation of much less than all new isotope products recorded in experiments. Therefore, the  
 339 processes with  ${}^2n_{isu}$  nuclei are considered as basic for the initiated decays of  $W$  stable  
 340 isotopes.

341



342 The above data allow one to state that the nuclear decay of initially non-radioactive tungsten  
 343 isotopes accompanied by the formation of lighter elements (erbium, lutetium, ytterbium,  
 344 hafnium), which is initiated in a low-temperature plasma (glow discharge), can be considered  
 345 as a new type of artificial radioactivity, which is different from the artificially induced  
 346 radioactivity initiated by nuclear reactions (e.g., by bombardment with alpha particles or  
 347 neutrons, giving rise to radioisotopes). It should be remembered that the stable isotopes of  
 348 many nuclei, from neodymium to bismuth, including a tantalum-181 isotope, for which  
 349 initiated decays similar to those described above are also recorded [9, 10], are potentially  $\alpha$ -  
 350 radioactive in the same sense as tungsten isotopes.



351

352

353 *3. Harpoon mechanism.*

354

355 The reactions between multi-electron atoms are of highest complexity for understanding the  
 356 mechanism of low-energy nuclear processes. These processes are usually considered in the  
 357 study of transformation processes in native systems [20-22]. It was recently shown [23],  
 358 however, that reactions of this type can occur in the initiation of self-propagating high-  
 359 temperature synthesis (SHS) [24]. The composition of the condensed products of thermite  
 360 powder mixture (Al + Fe<sub>2</sub>O<sub>3</sub>) combustion in air was studied [23]. The purity of the initial  
 361 materials was 99.7 to 99.9 mass %. It was shown that in the combustion of iron-oxide  
 362 aluminum thermites with a flame temperature higher than 2800 K, 0.55 mass % of stable  
 363 calcium is formed. The initial thermite powder systems (Al + Fe<sub>2</sub>O<sub>3</sub>) did not contain any  
 364 calcium. According to [23], the calcium could be formed in the following nuclear reactions:



367



368 The formation of calcium in the experiments [23] implies that the temperature of  
 369 electrons in the flame of combustion of iron-oxide aluminum thermites in air can be much  
 370 higher than the flame temperature estimated using the energy of atoms and ions. It is the  
 371 case that is typical for the low-temperature plasma in a glow discharge. In this case, the  
 372 interaction of high-energy electrons with nuclei  ${}_{13}^{27}\text{Al}$  and  ${}_7^{14}\text{N}$  could produce nuclei  ${}_{13}^{27}\text{Mg}_{isu}$   
 373 and  ${}_{6}^{14}\text{C}_{isu}$ , respectively. Among these nuclei, the nucleus  ${}_{13}^{27}\text{Mg}_{isu}$  shows the highest activity  
 374 in the nuclear interactions because the deficit of its energy relative to the nucleus  ${}_{13}^{27}\text{Mg}$  is  
 375  $\Delta Q = -2.61\text{ MeV}$ , whereas the energy deficit for the nucleus  ${}_{6}^{14}\text{C}_{isu}$  is much less,  $\Delta Q = -0.16$   
 376  $\text{MeV}$ .

377

378 Following [7], assume that when the nucleus of an atom, or ion is in a pre-decay metastable  
 379 *isu*-state (supposedly,  ${}_{13}^{27}\text{Mg}_{isu}$ ), the lability of its electron subsystem is higher and it is likely  
 380 that this subsystem can partially overlap the electron subsystems of the neighboring atoms  
 381 (specifically, the nitrogen atom). It is obvious that the high values of energy release in the  
 382 overall processes (22) and (23) should act as the factor initiating the spin-spin interaction of  
 383 the electron subsystems of both atoms and the formation of common “molecular” orbitals  
 384 with the correcting action of spin electron-nuclear interactions for each atom. The emerging  
 385 bonds bring both atoms closer to each other, and the formation of common orbitals is more  
 386 intense as the nuclei are brought closer to each other. This brings about a kind of “harpoon  
 387 mechanism” in which the atom with an *isu*-state nucleus captures the adjacent atom. The  
 388 complete integration of the electron subsystems of both atoms initiates the fusion of the

389 nuclear matter of the *isu*-state nucleus ( ${}_{13}^{27}\text{Mg}_{isu}$ ) and the adjacent nucleus ( ${}_{7}^{14}\text{N}$ ). In this case,  
 390 the overall processes can be written as

391



394 Earlier, the harpoon mechanism was considered for the nuclear transmutations in native  
 395 systems [7].

396 Because weak nuclear interactions are involved in the formation of the nuclear matter  
 397 in the final nucleus as a set of interacting nucleons, a significant part of the energy can also  
 398 be released by emitting neutrinos and antineutrinos when the final nucleus can be formed in  
 399 the ground state, obeying the spin and parity conservation laws. At the same time, when the  
 400 final nuclei are formed in the excited state, the non-ionizing radiation of neutrinos and  
 401 antineutrinos will be accompanied by the emission of X-rays or gamma quanta. These X-rays  
 402 were already reported experimentally [23].

403 The above phenomenological analysis shows that in order to understand the  
 404 mechanism of nuclear transformations observed in the burning of thermite mixtures, of great  
 405 importance is the development of new theoretical approaches to simulating the dynamics of  
 406 nuclear processes on the basis of quantum-chemical analysis rather than estimating the  
 407 quantum mechanical probabilities of some processes. This simulation will involve (1)  
 408 calculations of the electron structure of an atom when *isu*-state nuclei with a shaken-up  
 409 nucleonic structure are formed; (2) model calculations of the spatial instability of the atom  
 410 electron subsystem, which is caused by the loss of the nucleus stability; (3) calculations of the  
 411 overlapping of these “mobile” orbitals with the electron orbitals of the adjacent atoms and  
 412 formation of molecular orbitals that initiate the approach and fusion of the corresponding  
 413 nuclei. Analysis of the nuclear radioactive decay may require the discrete Kramers’ activation  
 414 mechanism (“roaming” over energy levels to reach a certain boundary) [25], which is  
 415 commonly used in the physicochemical kinetics. Here we imply the dynamics of energy  
 416 accumulation by an unstable *isu*-state nucleus on its “last” bond, the disruption of which  
 417 leads to the decay of the nucleus along a certain path.

418

#### 419 4. NUCLEAR CHEMICAL PROCESSES IN ANDREA ROSSI'S E-CAT REACTOR

420 The above concept of initiating low-energy nuclear-chemical reactions by the  
 421 mechanisms of nuclei fusion and  $e^{-}$ -catalysis can be used to physically interpret the results  
 422 of testing A. Rossi's energy E-Cat reactors as well [26]. Let us briefly discuss the results of  
 423 testing the E-Cat working chamber of the Rossi's reactor, presented by a group of  
 424 international experts [27]. The working chamber was a hollow ceramic cylinder 2 cm in  
 425 diameter and 20 cm long, into which the researchers loaded a fuel: about 0.9 g of finely  
 426 dispersed nickel powder with all stable isotopes present ( ${}_{28}^{58}\text{Ni}$ ,  ${}_{28}^{60}\text{Ni}$ ,  ${}_{28}^{61}\text{Ni}$ ,  ${}_{28}^{62}\text{Ni}$ , and  
 427  ${}_{28}^{64}\text{Ni}$  of 67, 26.3, 1.9, 3.9 and 1 %, respectively), and 0.1 g of  $\text{LiAlH}_4$  powder ( ${}_{3}^6\text{Li}$  and  ${}_{3}^7\text{Li}$   
 428 isotopes of 8.6 and 91.4%, respectively). The cylinder was sealed and then heated. The tests  
 429 were carried out for 32 days at chamber heating temperatures up to 1260 °C (first half of the  
 430 time) and 1400 °C (second half of the time). The energy released in the tests was measured  
 431 using the value of the heat flux produced by the chamber. In the tests, the overall excess  
 432 energy of 1.5 MWh was produced, corresponding to the chamber efficiency higher than 3.5.  
 433 The researchers recorded changes in the isotopic composition of the main fuel components  
 434 (nickel, lithium), for which the initial composition of stable elements was close to the  
 435 tabulated natural composition. After the tests, the isotopic composition of the recorded  
 436 components was dramatically changed: almost all nickel powder, more than 98%, was a









437 nickel-62 isotope (about 4% initially); the fraction of lithium-7 dropped to about 8% and  
 438 lithium-6 jumped to about 92%. The isotope abundances of the initial fuel and final “ash” in  
 439 the tests are listed in Table 1 [27].  
 440

441 Table 1. Isotope abundances for the initial fuel and final ash in the tests [27]

Ion	Fuel		Ash		Natural abundance [%]
	Counts in peak	Measured abundance [%]	Counts in peak	Measured abundance [%]	
${}^6\text{Li}^+$	15804	8.6	569302	92.1	7.5
${}^7\text{Li}^+$	168919	91.4	48687	7.9	92.5
${}^{58}\text{Ni}^+$	93392	67	1128	0.8	68.1
${}^{60}\text{Ni}^+$	36690	26.3	635	0.5	26.2
${}^{61}\text{Ni}^+$	2606	1.9	~0	0	1.8
${}^{62}\text{Ni}^+$	5379	3.9	133272	98.7	3.6
${}^{64}\text{Ni}^+$	1331	1	~0	0	0.9

442  
 443

444 According to the above concept, the recorded change in the isotopic composition of  
 445 main fuel components, nickel and lithium, in the presence of the hydrogen given off in the  
 446 decomposition of  $\text{LiAlH}_4$  at the above temperatures may be caused by the formation of a  
 447 protium-containing plasma in the reaction volume and the occurrence of neutral metastable  
 448 nuclei  ${}^1n_{isu}$ . Like neutrons, these neutral nuclei can interact with the nuclei of elements  
 449 constituting the fuel, accounting for the changes occurring in its elemental and isotopic  
 450 composition, which is accompanied by the corresponding energy release:



459

460 The above list of reactions implies that the specific (per component unit mass) energy release  
 461 is the highest for the lithium-7 nuclei. At the same time, when the mass fraction of the  
 462 lithium-7 isotope in the system is low, the total contribution to the heat release of the nuclear  
 463 reactions of  ${}^1n_{isu}$  nuclei with all other fuel elements, such as aluminum and nickel isotopes,  
 464 can become dominating. The almost complete disappearance of isotopes  ${}^7_3\text{Li}$  and  ${}^{58}_{28}\text{Ni}$  in the  
 465 ashes, which was recorded after the chamber was tested for more than a month, implies that  
 466 the values of rate constants are rather high not only for the processes (24) and (27), but also  
 467 for the other nuclear processes in which the new chemical elements are formed.  
 468

469 To understand the specific mechanisms accounting for the major changes in the fuel  
 470 composition during the E-Cat operation, including the almost complete exhaustion of the  
 471 lithium-7 isotope and the dominant growth of the nickel-62 isotope in the ash, it is necessary  
 472 to consider the other nuclear reactions, which can also change the isotopic composition of the

473 initial nickel. In these reactions, the energy carried away by the formed neutrinos and  
474 antineutrinos can noticeably reduce their heat releases, as compared to the above reactions:

$$475 \quad {}_{28}^{58}\text{Ni} + {}^1n_{isu} \rightarrow {}_{28}^{59}\text{Ni} + \nu\bar{\nu} + Q(8.22\text{MeV}), \quad T_{1/2}({}_{28}^{59}\text{Ni}) = 7.6 \cdot 10^4 \text{ yr}, \quad (32)$$

$$476 \quad {}_{28}^{60}\text{Ni} + {}^1n_{isu} \rightarrow {}_{28}^{61}\text{Ni} + \nu\bar{\nu} + Q(7.04\text{MeV}), \quad (33)$$

$$477 \quad {}_{28}^{61}\text{Ni} + {}^1n_{isu} \rightarrow {}_{28}^{62}\text{Ni} + \nu\bar{\nu} + Q(9.81\text{MeV}), \quad (34)$$

$$478 \quad {}_{28}^{62}\text{Ni} + {}^1n_{isu} \rightarrow {}_{28}^{63}\text{Ni} + \nu\bar{\nu} + Q(6.05\text{MeV}), \quad T_{1/2}({}_{28}^{63}\text{Ni}) = 100.1 \text{ yr}, \quad (35)$$

$$479 \quad {}_{28}^{64}\text{Ni} + {}^1n_{isu} \rightarrow {}_{28}^{65}\text{Ni} + \nu\bar{\nu} + Q(5.32\text{MeV}), \quad T_{1/2}({}_{28}^{65}\text{Ni}) = 2.52 \text{ h}, \quad (36)$$

480

481 The long half-life of the  ${}_{28}^{59}\text{Ni}$  isotope practically excludes the process in which the  
482 other nickel isotopes decayed in the tests are “replenished” with the  ${}_{28}^{58}\text{Ni}$  isotope, whose  
483 fraction is twice the fractions of the other nickel isotopes. Therefore, the almost complete  
484 absence of the  ${}_{28}^{60}\text{Ni}$  isotope in the ash should be attributed to the processes (28) and (33). It  
485 can also be assumed that the processes (29) and (34) account for the disappearance of the  
486  ${}_{28}^{61}\text{Ni}$  isotope in the ash; meanwhile, the process (34) brings the isotope  ${}_{28}^{62}\text{Ni}$  to the ash,  
487 providing its prevailing abundance among the other nickel isotopes in the ash. The additional  
488 contribution to this prevailing abundance is made by the “low” value of rate constant for the  
489 decline of the  ${}_{28}^{62}\text{Ni}$  isotope in the reaction (30), which describes the formation of cobalt with  
490 a low energy release in the process. It is also important to note that the long half-life of the  
491  ${}_{28}^{63}\text{Ni}$  isotope practically prevents from increasing the abundance of the isotope  ${}_{28}^{64}\text{Ni}$  in the  
492 ash, and the process (31) provides an almost complete conversion of this isotope in the initial  
493 nickel to the copper-65 isotope.

494

495 Admittedly, the above arguments can only qualitatively explain the ash composition  
496 recorded in the test. In this case, of high interest could be a comparative study of the  
497 elemental and isotopic composition of the ash and initial fuel by inductively coupled plasma  
498 mass spectrometry [28], successfully used before for studying the isotopic composition of  
499 impurities in the nickel in the laser ablation of a nickel sample in water. Here, it is important  
500 to study the changes in isotope ratios for various elements in the ash and initial fuel, primarily  
501 for the base element (nickel), as well as for the elements formed in the processes (24) - (31),  
502 such as magnesium, silicon, manganese, iron, cobalt, and copper.

503

## 504 5. TIME VARIATION OF THE ${}^{234}\text{U}/{}^{238}\text{U}$ ACTIVITY RATIO IN GROUND-WATER 505 FLOW SYSTEMS

506 One of the manifestations of the above-described initiated  ${}^{238}\text{U}$  nucleus decays is the  
507 well-known time variation of the basic, close to unity, ratio of the activity levels of uranium-  
508 234 and uranium-238 involved in the same decay chain of radioactive transformations  
509 (uranium/radium series) in the surface ground waters of seismic and volcanic regions [29-31].  
510 The activity  $\eta_i$ , introduced as the decay rate of the  $i$ -th uranium isotope, is defined as  
511  $\eta_i = k_i N_i$ , where  $k_i$  and  $N_i$  are the decay rate constant and the number of  $i$ -th isotope nuclei  
512 to be decayed, respectively. Note that the fraction  $\theta_i$  of the uranium-234 isotope in natural  
513 uranium ores is as low as  $\theta_{234} \approx 0.0055\%$ , with a half-life  $T_{1/2}({}^{234}\text{U}) \approx 2.45 \cdot 10^5$  years. At the  
514 same time, the fraction of the uranium-238 isotope is  $\theta_{238} \approx 99.3\%$ ,

515 yielding  $\chi \equiv \theta_{234}/\theta_{238} \approx 5.54 \cdot 10^{-5}$ , with a much longer half-life  $T_{1/2}(^{238}\text{U}) \approx 4.47 \cdot 10^9$  years.

516 The corresponding decay rate constants are related by the formula  $k_{234} = k_{238} \cdot \chi^{-1}$ .

517 It means [30] that in undisturbed minerals older than several million years, the  
518 abundances of  $^{238}\text{U}$  and its intermediate  $\alpha$ -decay product,  $^{234}\text{U}$ , reach a state of secular  
519 equilibrium. Under these conditions, the activity ratio (AR),

520  $\eta_{234}/\eta_{238} = \frac{T_{1/2}(^{238}\text{U})\theta_{234}}{T_{1/2}(^{234}\text{U})\theta_{238}} \equiv ^{234}\text{U}/^{238}\text{U AR}$ , will equal unity. However, natural waters,

521 especially in seismically active regions, typically are enriched in  $^{234}\text{U}$  with  $^{234}\text{U}/^{238}\text{U AR}$   
522 between 1 and 10 [30]. The uranium concentrations and  $^{234}\text{U}/^{238}\text{U AR}$  ratios in saturated-  
523 zone and perched ground waters were used to study the hydrologic flow in the vicinity of  
524 Yucca Mountain [30]. The U data were obtained by thermal ionization mass spectrometry for  
525 more than 280 samples from the Death Valley regional flow system. Wide variations in both  
526 U concentrations (commonly 0.6–10  $\mu\text{g l}^{-1}$ ) and  $^{234}\text{U}/^{238}\text{U AR}$  (commonly 1.5–6) were  
527 observed on both local and regional scales. The ground water beneath the central part of  
528 Yucca Mountain had intermediate U concentrations but a distinctive  $^{234}\text{U}/^{238}\text{U AR}$  of about  
529 7–8. It is necessary to add that about 600 seismic events have occurred near the site in the last  
530 20 years alone, with a 5.6-magnitude earthquake that happened as recently as 1992. There is  
531 also an evidence of relatively recent volcanic activity in the area.

532 Similar results were reported elsewhere [31], where the measurements of the  
533  $^{234}\text{U}/^{238}\text{U AR}$  in groundwater samples were used for monitoring the current deformations in  
534 the active faults at the Kultuk polygon, West Shore of Lake Baikal, for earthquake prediction.  
535 It was observed that the  $^{234}\text{U}/^{238}\text{U AR}$  fluctuated in time, with the duration of cycles and  
536 amplitudes of  $^{234}\text{U}/^{238}\text{U AR}$  fluctuations were variable in the range of 1.5–3.3, and the cycles  
537 of  $^{234}\text{U}/^{238}\text{U AR}$  in water were synchronized in the lines of the monitoring stations in the  
538 sublatitudinal and submeridional direction at the time intervals when seismic shocks occurred  
539 at the Kultuk polygon. The U concentrations in the ground-water samples of the Kultuk  
540 polygon ranged from 0.0087 to 5 mcg/l. The basic scenario of  $^{234}\text{U}/^{238}\text{U AR}$  variations in  
541 groundwater, recorded in the Kultuk polygon during the monitoring session, was examined in  
542 connection with the seismogenic activation of the western end of the Obruchev fault.

543 It is commonly believed that  $^{234}\text{U}$  enters solutions preferentially as a result of several  
544 mechanisms related to its origin by radioactive decay of  $^{238}\text{U}$  [30]. These mechanisms  
545 include damage of crystal-lattice sites containing  $^{234}\text{U}$  and the preferential release of  $^{238}\text{U}$  not  
546 bound to the crystal lattice from the defects of minerals, as well as direct ejection of the recoil  
547 nucleus into the water near the boundaries of mineral grains.

548 At the same time, the results of [29–31] suggest that the mechanochemical processes  
549 in relatively small volumes of uranium ore in ore deposits located in the geologically active,  
550 including seismically and volcanically active, zones of the Earth's crust are the important  
551 factor that can account for the significant changes of  $^{234}\text{U}/^{238}\text{U AR}$  under study [31]. These  
552 zones can be characterized by the emergence of high mechanical stresses, initiated shifts in  
553 the ore, and the formation of cracks and fissures. These processes in the U ore at high local  
554 mechanical pressures can not only change the structure of groundwater flows in the zone, but  
555 also give rise to high local electric fields and initiate the decomposition of water molecules  
556 and the formation of high-energy (on chemical scales) electrons. In this case, the concept  
557 developed in this paper allows us to expect that the formation of cracks and fissures in a

558 uranium ore can initiate the radioactive decay of uranium-238 nuclei by the  $e^-$ -catalytic  
 559 mechanism, producing *isu*-state  $\beta$ -protactinium nuclei. Note that it is the phenomenon of  
 560 mechanically activated nuclear processes discovered in the works of Deryagin et al. [32, 33]  
 561 that can be regarded as the starting point in the new stage of studying LENRs, which is  
 562 usually attributed to the work of Fleischmann and Pons [34]. For instance, it was  
 563 experimentally recorded that the destruction of targets made of a heavy ( $D_2O$ ) ice by a metal  
 564 striker with an initial velocity of 100–200 m/s produces neutrons, and their number is several  
 565 times higher than the background level [32]. In contrast, no new neutrons were recorded  
 566 when the same action was applied to the target made of an ordinary ( $H_2O$ ) ice.

567 Assume that when a fissure is formed, a fraction of uranium atoms leaves the fissure  
 568 surface layer of the uranium ore and is dissolved in the aqueous phase, with each isotope  
 569 dissolved according to its abundance in the ore. Additionally, assume that a very small  
 570 fraction  $\xi$  ( $\xi \ll 1$ ) of  $N_{238}$  nuclei of the main uranium-238 isotope that pass to the aqueous  
 571 medium is activated in the fissure formation by the  $e^-$ -catalytic mechanism and converted to  
 572 *isu*-state  $\beta$ -protactinium nuclei. Without this activation, the activity level of  $N_{238}$  nuclei of the  
 573 atoms of uranium-238 isotope in the aqueous medium,  $\eta_{238} = k_{238} N_{238}$ , was equal to the  
 574 activity level  $\eta_{234} = k_{234} N_{234}$  for the  $N_{234}$  nuclei of uranium-234 isotope that passed to the  
 575 aqueous medium. Section 2 implies that in the initiated radioactive decay, the effective decay  
 576 rate constant of  $^{238}U$  nuclei,  $\tilde{k}_{238}$ , for a relatively small number  $\xi$  of  $N_{238}$  nuclei in the aqueous  
 577 medium can dramatically change. It is wise to use the above in considering the simplified  
 578 decay of uranium-238 and uranium-234 isotopes. In this case, the decay of “intermediate”  
 579 thorium-234 and protactinium-234m isotopes with short lifetimes, which are also involved in  
 580 the radioactive uranium/radium series, is taken out of consideration. The balance equations  
 581 for the numbers of  $N_{238}$  and  $N_{234}$  nuclei at a steady-state concentration of uranium-234 isotope  
 582 in the aqueous medium can be written as  
 583

$$584 \quad \frac{dN_{238}}{dt} = -k_{238}(1-\xi)N_{238} - \xi\tilde{k}_{238}N_{238} = -k_{238}^{eff}N_{238}, \quad (37)$$

$$585 \quad \frac{dN_{234}}{dt} = -k_{234}N_{234} + k_{238}(1-\xi)N_{238} + \xi\tilde{k}_{238}N_{238} = 0. \quad (38)$$

586

587 Here,

588

$$589 \quad k_{238}^{eff} = k_{238} \left[ 1 + \xi \left( \frac{\tilde{k}_{238}}{k_{238}} - 1 \right) \right] \quad (39)$$

590

591 is the effective rate constant for the decay of the uranium-238 isotope when the radioactive  
 592 decay of the fraction  $\xi$  of uranium-238 nuclei is initiated by external factors and  
 593 characterized by the decay rate constant  $\tilde{k}_{238}$ . Equations (37)-(38) yield the desired formula  
 594 for the ratio of activity levels of uranium-234 and uranium-238 isotopes in open systems in  
 595 which the initiated accelerated decay of uranium-238 is effectuated:  
 596

$$597 \quad \eta_{234}/\eta_{238} = \frac{k_{234} N_{234}}{k_{238} N_{238}} = {}^{234}U/{}^{238}U \text{ AR} = 1 + \xi \left( \frac{\tilde{k}_{238}}{k_{238}} - 1 \right). \quad (40)$$

598 In this case, the apparent higher activity level of the uranium-234 isotope cannot be  
 599 attributed to the fact that the groundwater is directly enriched with  $^{234}\text{U}$  nuclides because its  
 600 release to the aquatic medium is easier due to the decay of the main  $^{238}\text{U}$  isotope, as is  
 601 usually assumed [30, 31]. The increased content of  $^{234}\text{U}$  nuclei in the aqueous medium is a  
 602 result of the decay of  $^{238}\text{U}$  nuclei initiated by the formation of cracks and fissures, which  
 603 produces  $\beta$ -protactinium nuclei by the  $e^-$ -catalytic mechanism; their release to the aqueous  
 604 medium, and their subsequent decay along the chain of the radioactive uranium/radium  
 605 series. The reference value of  $\tilde{k}_{238}/k_{238} \sim 10^9$  estimated in [4], showing a possible increase by  
 606 9 orders of magnitude of the decay rate constant for the  $^{238}\text{U}$  nuclei in the laser ablation,  
 607 implies that for the ratios  $^{234}\text{U}/^{238}\text{U AR} \sim 5-10$ , characteristic for the system studied in [30],  
 608 to take place the fraction  $\xi$  of activated  $^{238}_{91}\text{Pa}_{isu}$  nuclei relative to  $^{238}\text{U}$  nuclei in the aqueous  
 609 media must be  $\sim (0.5-1) \cdot 10^{-8}$ .

610

## 611 6. NON-BARYONIC STATES OF NUCLEAR MATTER AND “HEAVY” QUARKS

612

613 It is well known that quarks as subunits of hadrons manifest themselves as free point  
 614 objects in the energy and momentum transfer in the proton collisions occurring in the  
 615 colliding beams with characteristic energies of more than 1 TeV for each pair of the colliding  
 616 nucleons [35, 36]. As a result, the quarks can be associated with the independent degrees of  
 617 freedom of nuclear matter. When the decays of excited hadrons that were formed in these  
 618 high-energy collisions of particles are considered, the quarks are traditionally regarded as  
 619 *elementary particles* with “point” electric charges of  $-1/3 e$  or  $+2/3 e$ , where  $e$  is the absolute  
 620 value of the electron charge. It is as such particles that the quarks are involved in the Standard  
 621 Model of Elementary Particles [36].

622

623 In this section, the concept stating that non-nucleonic *isu*-states may occur in the  
 624 nuclear matter will be used in considering a set of problems arising in the study of decays of  
 625 the excited baryons and mesons that were formed in the high-energy collisions of protons and  
 626 characterized by highly excited states of “decay”. Their characteristic half-lives are quite  
 627 long,  $\sim 10^{-13}-10^{-8}$  s, which implies the dominant role of weak nuclear interactions in these  
 628 decays. These times are higher than the characteristic nuclear times by 10 or more orders of  
 629 magnitude. According to the Standard Model [36, 37], in addition to  $u$ - and  $d$ -quarks, which  
 630 are characterized by the so-called current quark masses of 2.3 and 4.8  $\text{MeV}/c^2$ , respectively,  
 631 the excited hadrons contain heavier  $s$ -,  $c$ -, and  $b$ -quarks with current quark masses of 95,  
 632 1275, and 4180  $\text{MeV}/c^2$ , respectively [37]. When there is, at least, one heavy quark among the  
 633 three quarks of a baryon, the baryon is called heavy. The mesons formed by a quark-  
 antiquark pair are called heavy when the quark is heavy.

634

635 When the decay of heavy hadrons is discussed, an accent is usually made on purely  
 636 formal aspects related to the classification of quarks in the Standard Model of Elementary  
 637 Particles [36], which is based on the requirements of symmetry for the wave functions of  
 638 baryons as fermions and mesons as bosons with regard for the quantum numbers additively  
 639 introduced for heavy quarks. This allows one to predict possible decay paths for the heavy  
 640 hadrons to be formed, using a given set of their quantum numbers. However, the nature of the  
 641 introduced new quantum numbers, defined as  $s$  (strange),  $c$  (charmed), and  $b$  (beauty or  
 642 bottom), remains unclear. It is not clear what kind of physically interpreted parameters can  
 643 account for the above differences in the masses of heavy quarks. In addition, it is unclear how  
 644 each of the heavy quarks is converted to light  $u$ - or  $d$ -quarks in the decay of heavy hadrons,  
 645 because the hadron-products formed in the high-energy collisions of protons or nuclei contain  
 only light  $u$ - or  $d$ -quarks. There is no discussion yet about the nature of the confinement of

646 quarks, defined as the impossibility of separating quarks from the nuclear matter and studying  
 647 them in a free state. Instead, an assumption is made that the force of mutual attraction of  
 648 quarks rises as the distance between them increases, without any discussion about the nature  
 649 of this force.

650 It is suggested in this study that the above problems can be studied in terms of the  
 651 phenomenological approach (qualitatively rather than quantitatively) if it is assumed that  
 652 quarks are not elementary particles but kinetic markers, three for baryons and two for  
 653 mesons, for the large fragments of the nuclear matter, which are bound to each other by the  
 654 strong nuclear interaction effectuated by the exchange of pions. It is these interactions that  
 655 account for the confinement of quarks as quasi-particles. In addition, assume that the highly  
 656 excited hadrons, such as charged baryons  $p^*$ , neutral baryons  $n^*$ , and mesons  $m^*$ , which are  
 657 formed in the collisions of high-energy particles, lose their stability, which is provided by the  
 658 exchange of pions, because of the relative high-energy movements of the current quarks, and  
 659 come to an *isu*-state of decay. The subsequent relaxation of the *isu*-state of such excited  
 660 hadrons and the formation of decay products, such as nucleons, pions, and leptons, are  
 661 initiated by the weak nuclear interaction between the quarks found in the hadrons using the  
 662 formation and absorption of gauge vector bosons.

663 Let us consider a working hypothesis, assuming that highly excited heavy baryons  
 664 and mesons are particles with local soliton-like excited states of nuclear matter that can be  
 665 based only on  $u$ - and  $d$ -quarks, the corresponding  $\tilde{u}$ - and  $\tilde{d}$ -antiquarks, and virtual pions  
 666 and vector bosons. We will try to understand whether it is possible to hypothetically identify  
 667 the degrees of freedom, which can be defined as heavy quarks, in the excited system. Here, it  
 668 is implied that the polarization of the nuclear medium in the vicinity of its quarks could  
 669 effectively cause an increase in the current masses of the  $u$ - and  $d$ -quarks, converting them to  
 670 heavy quarks.

671 To date, the researchers have discovered many types of decays of excited baryons,  
 672 including those producing both neutral and charged particles at intermediate stages, which  
 673 leads to a wide variety of final neutral and charged particles, such as nucleons, pions, and  
 674 leptons [36, 37]. Consider several examples of the decay of excited nucleons  $n^*$  and  $p^*$ ,  
 675 parenthesizing the hyperon (heavy baryon) that decayed to form the products of interest. Let  
 676 it be hyperons  $\Lambda(uds)$ ,  $\Lambda_c^+(udc)$ ,  $\Sigma_c^0(ddc)$  and  $\Lambda_b^0(udb)$ , containing  $s$ -,  $c$ -, и  $b$ -quarks. These  
 677 quarks are symbolically shown in the parenthesized quark composition of the hyperons. The  
 678 processes below show the intermediate and final decay products for these hyperons, with two  
 679 possible decay paths for the hyperon  $\Lambda$ :

680  
 681

$$682 \quad n^*(\Lambda) \rightarrow p + \pi^-, \quad (41)$$

$$683 \quad n^*(\Lambda) \rightarrow n + \pi^0, \quad (42)$$

$$684 \quad p^*(\Lambda_c^+) \rightarrow p + K^- + \pi^+ \rightarrow p + \mu^- + \tilde{\nu}_\mu + \pi^+, \quad (43)$$

$$685 \quad n^*(\Sigma_c^0) \rightarrow \Lambda_c^+ + \pi^- \rightarrow p + \mu^- + \tilde{\nu}_\mu + \pi^+ + \pi^-, \quad (44)$$

$$686 \quad n^*(\Lambda_b^0) \rightarrow \Lambda_c^+ + \pi^+ + \pi^- + \pi^- \rightarrow p + \mu^- + \tilde{\nu}_\mu + \pi^+ + \pi^+ + \pi^- + \pi^-. \quad (45)$$


687  
 688

689 Feynman diagrams representing the above processes, which present the complicated  
 690 dynamics of  $u$ - and  $d$ -quark conversions in the formation and decay of charged and neutral  
 691 virtual vector bosons, are plotted in Fig. 2.



692



693  
694



695  
696



697  
698

699 Fig. 2. Feynman diagrams for the decay of hyperons that contain (a, b) s-quark, (c, d) c-  
700 quark, and (e) b-quark  
701

702 It is obvious that the number of pions and muons formed along with a nucleon in the  
703 decay of an excited baryon depends on the total excitation energy of the baryon. As the birth  
704 of, at least, one virtual vector boson is needed for one pion or muon to be formed in the final  
705 state (Fig. 2 a, b, c), one could expect that the processes in which the number of formed pions  
706 and muons in the final state increases would be strongly suppressed because the  
707 dimensionless constant  $\alpha_F$  of weak nuclear interaction is low. Note, however, that when the  
708 processes in Fig. 2 are analyzed quantitatively, it should be taken into account that the weak  
709 nuclear interactions are not as weak as it is often assumed. The value of the corresponding  
710 dimensionless constant  $\alpha_F$  is almost an order of magnitude greater than the value of fine-  
711 structure constant  $\alpha_e$  [18, 38]. Indeed, if the dimensionless constant of strong nuclear  
712 interaction is taken to be  $\alpha_s = \sqrt{2}$  [18] and the value of squared elementary charge of weak

713 nuclear interaction is estimated by  $q_F^2 \equiv G_F/a_Z^2$  [36], where  $a_Z = 2^{1/2}\hbar/m_Zc \approx 3.3 \cdot 10^{-16}$  cm  
 714 is the characteristic radius related to the mass of the intermediate  $Z^0$  vector boson  $m_Z = 91.2$   
 715  $\text{GeV}/c^2 = 1.62 \cdot 10^{-22}$  g and  $G_F = 1.17 \cdot 10^{-5}(\hbar c)^3/(\text{GeV})^2$  is the Fermi constant of four-  
 716 fermion interaction, we obtain  $\alpha_F = \frac{q_F^2}{\hbar c} \approx 4.9 \cdot 10^{-2}$  and, hence,  $\alpha_F/\alpha_s \approx 3.45 \cdot 10^{-2}$ . In this  
 717 case,  $\alpha_e = 1/137 \approx 0.73 \cdot 10^{-2}$  and, hence,  $\alpha_e/\alpha_s \approx 5.2 \cdot 10^{-3}$  and  $\alpha_F/\alpha_e = 6.7$ .

718 Unfortunately, the proton mass is often used in the literature [35] as the normalizing  
 719 mass in estimating the dimensionless constant of weak nuclear interaction, though it is almost  
 720 100 times smaller than the mass of a  $Z^0$  vector boson [36]. As a result, the value of constant  
 721  $\alpha_F$  is underestimated by almost 4 orders of magnitude. According to our above estimates, the  
 722 correct value of this constant is only 35 times, rather than 5 orders of magnitude, less than the  
 723 value for the dimensionless constant of strong nuclear interaction. It is this correct value that  
 724 accounts for the existence of reliable experimental data on the decays of highly excited heavy  
 725 baryons that produce five or more pions and a muon [39].

726 As noted above,  $u$ - and  $d$ -quarks can be regarded as kinetic markers for the subunits of  
 727 nuclear matter. The diagrams plotted in Fig. 2 imply that every decay process given above is  
 728 a complicated process with respect to the  $u$  and  $d$  dynamic variables, in which the nonlinear  
 729 interrelations with respect to these variables represent the processes of energy redistribution  
 730 in the transfer of charged and neutral vector bosons and account for the emergence of pions  
 731 as decay products.

732 The diagrams demonstrate both the variety of possible baryon decays for the same  
 733 excited state (Fig. 2 *a, b*) and the fact that it is *impossible to introduce*, in addition to  $u$  and  $d$ ,  
 734 new types of degrees of freedom – heavy quarks as effective dynamical variables for  
 735 describing the entire diversity of complicated decay dynamics of excited baryons  $p^*$  and  $n^*$ .  
 736 The latter follows from the comparison of Figs. 2 *c* and Figs. 2 *d* showing the decays of  
 737  $\Lambda_c^+(udc)$  and  $\Sigma_c^0(ddc)$  hyperons, respectively, which contain, as supposed, the same heavy  $c$ -  
 738 quark, but are accomplished by different mechanisms with 2 and 3 virtual vector bosons  
 739 formed, respectively.

740 The number of examples of this kind that illustrate different decay mechanisms for  
 741 hadrons, which according to the modern theory [35, 36] contain light quarks together with  
 742 one of the heavy quarks ( $s$ ,  $c$  or  $b$ ), could be larger if we consider the decays of heavy  
 743 baryons and mesons. It will be wise, however, to use the general phenomenological approach  
 744 when only  $u$ - и  $d$ -quarks are considered as the current degrees of freedom, which are related  
 745 to certain fragments of nuclear matter and change their electric charge in the absorption and  
 746 emission of charged vector bosons.

747 Concluding this section, it should be noted that, proton-proton collisions with an  
 748 energy of 7-8 TeV produce particles with a four-quark [40] and five-quark arrangement of  
 749 nuclear matter [41]. The lifetimes of these particles are  $\sim 10^{-24}$ - $10^{-23}$  s, which is typical for  
 750 the resonances. This means that the nuclear exchange forces in the excited systems of 4 and 5  
 751 quarks are too weak to keep these systems so long as it is necessary for their relaxation to be  
 752 accounted for by the weak nuclear interaction, which is true for the systems of 2 and 3  
 753 quarks. Experiments of this kind could not record the formation of a weakly decaying strange  
 754 heavy dibaryon either [42], in contrast to the low-energy experiments, which produce a  $\beta$ -  
 755 dineutron.

756  
 757

758 7. METASTABLE NON-NUCLEONIC STATES OF NUCLEAR MATTER IN DOUBLE  
759 BETA-DECAYS

760

761

762

763

764

765

766

767

768

769

770

771

772

773

774

775

776

777

778

779

780

781

782

783

784

785

786

The double  $\beta$ -decays of some even-even nuclei [43-46] could be a future candidate for one of the manifestations of the initiation of metastable states in the nuclear matter, in which the nuclear forces are not strong enough to bind a part of the quarks into stable nucleons and the nucleonic structure in the nucleus is locally shaken up. All cases in which this type of decay is reliably recorded are characterized by half-lives longer than  $10^{18}$  years, which is several orders of magnitude greater than the existence time of the Universe. The main difficulties to overcome in studying the double  $\beta$ -decay are represented by its low probability and the long-run experiments needed to minimize background events as possible, as well as by close and thorough analysis of experimental results. So far, these decays are experimentally recorded in 10 out of more than 30 pairs of even-even isotopes that can be bound by the double  $\beta^-$ -decay. At the same time, in about the same number of pairs of even-even isotopes that can be bound by the double  $\beta^+$ -decay, no decays of this type have not been recorded yet. The latter can be a result of a noticeable difference in the probabilities of  $\beta^-$ - and  $\beta^+$ -decay. The double  $e^-$ -capture was recorded for only one nucleus:  $^{130}\text{Ba}$  isotope.

The above phenomenological concept of the radioactive decay of nuclei initiated by the  $e^-$ -catalysis mechanism [5, 7] allows us not only to understand the possible reason for the difference in the probabilities  $\beta^-$ - and  $\beta^+$ -decay, but also open new opportunities in studying double  $\beta^-$ - and  $\beta^+$ -decays because their rates dramatically rise when the processes are initiated using low-energy excitations. The latter is implied not only by the general result of [5, 7] regarding the loss of stability of *isu*-state nuclei in the nuclear matter, but also by the experimental data recorded in a group of studies, for example, [9, 10], dealing with the  $\alpha$ -decay of all so-called stable tungsten isotopes (half-lives  $\sim 10^{17}$ - $10^{19}$  years) initiated in the glow discharge.

As a future experiment idea, it is of interest to compare the characteristic parameters  $Q$  and  $|\Delta Q|$  for already and not yet recorded double  $\beta^-$ -decays of different isotopes, including those of the same element:

787 
$$^{48}_{20}\text{Ca} + e^-_{he} \rightarrow ^{48}_{19}\text{K}_{isu} + \nu \rightarrow ^{48}_{22}\text{Ti} + 3e^- + 2\tilde{\nu} + \nu + Q(4.27\text{MeV}), \quad \Delta Q = -12.09\text{MeV}, \quad (46)$$

788 
$$^{46}_{20}\text{Ca} + e^-_{he} \rightarrow ^{46}_{19}\text{K}_{isu} + \nu \rightarrow ^{46}_{22}\text{Ti} + 3e^- + 2\tilde{\nu} + \nu + Q(0.98\text{MeV}), \quad \Delta Q = -7.72\text{MeV}, \quad (47)$$

789 
$$^{82}_{34}\text{Se} + e^-_{he} \rightarrow ^{82}_{33}\text{As}_{isu} + \nu \rightarrow ^{82}_{36}\text{Kr} + 3e^- + 2\tilde{\nu} + \nu + Q(3.01\text{MeV}), \quad \Delta Q = -7.27\text{MeV}, \quad (48)$$

790 
$$^{80}_{34}\text{Se} + e^-_{he} \rightarrow ^{80}_{33}\text{As}_{isu} + \nu \rightarrow ^{80}_{36}\text{Kr} + 3e^- + 2\tilde{\nu} + \nu + Q(0.136\text{MeV}), \quad \Delta Q = -5.64\text{MeV}, \quad (49)$$

791 
$$^{116}_{48}\text{Cd} + e^-_{he} \rightarrow ^{116}_{47}\text{Ag}_{isu} + \nu \rightarrow ^{116}_{50}\text{Sn} + 3e^- + 2\tilde{\nu} + \nu + Q(2.81\text{MeV}), \quad \Delta Q = -6.15\text{MeV}, \quad (50)$$

792 
$$^{114}_{48}\text{Cd} + e^-_{he} \rightarrow ^{114}_{47}\text{Ag}_{isu} + \nu \rightarrow ^{114}_{50}\text{Sn} + 3e^- + 2\tilde{\nu} + \nu + Q(0.54\text{MeV}), \quad \Delta Q = -5.08\text{MeV}, \quad (51)$$

793 
$$^{186}_{74}\text{W} + e^-_{he} \rightarrow ^{186}_{73}\text{Ta}_{isu} + \nu \rightarrow ^{186}_{76}\text{Os} + 3e^- + 2\tilde{\nu} + \nu + Q(0.49\text{MeV}), \quad \Delta Q = -3.9\text{MeV}. \quad (52)$$

794

795

796

797

798

The value of released energy  $Q$  accounts for the phase volume of the reaction products to be formed because the process probability is proportional to this factor, whereas the value of deficit  $|\Delta Q|$  for the structural energy of the *isu*-state nucleus accounts for the extent to which the stability of the nucleus is lost.

799 Comparison of the values of  $|\Delta Q|$  and  $Q$  for the experimentally studied processes of  
 800 double  $\beta^-$ -decay of calcium-48, selenium-82, and cadmium-116 isotopes with those for the  
 801 processes not studied yet due to, as we can suggest, their much lower probability gives us a  
 802 reason to consider the released energy  $Q$  as the main parameter accounting for the double  
 803  $\beta^-$ -decay – the processes with lower energy releases. This suggestion is supported by the  
 804 data in Table 2 (was compiled on the basis of [43, 44]) which lists the values of  $Q$  in the other  
 805 7 processes for which the  $\beta^-$ -decay was experimentally recorded. Note that the smallest  
 806 value of parameter  $Q$  refers to the double  $\beta^-$ -decay of uranium-238 nuclei, and this value, as  
 807 we can suggest, may account for the highest value of its half-life, which is 3-4 orders of  
 808 magnitude higher the values for the other 9 nuclei.

809  
 810 Table 2. Isotopes with experimentally recorded double  $\beta^-$ -decay

Isotope	$Q$ , MeV	$ \Delta Q $ , MeV	$T_{1/2}$ , years
$^{48}_{20}\text{Ca}$	4.27	12.09	$(4.3 \pm 2.3) \times 10^{19}$
$^{76}_{32}\text{Ge}$	2.04	7.01	$(1.3 \pm 0.4) \times 10^{21}$
$^{82}_{34}\text{Se}$	3.01	7.27	$(9.2 \pm 0.8) \times 10^{19}$
$^{96}_{40}\text{Zr}$	3.35	7.1	$(2.0 \pm 0.4) \times 10^{19}$
$^{100}_{42}\text{Mo}$	3.03	6.24	$(7.0 \pm 0.4) \times 10^{18}$
$^{116}_{48}\text{Cd}$	2.81	6.15	$(3.0 \pm 0.3) \times 10^{19}$
$^{128}_{52}\text{Te}$	0.87	4.38	$(3.5 \pm 2.0) \times 10^{24}$
$^{130}_{52}\text{Te}$	2.53	4.96	$(6.1 \pm 4.8) \times 10^{20}$
$^{150}_{60}\text{Nd}$	3.37	5.69	$(7.9 \pm 0.7) \times 10^{18}$
$^{238}_{92}\text{U}$	1.11	3.46	$(2.0 \pm 0.6) \times 10^{21}$

811  
 812 Below we list several double  $\beta^+$ -decays of even-even isotopes among which there  
 813 may be processes with the values of  $Q$  commensurate with those at which the double  $\beta^-$ -  
 814 decay is effectuated:

815  
 816  $^{106}_{48}\text{Cd} + e^-_{he} \rightarrow ^{106}_{47}\text{Ag}_{isu} + \nu \rightarrow ^{106}_{46}\text{Sn} + 2e^+ + e^- + 3\nu + Q(2.78\text{MeV}), \quad \Delta Q = -0.094\text{MeV}, \quad (53)$

817  $^{108}_{48}\text{Cd} + e^-_{he} \rightarrow ^{108}_{47}\text{Ag}_{isu} + \nu \rightarrow ^{108}_{46}\text{Sn} + 2e^+ + e^- + 3\nu + Q(0.27\text{MeV}), \quad \Delta Q = -1.65\text{MeV}, \quad (54)$

818  $^{112}_{50}\text{Sn} + e^-_{he} \rightarrow ^{112}_{49}\text{In}_{isu} + \nu \rightarrow ^{112}_{48}\text{Cd} + 2e^+ + e^- + 3\nu + Q(1.92\text{MeV}), \quad \Delta Q = -0.66\text{MeV}, \quad (55)$

819  $^{152}_{64}\text{Gd} + e^-_{he} \rightarrow ^{152}_{63}\text{Eu}_{isu} + \nu \rightarrow ^{152}_{62}\text{Sm} + 2e^+ + e^- + 3\nu + Q(0.058\text{MeV}), \quad \Delta Q = -1.82\text{MeV}. \quad (56)$

820  
 821 In the above processes, the possible candidate isotopes are cadmium-106 and tin-112.  
 822 At the same time, the comparison between the  $2\beta^+$ - and  $2\beta^-$ -decays shows that their values  
 823 of  $|\Delta Q|$  are much different from each other (see the Table 3 compiled on the basis of [43,  
 824 45]). It is the low values of  $|\Delta Q|$  that may account for the low probabilities of  $2\beta^+$ -decays,  
 825 because any radioactive decay of nuclei begins with the initiated (either due to fluctuation or  
 826 by the action of external factors) interaction of an electron in the inner shells of the


827 radioactive atom with its nucleus  ${}^A_Z N$ , which produces an intermediate *isu*-state nucleus  
 828  ${}_{Z-1}^A M_{isu}$  [5, 7]. As the rates at the stage where the  ${}_{Z-1}^A M_{isu}$  nucleus is formed are much higher  
 829 when the process is initiated in a low-temperature plasma, the possibility of recording  $2\beta^+$ -  
 830 decays may be higher for the isotopes characterized by the highest values of  $|\Delta Q|$  and  $Q$ .  
 831 Probable candidate nuclei for these decays among all  $2\beta^+$ -active nuclei are listed in Table 2,  
 832 where the barium-132, xenon-126, and tellurium-120 isotopes may be expected to become  
 833 the most promising candidates for experimental studies of  $2\beta^+$ -decay.

834 The above difference in the values of  $|\Delta Q|$  for the  $2\beta^-$ - and  $2\beta^+$ -decays is  
 835 reasonable to attribute to the initiating role of the electron factor in the radioactive decays of  
 836 nuclei [5] occurring in our asymmetric Universe, which is characterized by the recorded  
 837 existence of the matter with atoms composed of elementary particles and the absence of the  
 838 antimatter with atoms composed of antiparticles. This difference in probability of the  $2\beta^-$ -  
 839 and  $2\beta^+$ -decays can be considered as one of the arguments in favor of the hypothesis of the  
 840 activating role of electrons in radioactive decays. Indeed, the nature of these differences for  
 841 radioactive decays cannot be understood in terms of generally accepted approaches.  
 842

843 Table 3. Possible double  $\beta^+$ -decay candidates for even-even isotopes


Isotope	$Q$ , MeV	$ \Delta Q $ , MeV
${}^{96}_{44}\text{Ru}$	2.72	0.254
${}^{106}_{48}\text{Cd}$	2.78	0.194
${}^{112}_{50}\text{Sn}$	1.92	0.664
${}^{120}_{52}\text{Te}$	1.70	0.982
${}^{124}_{54}\text{Xe}$	3.07	0.295
${}^{126}_{54}\text{Xe}$	0.90	1.258
${}^{130}_{56}\text{Ba}$	2.58	0.368
${}^{132}_{56}\text{Ba}$	0.833	1.28
${}^{138}_{64}\text{Ce}$	2.01	0.433
${}^{156}_{58}\text{Dy}$	0.708	1.045
${}^{162}_{68}\text{Er}$	1.85	0.296

844 We suggest that the above double  $\beta$ -decays must be accompanied by the coupled  
 845 conversion of quarks of two neutrons in the  $\beta^-$ -decays or two protons for the  $\beta^+$ -decays in  
 846 the emission and absorption of vector bosons. In these processes, the initiating role is played  
 847 by the quasi-free (within the nucleus) quarks emerged in the formation of the *isu*-state of  
 848 nuclear matter. The corresponding Feynman diagrams, which schematically represent the  
 849 dynamics of the interactions effectuated in double  $\beta$ -decays, are plotted in Fig. 3.  
 850  
 851




a

852  
853  
854



855



b

c

856  
857  
858

859 Fig. 3. Feynman diagrams for initiated (a)  $2\beta^-$  - and (b, c)  $2\beta^+$  -decays

860

861

862

863

864

865

866

867

868

869

870

871

872

873

874

875

876

877

Note that when double  $\beta$ -decay diagrams are discussed, the dynamics of coupled conversions are taken into account only when the decays without neutrinos, which are possible when the neutrino and antineutrino are the same particle (Majorana neutrino), are considered [46]. As we can expect that the characteristic times of  $2\beta^-$ -decays initiated by low-energy actions will be many orders of magnitude less than the values usually recorded in experiments, there is a hope to see in the future not only new experimental data on double  $\beta^-$ -decays, but also the first data on  $\beta^+$ -decays, as well as to get some idea about the extension to which the lepton number conservation law in the  $2\beta^-$ -decays is violated.

## 8. NATURE OF THE ANOMALY RECORDED IN THE BERYLLIUM-8 DECAY

Let us discuss the anomalies in the angular correlations between the positrons and electrons emitted in the radioactive decays of excited  ${}^8_4\text{Be}^*$  nuclei [3]. As in [5], we assume that the decay of a  ${}^8_4\text{Be}^*$  nucleus is preceded by its interaction with one of the electrons in the inner electron shells of the atom, which emits a neutrino  $\nu$  and produces an excited metastable  $isu$ -state  ${}^8_3\text{Li}^*_{isu}$  nucleus:

$$878 \quad {}^7_3\text{Li} + p \rightarrow {}^8_4\text{Be}^*, \quad {}^8_4\text{Be}^* + e^-_{he} \rightarrow {}^8_3\text{Li}_{isu}^* + \nu \rightarrow 2{}^4_2\text{He} + 2e^- + e^+ + \nu + \tilde{\nu}. \quad (57)$$

879

880 Based on the above decay diagram for the excited  ${}^8_4\text{Be}^*$  nucleus, we can suggest a new  
 881 formation mechanism for a correlated  $e^+e^-$  pair in the reaction  ${}^7\text{Li}(p, \gamma){}^8\text{Be}$  when two states  
 882 of the  ${}^8_4\text{Be}^*$  nucleus, 17.64 MeV and 18.15 MeV, are excited, as in the experiment [3], which  
 883 is alternative to the one proposed in [1-3].

884 As noted above, the nature of the excitation of  ${}^8_4\text{Be}$  nuclei initiated by the collisions  
 885 with nucleons, when the entire nucleonic subsystem of the nucleus is excited, is substantially  
 886 different from the nature of the local metastable *isu*-excitation caused by a shake-up in the  
 887 nucleonic structure of  ${}^8_3\text{Li}_{isu}$  nucleus and the loss of the overall stability of these  $\beta$ -nuclei.  
 888 The latter depends on the absolute value of structural energy deficit  $\Delta Q = (m_{{}^8_4\text{Be}} - m_{{}^8_3\text{Li}})c^2$  and  
 889 the difference in the masses of  ${}^8_4\text{Be}$  and  ${}^8_3\text{Li}$  nuclei in the ground state. Therefore, we can  
 890 assume that the excitation energy of the nucleonic subsystem of the  ${}^8_4\text{Be}^*$  nucleus can be  
 891 almost completely kept in the nucleonic subsystem of the  ${}^8_3\text{Li}_{isu}^*$  nucleus if the latter has the  
 892 corresponding excited state. In this case, it becomes obvious that the efficiency of the decay  
 893 of the excited  ${}^8_3\text{Li}_{isu}^*$  nucleus that emits two alpha particles and a correlated  $e^+e^-$  pair will  
 894 depend on how close one of the excited states of the  ${}^8_3\text{Li}$  nucleus approximates the excited  
 895 state of the  ${}^8_4\text{Be}$  nucleus [47].

896


897 Here we must take into account that the probability of emitting  $\gamma$  quanta by excited  
 898 nuclei, which depends on the width of the corresponding excited state, and the probability of  
 899 emitting photons in the transition of a single atom from the excited state to the ground state  
 900 are initiated by the zero-point oscillations of the EM vacuum [48]. Virtually, the main factor  
 901 is the average of squared fluctuating values of the electric field intensity for the EM vacuum.  
 902 As noted above, when a metastable *isu*-state with a local shake-up in the nucleonic structure  
 903 is initiated in the nuclear matter, the irreversible loss of the nucleus stability is likewise  
 904 accounted for by the EM vacuum as a result of changes in the boundary conditions at the  
 905 nucleus surface [5, 18]. However, these two emissions are independent of each other.

906 As the ground-state energy for the  ${}^8_3\text{Li}$  nucleus is 16.005 MeV higher than that for the  
 907  ${}^8_4\text{Be}$  nucleus [47], the excited states of 1.635 and 2.145 MeV for the  ${}^8_3\text{Li}$  nucleus could  
 908 formally correspond to the excited states of 17.64 and 18.15 MeV for the  ${}^8_4\text{Be}$  nucleus. For  
 909 the  ${}^8_3\text{Li}$  nucleus, the excited states closest to the ground one are 0.891 MeV, which is not  
 910 high enough for producing an  $e^-e^+$  pair, and 2.255 MeV, which is 0.11 MeV higher than the  
 911 above value of 2.145 MeV. If the anomaly in angular correlations between positrons and  
 912 electrons recorded in [3] is effectuated by the above excited states of  ${}^8_4\text{Be}$  and  ${}^8_3\text{Li}_{isu}$  nuclei, it  
 913 implies that in this case the width of the 2.255 MeV state for the  ${}^8_3\text{Li}_{isu}$  nucleus is larger than  
 914 0.11 MeV, and we can speak about a direct correspondence between the 18.15 MeV excited  
 915 state for the  ${}^8_4\text{Be}$  nucleus and the 2.255 MeV excited state for the  ${}^8_3\text{Li}_{isu}$  nucleus. Obviously,  
 916 this correspondence needed for the anomaly recorded in [3] to take place may be achieved by  
 917 adjusting the kinetic energy  $E_p$  of the bombarding protons, though not always.


918

919 Assume that the anomaly recorded in [3], which is the formation of correlated  
 920  $e^-e^+$  pairs at their opening angles  $\Theta \sim 130-140^\circ$ , is mostly due to the exchange of  $d$ - and  $u$ -  
 921 quarks localized in the region of non-nucleonic metastability of the  $isu$ -state nucleus, which  
 922 can migrate over the nucleus, and  $d$ - and  $u$ -quarks of the superpositions  $d\tilde{d}$  and  $u\tilde{u}$  among  
 923 the quark-antiquarks pairs produced in the decay of vector  $Z^0$ -mesons in the same  $isu$ -region  
 924 of the nuclear matter. Virtually, this exchange is effectuated in the annihilation of these  
 925 quarks and antiquarks of  $d\tilde{d}$  and  $u\tilde{u}$  pairs producing a correlated  $e^-e^+$  pair, which is in good  
 926 agreement with the decay of a neutral boson studied in [1-3].  
 927

928




929  
930



a

b



c

931  
932

933 Fig. 4. Feynman diagrams for the formation of correlated  $e^-e^+$ ,  $e^-\tilde{\nu}$  и  $e^+\nu$  pairs initiated by  
 934 the interaction between a  $\pi^0$ -meson (quark-antiquark superposition  $d\tilde{d}$  and  $u\tilde{u}$ ) and the  
 935 nucleons of the excited  ${}^8\text{Be}^*$  nucleus  
 936

937 Possible diagrams for the formation of  $e^-e^+$  pairs accompanied by the production of  
 938  $e^-\tilde{\nu}$  and  $e^+\nu$  pairs are plotted in Fig. 4. In contrast to the decays of bosons as particles with a  
 939 certain set of quantum characteristics analyzed in [1-3], the quark pairs to be annihilated in  
 940 these diagrams can have different relative orbital moments, as in [49], which does not impose  
 941 any substantial restrictions on the sets of quantum numbers for the excited states of the  
 942 nuclei; specifically,  ${}^8_4\text{Be}^*$  and  ${}^8_3\text{Li}_{isu}^*$ . For this reason, the results of [3] were discussed above  
 943 without referring to the quantum numbers for the excited states of these nuclei.



944 The developed concept stating the existence of the metastable states of the nuclear  
 945 matter with a local shake-up of its nucleonic structure makes it possible to qualitatively  
 946 interpret the formation of correlated  $e^-e^+$  pairs in the experiment under discussion [3] without  
 947 involving the hypothesis of a fifth fundamental interaction into the physical science.  
 948 Admittedly, it is possible so far to speak only about a qualitative understanding of the  
 949 correlation of  $e^-e^+$  pairs in the above process because the exchanges of quarks in the region  
 950 of nucleus non-nucleonic metastability have not been studied yet.

951 An additional clarity in discussing the above alternative could be brought by the new  
 952 experiments proposed in [2] to record the anomalies in the angular correlations between  
 953 positrons and electrons, like those in [3], that are emitted in the radioactive decay of other  
 954 excited nuclei. The study [2] deals with the reactions  ${}^7\text{Li}({}^3\text{He}, \gamma){}^{10}\text{B}^*$  (19.3 MeV) [50] and  
 955  ${}^7\text{Li}(t, \gamma){}^{10}\text{Be}^*$  (17.79 MeV) [51] and assumes that the decay of these excited states of the  
 956 daughter nuclei can produce  $e^+e^-$ -pairs with the same type of opening-angle anomaly as in  
 957 [3].

958 The developed concept of the radioactive decays of the excited nuclei producing the  
 959  ${}^{10}\text{B}^*$  (19.3 MeV) nucleus implies that, in the first of the above reactions, the  ${}^{10}_4\text{Be}_{isu}^*$  nucleus  
 960 rather than the  ${}^{10}_5\text{B}^*$  nucleus would decay, producing the final products  ${}^7\text{Li}$ ,  ${}^3\text{He}$ , and an  $e^+e^-$ -  
 961 pair. In the second reaction, the final products  ${}^7\text{Li}$ ,  ${}^3\text{H}$ , and an  $e^+e^-$ -pair are formed in the  
 962 decay of the  ${}^{10}_3\text{Li}_{isu}^*$  nucleus rather than the  ${}^{10}_4\text{Be}^*$  nucleus. The above differences are  
 963 significant due to the high difference in the ground-state energies of the nuclei to be decayed  
 964 when these energies are referred to the unified energy scale. It is this kind of analysis that will  
 965 enable us to make an unambiguous choice in conducting appropriate experimental studies in  
 966 favor of the hypothesis of the existence of a fifth fundamental interaction or developed  
 967 concept of nuclear radioactive decays. The most significant differences are seen for the  
 968 energy levels of  ${}^{10}_3\text{Li}_{isu}^*$  and  ${}^{10}_4\text{Be}^*$  nuclei: the ground state for the lithium-10 nucleus is 20.444  
 969 MeV higher than the one for the beryllium-10 nucleus. The corresponding difference between  
 970 the  ${}^{10}_4\text{Be}_{isu}^*$  nucleus and  ${}^{10}_5\text{B}^*$  nucleus is 0.556 MeV [47].

971 In view of the above differences, the excited state of 19.3 MeV for the boron-10  
 972 nucleus should formally be in correspondence with the excited state of 18.74 MeV for the  
 973 beryllium-10 nucleus, in which the excited-state energy closest to the latter value is 18.55  
 974 MeV. When the width of this state is greater than 0.2 MeV, the above correspondence can be  
 975 true and the anomaly in the angular correlations between positrons and electrons emitted in  
 976 the radioactive decays of excited  ${}^{10}\text{B}^*$  nuclei can, in principle, be recorded. The situation is  
 977 substantially different when we look for these anomalies in the decays of excited  ${}^{10}\text{Be}^*$   
 978 nuclei. The excited state of 17.79 MeV for the beryllium-10 nucleus cannot even formally be  
 979 in correspondence with the ground state for the lithium-10 nucleus because the energy  
 980 difference between the ground states for the lithium-10 nucleus and beryllium-10 nucleus is  
 981 higher than the above excited-state energy for the beryllium-10 nucleus. Therefore, the  
 982 desired correlations in the  ${}^7\text{Li}(t, \gamma){}^{10}\text{Be}^*$  reaction should be sought at kinetic energies  $E_t$  of the  
 983 tritium nuclei higher than those suggested in [2]. For example, the excited states of 1.4, 2.35,  
 984 and 2.85 MeV for the lithium nucleus, whose decay into  ${}^7\text{Li}$  and  ${}^3\text{H}$  may be accompanied by  
 985 the anomaly in the angular correlations of the recorded  $e^+e^-$ -pair can be achieved using the  
 986 tritium nucleus energies of 21.8, 22.8, and 23.3 MeV, respectively. This experiment can  
 987 become an *experimentum crucis* in selecting between the discussed nature alternatives for the  
 988 correlated opening angle of the  $e^+e^-$ -pair, as well as in deciding whether it is possible to

989 initiate metastable states with a shaken-up nucleonic structure in the nuclear matter and,  
990 hence, validate the new concept of radioactive decays of nuclei.

991

## 992 9. CONCLUSION

993

994 This study may be the first attempt to discuss the existence of metastable states of the  
995 nuclear matter when the nuclear forces are not strong enough to bind a part of the quarks into  
996 stable nucleons, which results in a local nucleonic structure shake-up in the nucleus. With this  
997 anomalous excited state of the nuclear matter, called inner shake-up or *isu*-state, the  
998 relaxation of the nuclei is initiated by the weak nuclear interaction. Apparently, the most  
999 unexpected result of this study is represented by the fact that we discovered the unified  
1000 physical nature of decays in the nuclear matter under the action of weak nuclear forces. These  
1001 decays can be initiated by both a low-temperature plasma and the collision of countermoving  
1002 proton beams with characteristic energies higher than 1 TeV per colliding proton pair. In  
1003 either way of initiation, the necessary condition for this type of decay – large enough drop in  
1004 the strong nuclear interaction, is met. In the first way, this is achieved by the “soft force”: by  
1005 initiating an inelastic interaction between the hot (on chemical scales) electron and the  
1006 nucleus denying the *K*-trapping, which produces a certain mass deficit in the resulting  
1007 nucleus. In the second way, it is achieved by a direct high-energy action: by increasing the  
1008 kinetic energy of the baryon quarks.

1009 It is the above approach that we successfully used before to understand a large enough  
1010 set of experimental data on the initiation of low energy nuclear reactions and acceleration of  
1011 radioactive  $\alpha$ - and  $\beta$ -decays in a low-temperature plasma. Taking the metastable non-  
1012 nucleonic states of the nuclear matter into account in the study of the above high-energy  
1013 collisions made it possible to understand the nature of various recorded decays of highly  
1014 excited hadrons, which are effectuated by the weak nuclear interaction.

1015 The concept stating the existence of metastable states of the nuclear matter, in which  
1016 the nucleonic structure is locally shaken up, enabled us to present the arguments  
1017 substantiating an alternative approach to interpreting the experimental results [1-3] and  
1018 question the need to introduce a fifth fundamental interaction into the physical science,  
1019 additional to the electromagnetic, nuclear strong/weak, and gravitational interactions, which  
1020 can relate ordinary matter and hypothetical dark matter.

1021

1022

## 1023 REFERENCES

1024

1025 1. Feng J.L., Fornal B., Galon I., et al., Protophobic fifth force interpretation of the observed  
1026 anomaly in  $8\text{Be}$  nuclear transitions, *Phys. Rev. Lett.* 117 (2016), no. 7, 071803;  
1027 arXiv:1604.07411v2 [hep-ph].

1028 2. Feng J.L., Fornal B., Galon I., et al., Particle Physics Models for the 17 MeV Anomaly in  
1029 Beryllium Nuclear Decays; arXiv:1608.03591v2 [hep-ph].

1030 3. Krasznahorkay A.J., Csatlós M., Csige L., et al., “Observation of anomalous internal pair  
1031 creation in  $\text{Be}8$  : a possible indication of a light, neutral boson,” *Phys. Rev. Lett.* 116 (2016)  
1032 042501; arXiv:1504.01527 [nucl-ex].

1033 4. Timashev S. F., Simakin A. V., Shafeev G. A. Nuclear–Chemical processes under the  
1034 conditions of laser ablation of metals in aqueous media (Problems of “cold fusion”) //  
1035 *Russian Journal of Physical Chemistry A.* 2014. V. 88. N. 11. P. 1980-1988).

1036 5. Timashev S.F. Radioactive decay as a forced nuclear chemical process: phenomenology //  
1037 *Russian Journal of Physical Chemistry A.* 2015. V. 89. N. 11. P. 2072–2083.

- 1038 6. Timashev Serge. Nuclear-chemical processes in the solar atmosphere // International  
1039 Journal of Astrophysics and Space Science, 2014. V. 2(6). P. 88-92.
- 1040 7. Timashev S.F.. Initiating nuclear-chemical transformations in native systems:  
1041 phenomenology // Russian Journal of Physical Chemistry A. 2016. V. 90. N. 10. P. 2089-  
1042 2095.
- 1043 8. Vysotskii V.I., Kornilova A.A. Microbial transmutation of Cs-137 and LENR in growing  
1044 biological systems // Curr. Sci. 2015. V. 108. P. 636-640.
- 1045 9. Savvatimova I. Creation of more light elements in tungsten irradiated by low-energy  
1046 deuterium ions. Proc.13th Int. Conf.ICCF13, Sochy, Russia, 2007. P. 505-517.
- 1047 10. Savvatimova I., Savvatimov G., Kornilova A. Decay in tungsten irradiated by low energy  
1048 deuterium ions, Proc.13th Int. Conf.ICCF13, Sochy, Russia, 2007
- 1049 11. Shafeev G.A., Simak A.V., Bozon-Verduraz F., Robert M. Excitation of high energy  
1050 levels under laser exposure of suspensions of nanoparticles in liquids // Applied Surface  
1051 Science. 2007. V. 254 P. 1022-1026.
- 1052 12. Simak A.V., Shafeev G.A. Initiation of nuclear reactions under laser irradiation of  
1053 metal nanoparticles in the presence of thorium aqua ions // Physics of Wave Phenomena.  
1054 2008. V. 16. N. 4. P. 268–274.
- 1055 13. Barmina E.V., Timashev S.F., Shafeev G.A. Laser-induced synthesis and decay of  
1056 Tritium under exposure of solid targets in heavy water // Journal of Physics: Conference  
1057 Series. 2016. V. 688. 012106. (8th International Conference on Inertial Fusion Sciences and  
1058 Applications (IFSA 2013) IOP Publishing; <https://arXiv.org/abs/1306.0830>.
- 1059 14. Jung M., Bosch F., Beckert K. et al. First observation of bound-state decay // Phys. Rev.  
1060 Lett. 1992. V. 69. 2164-2167.
- 1061 15. Bosch F., Faestermann T., Friese J. et al. Observation of bound-state  $\beta^-$  decay of fully  
1062 ionized  $^{187}\text{Re}$ :  $^{187}\text{Re} - ^{187}\text{Os}$  cosmochronometry // Phys. Rev. Lett. 1996. V. 77. 5190-5193.
- 1063 16. Thomas S.A., Abdalla F.D., Lahav O. Upper Bound of 0.28 eV on Neutrino Masses from  
1064 the Largest Photometric Redshift Survey // Phys. Rev. Lett. 2010. V. 105. N 3. P. 031301
- 1065 17. Lattimer J.M., Pethick C.J., Prakash M., Haensel P. Direct URCA process in neutron stars  
1066 // Phys. Rev. Lett. 1991. V. 66. P. 2701-2704
- 1067 18. Timashev Serge F. The Planck numbers and the essence of gravity: phenomenology //  
1068 <http://arxiv.org/abs/1701.08073> [physics.gen-ph].
- 1069 19. V. M. Sharapov and S. L. Kanashenko, Vopr. At. Nauki Tekh., Ser. Termoyad. Sintez //  
1070 2008. No. 2, P. 20 [in Russian].
- 1071 20. Kervran C.L. Biological Transmutation. Happiness Press. USA. Magalia, California.  
1072 1998.
- 1073 21. Biberian J.-P. Review Article Biological Transmutations: Historical Perspective // J.  
1074 Condensed Matter Nucl. Sci. 2012. V. 7. P. 11–25.
- 1075 22. Vysotskii V.I., Kornilova A.A. Nuclear Transmutation of Stable And Radioactive  
1076 Isotopes In Biological Systems. Pentagon Press. New Delhi. 2010.
- 1077 23. Gromov A.A., Gromov A.M., Popenko E.M. et al. Formation of calcium in the products  
1078 of iron–aluminum thermite combustion in air // Russian Journal of Physical Chemistry A.  
1079 2016. V. 90 (10). P. 2104-2106.
- 1080 24. Borisov A.A., De Luca L.T., Merzhanov A.G. Self-Propagating High-Temperature  
1081 Synthesis of Materials. CRC Press. Taylor and Francis. 2000, 400 p.
- 1082 25. Tunitskii N. N., Kaminskii V. A., Timashev S. F. Methods of Physicochemical Kinetics.  
1083 Moscow: Khimiya. 1972, 198 p. [in Russian].
- 1084 26. United State Patent – US 9,115,913 B1, Aug. 25. 2015;  
1085 <https://animpossibleinvention.files.wordpress.com/2015/08/us9115913b1.pdf>; Andrea Rossi
- 1086 27. Levi G., Foschi E., Höistad B., et al., Observation of Abundant Heat Production from a  
1087 Reactor Device and of Isotopic Changes in the Fuel.

- 1088 <http://amsacta.unibo.it/4084/1/LuganoReportSubmit.pdf>.
- 1089 28. Revel'skiy I.A., Buryak A.K., Sajti P. L., et al. Isotopic ratio changes of several trace  
1090 elements in nickel as a result of laser ablation in aqueous medium, in press.
- 1091 29. Finkel R.C., 1981. Uranium concentrations and  $^{234}\text{U}/^{238}\text{U}$  activity ratios in fault-  
1092 associated groundwater as possible earthquake precursors // *Geophysical Research Letters*.  
1093 1981. V. 8(5). P. 453–456.
- 1094 30. Paces J.B., Ludwig K.R., Peterman Z.E., Neymark L.A.  $^{234}\text{U}/^{238}\text{U}$  evidence for local  
1095 recharge and patterns of ground-water flow in the vicinity of Yucca Mountain, Nevada, USA  
1096 // *Applied Geochemistry*. 2002. V. 17. Issue 6. P. 751–779.
- 1097 31. Rasskazov S.V., Chebykin E.P., Ilyasova A.M. et al. Creating the Kultuk polygon for  
1098 earthquake prediction: variations of ( $^{234}\text{U}/^{238}\text{U}$ ) and  $^{87}\text{Sr}/^{86}\text{Sr}$  in groundwater from active  
1099 faults at the western shore of lake Baikal // *Geodynamics and Tectonophysics* (Published by  
1100 the Institute of the Earth's crust Siberian Branch of Russian Academy of Sciences). 2015.  
1101 V.6. Issue 4. P. 519-553.
- 1102 32. Derjagin B.V., Klyuev V.A., Lipson A.G., Toporov Yu.P. Possibility of nuclear reactions  
1103 during the fracture of solids // *Colloid Journal USSR*. 1986. V. 48 (1). P. 8-10.
- 1104 33. Tsarev V.A. Cold fusion // *Soviet Physics Uspekhi*. 1990. V. 33(11). P. 881-910.
- 1105 34. Fleishmann M., Pons S. and Hawkins M. Electrochemically induced nuclear fusion of  
1106 deuterium // *J. Electroanal. Chem*. 1989. V. 261. P. 301-308.
- 1107 35. Klapdor-Kleingrothaus H.V., Zuber K. *Teilchenastrophysik*. B.G. Teubner GmbH,  
1108 Stuttgart, 1997.
- 1109 36. Donnelly T.W., Formaggio J.A., Holstein B.R., Milner R.G., B. Surrow B. *Foundations*  
1110 *of Nuclear and Particle Physics*. Cambridge. University Press. 2017, 745 p.
- 1111 37. Khoze V.A., Shifman M.A. Heavy quarks // *Sov. Phys. Usp*. 1983. V. 26. P. 387–424.
- 1112 38. Akhiezer A.I., Rekalov M.P. *Elementary particles*. Moscow: Nauka. 1986. 256 p. (in  
1113 Russian)
- 1114 39. Amsler C. et al. (Particle Data Group), PL B667, 1 (2008) (URL: <http://pdg.lbl.gov>)
- 1115 40. Aaij R. et al (LHCb collaboration). Observation of the resonant character of the  $Z(4430)$ -  
1116 state; <https://arxiv.org/abs/1404.1903>.
- 1117 41. Aaij R. et al (LHCb collaboration). Observation of  $J/\psi p$  resonances consistent with  
1118 pentaquark states in  $\Lambda_b^0 \rightarrow J/\psi K^- p$  decays // <https://arXiv.org/abs/1507.03414v2> .
- 1119 42. ALICE Collaboration. Search for weakly decaying dibaryon states //  
1120 <https://arxiv.org/abs/1506.07499>.
- 1121 43. Lazarenko V.R. Double beta decay and the properties of the neutrino // *Physics-Uspekhi*.  
1122 1967. V.9.N.6. P. 860–873.
- 1123 44. Pritychenko B. On Double-Beta Decay Half-Life Time Systematics // Brookhaven  
1124 National Laboratory. BNL-91299-2010. April 14, 2010; <https://arxiv.org/abs/1004.3280>.
- 1125 45. Audi G., Bersillon O., Blachot J., Wapstra A.H. The NUBASE evaluation of nuclear and  
1126 decay properties // *Nuclear Physics A*. 2004. V. 729. Issue 1. P. 3-128.
- 1127 46. Zelevinsky V., Volya A. *Physics of Atomic Nuclei*. Wiley-VCH Verlag GmbH &  
1128 Co.KgaA. 2017, 667 p.
- 1129 47. D.R. Tilley et al. Energy levels of light nuclei  $A = 8, 9, 10$  // *Nuclear Physics A*. 2004. V.  
1130 745. Issue 3-4. P. 155-362.
- 1131 48. Vihman E. *Berkeley Physics: Volume 4: Quantum Physics*. McGraw-Hill Book Co.  
1132 1967.
- 1133 49. Shapiro I.S., Timashev S.F. Direct reactions with two nucleon transfer // *Nucl. Phys*.  
1134 1965. V. 79. P. 46-64.
- 1135 50. S. C. Ling and S. L. Blatt, “States in  $^{10}\text{B}$  between 18 and 22 MeV,” *Nucl. Phys. A*174  
1136 (1971) 375–384.

1137 51. K.M. Subotić, B. Lalović, B.Z. Stepančić. The  ${}^7\text{Li}({}^3\text{H}, \gamma){}^{10}\text{Be}$  reaction from 0.4–1.1 MeV  
1138 // Nucl. Phys. A296 (1978) 141–150.  
1139